

The completion problem for N -matrices *

CLAUDIA MENDES and JUAN R. TORREGROSA

Dpto. de Matematica \ Dpto. de Matemática Aplicada
Universidade do Minho \ Universidad Politécnica de Valencia
Campus de Gualtar, 4700 320 Braga \ Camino de Vera, 46071-Valencia
PORTUGAL \ SPAIN
clmendes@math.uminho.pt \ jrtorre@mat.upv.es

Abstract

An $n \times n$ matrix is called an N -matrix if all principal minors are negative. In this paper, we are interested in N -matrix completions problems, that is, when a partial N -matrix has an N -matrix completion. In general, a combinatorially or non-combinatorially symmetric partial N -matrix does not have an N -matrix completion. Here, we prove that a combinatorially symmetric partial N -matrix has an N -matrix completion if the graph of its specified entries is a 1-chordal graph. We also prove that there exists an N -matrix completion for a partial N -matrix whose associated graph is an undirected cycle.

Key-Words: Partial matrix, completion, combinatorially symmetric, N -matrix, 1-chordal graph, cycles.

1 Introduction

A *partial matrix* is a matrix in which some entries are specified and others are not. In this work we consider partial matrices where the diagonal entries are known. A *completion* of a partial matrix is the matrix resulting from a particular choice of values for the unspecified entries. A *completion problem* asks if we can obtain a completion of a partial matrix with some prescribed properties. A partial ma-

trix $A = (a_{ij})$ is said to be *combinatorially symmetric* when a_{ij} is specified if and only if a_{ji} is.

A natural way to describe an $n \times n$ partial matrix $A = (a_{ij})$ is via a graph $G_A = (V, E)$, where the set of vertices V is $\{1, 2, \dots, n\}$, and the edge or arc $\{i, j\}$, ($i \neq j$) is in set E if and only if position (i, j) is specified; as all main diagonal entries are specified, we omit loops. In general, a directed graph is associated with a non-combinatorially symmetric partial matrix and, when the partial matrix is combinatorially symmetric, an undirected graph can be used.

A *path* is a sequence of edges (arcs) $\{i_1, i_2\}$, $\{i_2, i_3\}$, \dots , $\{i_{k-1}, i_k\}$ in which the vertices are distinct. A *cycle* is a path with the first vertex equal to the last vertex. An undirected graph is *chordal* if it has no induced cycles of length 4 or more [2].

An $n \times n$ real matrix $A = (a_{ij})$ is called *N -matrix* if all its principal minors are negative. The principal submatrix of A lying in rows and columns α , $\alpha \subseteq N = \{1, 2, \dots, n\}$, is denoted by $A[\alpha]$.

N -matrices arise in the theory of global univalence of functions [3], in multivariate analysis [6], and in linear complementary problems [5, 7]. In [8], the class of N -matrices were studied in connection with Lemke's algorithm for solving linear and convex quadratic programming problems.

In the following proposition we give some important properties for N -matrices.

Proposition 1.1 *Let $A = (a_{ij})$ be an N -matrix of size $n \times n$. Then*

*Research supported by Spanish DGI grant number BFM2001-0081-C03-02

1. If P is a permutation matrix then PAP^T is an N -matrix.
2. If D is a positive diagonal matrix then DA and AD are N -matrices.
3. If D is a diagonal matrix then DAD^{-1} is an N -matrix.
4. $a_{ij} \neq 0$ and $\text{sign}(a_{ij}) = \text{sign}(a_{ji})$, $\forall i, j \in \{1, 2, \dots, n\}$.
5. $\forall \alpha \subset \{1, 2, \dots, n\}$, principal submatrix $A[\alpha]$ is an N -matrix.

From before properties, we can suppose, without lost of generality, that if A is an N -matrix of size $n \times n$, A is an element of set:

$$S_n = \{A = (a_{ij}) : a_{ij} \neq 0 \text{ and } \text{sign}(a_{ij}) = (-1)^{i+j+1}, \forall i, j\}$$

On the other hand, the last property of the before proposition allows us to give the following definition.

Definition 1.1 *A partial matrix is said to be a partial N -matrix if every completely specified principal submatrix is an N -matrix.*

The goal of this paper is the following N -matrix completion problem:

Problem 1 *Let A be a partial N -matrix.*

- (1.a) *Is there an N -matrix completion A_c of A ?*
- (1.b) *What conditions allow us to assure the existence of an N -matrix completion A_c of A ?*

In section 2 we analyze the above problem (1.a) for combinatorially and non-combinatorially symmetric partial N -matrices. In section 3 and 4 we study some types of undirected graphs whose the associated partial matrices have N -completions.

2 N -matrix completion problem

Let $A = (a_{ij})$ be a partial N -matrix of size $n \times n$. From property 4 of Proposition 1.1 the conditions

- (i) Specified entries of A are nonzero,
- (ii) $\text{sign}(a_{ij}) = \text{sign}(a_{ji})$, when a_{ij} and a_{ji} are specified,

are necessary conditions in order to obtain an N -matrix completion of A .

For matrices of size 2×2 conditions (i) and (ii) are also sufficient.

Proposition 2.1 *Let A be a partial N -matrix of size 2×2 . There exists an N -matrix completion A_c of A , if and only if A satisfies conditions (i) and (ii).*

Unluckily, the above proposition is false for partial matrices of size $n \times n$, $n \geq 3$, both when the partial matrix is combinatorially symmetric and it is not, as the following examples show:

- (a) The non-combinatorially symmetric partial N -matrix

$$A = \begin{bmatrix} -1 & 2 & x_{13} \\ 2 & -1 & 2 \\ 3 & 2 & -1 \end{bmatrix}$$

satisfies conditions (i) and (ii), but does not have an N -matrix completion since

$$\det A[\{1, 3\}] < 0 \Leftrightarrow 1 - 3x_{13} < 0 \Leftrightarrow x > 1/3,$$

and

$$\det A < 0 \Leftrightarrow 7x_{13} + 19 < 0 \Leftrightarrow x < -19/7.$$

- (b) The combinatorially symmetric partial N -matrix

$$A = \begin{bmatrix} -1 & 1 & x_{13} & -3 \\ 2 & -1 & 1 & x_{24} \\ x_{31} & 2 & -1 & 1 \\ -4 & x_{42} & 2 & -1 \end{bmatrix}$$

satisfies conditions (i) and (ii), but does not have an N -matrix completion since

$$\det A[\{2, 3, 4\}] < 0 \Leftrightarrow 3 + x_{24}x_{42} + 4x_{24} + x_{42} < 0 \\ \Rightarrow x_{24}, x_{42} < 0,$$

but

$$\det A[\{1, 2, 4\}] = 13 + x_{24}x_{42} - 4x_{24} - 6x_{42} > 0,$$

for $x_{24}, x_{42} < 0$.

If we add another condition to before conditions (i) and (ii) we can define the following set:

$$\begin{aligned} PS_n &= \{A = (a_{ij}), n \times n \text{ partial matrix} : \\ &\text{for } a_{ij} \text{ specified} \\ &a_{ij} \neq 0 \text{ and } \text{sign}(a_{ij}) = (-1)^{i+j+1}, \forall i, j\} \end{aligned}$$

Proposition 2.2 *Let A be a partial N -matrix of size 3×3 such that $A \in PS_3$. Then, there exists an N -matrix completion A_c of A .*

Corollary 2.1 *Let A be a combinatorially symmetric partial N -matrix of size 3×3 . Then, there exists an N -matrix completion A_c of A .*

Proposition 2.2 is not true for matrices of size $n \times n$, $n \geq 4$, as the following example shows.

Example 1

Consider the partial matrix

$$A = \begin{bmatrix} -1 & 1 & -11 & x_{14} \\ 2 & -1 & 1 & -200 \\ -0.1 & 10 & -1 & 1 \\ 1 & -10 & 1.01 & -1 \end{bmatrix}.$$

It is not difficult to verify that A is a partial N -matrix and $A \in PS_4$. However, A has no N -matrix completion because

$$\det A[\{1, 2, 4\}] = 1801 - 19x_{14} < 0 \Leftrightarrow x_{14} > 94.79,$$

and

$$\det A[\{1, 3, 4\}] = -9.89 + 0.899x_{14} < 0 \Leftrightarrow x_{14} < 11.$$

From this example, we can establish the following result:

Proposition 2.3 *For every $n \geq 4$, there is an $n \times n$ partial N -matrix, belong to PS_n , that has no N -matrix completion.*

Proof: We denote by \bar{I} the partial matrix, of size $(n-4) \times (n-4)$, with all entries unspecified except the entries of the main diagonal that are equal to -1. The partial matrix

$$B = \begin{bmatrix} A & X \\ Y & \bar{I} \end{bmatrix},$$

where X, Y are completely unspecified matrices and A is the matrix of the before example, is a partial N -matrix in PS_n that does not have N -matrix completion. \square

3 Chordal graphs

In order to get started, we recall some very rich clique structure of chordal graphs. See [2] for further information. A *clique* in an undirected graph G is simply a complete (all possible edges) induced subgraph. We also use clique to refer to a complete graph and use K_p to indicate a clique on p vertices. A useful view of chordal graphs is that they have a tree-like structure in which their maximal cliques play the role of vertices.

If G_1 is the clique K_q and G_2 is any chordal graph containing the clique K_p , $p < q$, then the *clique sum* (see [2]) of G_1 and G_2 along K_p is also chordal. The cliques that are used (to build chordal graphs) are the maximal cliques (see [2]) of the resulting chordal graph and the cliques along which the summing takes place are the so-called *minimal vertex separators* of the resulting chordal graph. If the maximum number of vertices in a minimal vertex separator is p , then the chordal graph is called *p -chordal*. In this section we are interested in 1-chordal graphs.

Proposition 3.1 *Let $A = (a_{ij})$ be a partial N -matrix of size $n \times n$, the graph of whose specified entries is 1-chordal with two maximal cliques, one of them with two vertices. Then there exists an N -matrix completion of A .*

Proof: We may assume, without loss of generality,

that A has the following form:

$$A = \begin{bmatrix} -1 & 1 & \cdots & x_{1n} \\ a_{21} & -1 & \cdots & (-1)^{n+1}a_{2n} \\ x_{31} & a_{32} & \cdots & (-1)^{n+2}a_{3n} \\ \vdots & \vdots & & \vdots \\ x_{n1} & (-1)^{n+1}a_{n2} & \cdots & -1 \end{bmatrix},$$

that can be partitioned as follows:

$$A = \begin{bmatrix} -1 & 1 & X \\ a_{21} & -1 & \bar{a}_{23}^T \\ Y & \bar{a}_{32} & A_{33} \end{bmatrix}.$$

It is easy to see that we obtain an N -matrix completion of A by replacing the unspecified entries in the following way:

$$\begin{aligned} x_{1j} &= -a_{2j}, & j \in \{3, 4, \dots, n\} \\ x_{i1} &= -a_{i2}, & i \in \{3, 4, \dots, n\} \end{aligned}$$

Proposition 3.2 *Let A be a partial N -matrix of size $n \times n$, the graph of whose specified entries is 1-chordal with two maximal cliques. Then there exists an N -matrix completion of A .*

Proof: We may assume, without loss of generality, that A has the following form:

$$A = \begin{bmatrix} A_{11} & a_{12} & X \\ a_{21}^T & -1 & a_{23}^T \\ Y & a_{32} & A_{33} \end{bmatrix}.$$

Consider the completion,

$$A_c = \begin{bmatrix} A_{11} & a_{12} & -a_{12}a_{23}^T \\ a_{21}^T & -1 & a_{23}^T \\ -a_{32}a_{21}^T & a_{32} & A_{33} \end{bmatrix}.$$

We are going to see that A_c is an N -matrix. Let α and β be the subsets of $N = \{1, 2, \dots, n\}$ such that

$$A_c[\alpha] = \begin{bmatrix} A_{11} & a_{12} \\ a_{21}^T & -1 \end{bmatrix}, \text{ and } A_c[\beta] = \begin{bmatrix} -1 & a_{23}^T \\ a_{32} & A_{33} \end{bmatrix},$$

and assume $|\alpha| = k$ (thus k is the index of the overlapping entry). Let $\gamma \subseteq N$. Then there are two cases to consider:

(a) $k \in \gamma$, then

$$\det A_c[\gamma] = (-1) \det A_c[\gamma \cap \alpha] \cdot \det A_c[\gamma \cap \beta] < 0$$

(b) $k \notin \gamma$. We consider

$$\gamma = \{1, 2, \dots, k-1, k+1, \dots, n\}.$$

For another γ we proceed in analogous way. By applying Jacobi's identity,

$$\det A_c[\gamma] = \det A_c^{-1}[\{k\}] \cdot \det A_c.$$

By case (a), $\det A_c < 0$, and we prove that $\det A_c^{-1}[\{k\}]$ is positive. \square

We can extend this result in the following way:

Theorem 3.1 *Let G be an undirected connected 1-chordal graph. Then any partial N -matrix, the graph of whose specified entries is G , has an N -matrix completion.*

Proof: The proof is by induction on the number p , of maximal cliques in G . The case of p -maximal cliques is reduced to that of $(p-1)$ -cliques by choosing a clique (the p th-clique) to be one that has only one vertex in common with any other maximal clique (the existence of such cliques follows from the way chordal graphs are built, see [2]). Then completing the subgraph induced by the remaining $(p-1)$ -cliques reduces the problem to the case of two maximal cliques. The case of a 1-chordal graph with two maximal cliques is handled in the before proposition. \square

4 Paths and cycles

In this section we are going to prove the existence of an N -completion for a partial N -matrix, combinatorially symmetric whose associated graph is a path or a cycle.

Proposition 4.1 *Let $A = (a_{ij})$ be an $n \times n$ combinatorially symmetric partial N -matrix, such that its associated graph is a path. Then, there exists an N -matrix completion.*

Proof: We can suppose, without loss of generality, that matrix A has the following form:

$$A = \begin{bmatrix} -1 & 1 & x_{13} & \cdots & x_{1n-1} & x_{1n} \\ a_{21} & -1 & 1 & \cdots & x_{2n-1} & x_{2n} \\ x_{31} & a_{32} & -1 & \cdots & x_{3n-1} & x_{3n} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ x_{n-11} & x_{n-12} & x_{n-13} & \cdots & -1 & 1 \\ x_{n1} & x_{n2} & x_{n3} & \cdots & a_{nn-1} & -1 \end{bmatrix},$$

with $a_{i+1i} > 0$, $i = 1, 2, \dots, n-1$.

It is easy to see that we obtain an N -matrix completion by replacing the unspecified entries in the following way:

$$\begin{aligned} x_{ij} &= (-1)^{i+j+1}, & i \in \{1, 2, \dots, n\}, \\ & & j \geq i+1 \\ x_{j+2j} &= -a_{j+1j}a_{j+2j+1}, & j \in \{1, 2, \dots, n-2\} \\ x_{ij} &= -c_{i-1j}a_{ii-1}, & j \in \{1, 2, \dots, n-2\}, \\ & & i > j+2 \end{aligned}$$

□

Lemma 4.1 *Let A be a combinatorially symmetric, partial N -matrix of size 4×4 , such that $A \in PS_4$ and its associated graph is a cycle. Then, there exists an N -matrix completion.*

Proof: We may assume that A has the following form:

$$A = \begin{bmatrix} -1 & 1 & x_{13} & a_{14} \\ a_{21} & -1 & 1 & x_{24} \\ x_{31} & a_{32} & -1 & 1 \\ a_{41} & x_{42} & a_{43} & -1 \end{bmatrix},$$

where $a_{21}, a_{32}, a_{43}, a_{14}, a_{41}$ are positive.

We consider the following partial N -matrix in PS_4

$$\bar{A} = \begin{bmatrix} -1 & 1 & x_{13} & a_{14} \\ a_{21} & -1 & 1 & x_{24} \\ -a_{32} & a_{32} & -1 & 1 \\ a_{41} & -a_{41} & a_{43} & -1 \end{bmatrix},$$

and we prove that there exist values for x_{13} and x_{24} such that \bar{A}_c is an N -matrix. Therefore, there exists an N -matrix completion A_c of A . □

We can extend this result for matrices of size $n \times n$, $n \geq 4$.

Theorem 4.1 *Let A be a combinatorially symmetric, partial N -matrix of size $n \times n$, such that $A \in PS_n$ and its associated graph is a cycle. Then, there exists an N -matrix completion.*

Proof: We may assume that A has the following form:

$$A = \begin{bmatrix} -1 & 1 & \cdots & x_{1n-1} & (-1)^n a_{1n} \\ a_{21} & -1 & \cdots & x_{2n-1} & x_{2n} \\ \vdots & \vdots & & \vdots & \vdots \\ x_{n-11} & x_{n-12} & \cdots & -1 & 1 \\ (-1)^n a_{n1} & x_{n2} & \cdots & a_{nn-1} & -1 \end{bmatrix},$$

where $a_{1n}, a_{n1} > 0$ and $a_{ii-1} > 0$, $i = 2, 3, \dots, n$.

The proof is by induction on n . For $n = 4$ see Lemma 4.1. Now, let A be an $n \times n$ matrix. Consider the following partial N -matrix in PS_n :

$$\bar{A} = \left[\begin{array}{cccc|c} -1 & 1 & \cdots & (-1)^{n-1} a_{1n} & x_{1n} \\ a_{21} & -1 & \cdots & x_{2n-1} & x_{2n} \\ \vdots & \vdots & & \vdots & \vdots \\ (-1)^{n-1} a_{n1} & x_{n-12} & \cdots & -1 & 1 \\ \hline x_{n1} & x_{n2} & \cdots & a_{nn-1} & -1 \end{array} \right].$$

$\bar{A}[\{1, 2, \dots, n-1\}]$ is a partial N -matrix in PS_{n-1} such that its associated graph is an $(n-1)$ -cycle. By induction hypothesis there exists an N -matrix completion $\bar{A}[\{1, 2, \dots, n-1\}]_c$. Let \hat{A} be the partial N -matrix obtained by replacing in \bar{A} the completion $\bar{A}[\{1, 2, \dots, n-1\}]_c$.

By applying Proposition 3.1 to matrix \hat{A} we obtain an N -matrix completion A_c of A . □

References

- [1] A. Berman, R.J. Plemmons, *Nonnegative matrices in the Mathematical Sciences*, SIAM, 1994.
- [2] J.R.S. Blair, B. Peyton, An introduction to chordal graphs and clique trees, *The IMA volumes in Mathematics and its Applications*, vol. 56, Springer, New York, 1993, pp. 1-31.
- [3] K. Inada, The production matrix and Stolpmer-Samuelson condition, *Econometrica*, 39: 219-239 (1971).

- [4] J.S. Maybee, Some aspects of the theory of PN-matrices, *SIAM J. Appl. Math.*, 312: 397-410, (1976).
- [5] S.R. Mohan, R. Sridhar, On characterizing N-matrices using Linear Complementarity, *Linear Algebra and its Applications*, 160: 231-245, (1992).
- [6] S.R. Paranjape, Simple proofs for the infinite divisibility of multivariate gamma distributions, *Sankhya Ser. A*, 40: 393-398, (1978).
- [7] T. Parthasarathy, G. Ravindran, N-matrices, *Linear Algebra and its Applications*, 139: 89-102, (1990).
- [8] R. Saigal, On the class of complementary cones and Lemke's algorithm, *SIAM J. Appl. Math.*, 23: 46-60, (1972).