

1 **Fungal diversity associated to the olive moth, *Prays oleae* Bernard: a survey for**
2 **potential entomopathogenic fungi**

3
4 Ivo Oliveira¹, José A. Pereira¹; Teresa Lino-Neto², Albino Bento¹, Paula Baptista^{1*}

5
6 ¹CIMO / School of Agriculture, Polytechnic Institute of Bragança, *Campus* de Santa
7 Apolónia, Apartado 1172, 5301-854 Bragança, Portugal.

8 ² Centre for Biodiversity Functional and Integrative Genomics (BioFIG), Plant
9 Functional Biology Centre, University of Minho, *Campus* de Gualtar, 4710-057 Braga,
10 Portugal.

11
12
13 Running title: Fungal associated to *Prays oleae*

14 Submission date: 28-05-2011

15
16
17
18 * Corresponding author. Tel.: + 351 273303332; Fax + 351 273 325405.

19 *E-mail address:* pbaptista@ipb.pt (P. Baptista).

20

21 **ABSTRACT**

22 Olive production is one of the main agricultural activities in Portugal. In the region of
23 Trás-os-Montes this crop has been considerably affected by *Prays oleae*. In order to
24 evaluate the diversity of fungi on *P. oleae* population of Trás-os-Montes olive orchards,
25 larvae and pupae of the three annual generations (phyllophagous, antophagous and
26 carpophagous) were collected and evaluated for fungal growth on their surface. From
27 the 3828 larvae and pupae, a high percentage of individuals exhibited growth of a
28 fungal agent (40.6%), particularly those from the phyllophagous generation. From all
29 the moth generations, a total of 43 species from 24 genera were identified, but the
30 diversity and abundance of fungal species differed between the three generations.
31 Higher diversity was found in the carpophagous generation, followed by the
32 antophagous and phyllophagous generations. The presence of fungi displaying
33 entomopathogenic features was highest in the phyllophagous larvae and pupae, being
34 *B. bassiana* the most abundant taxa. The first report of *B. bassiana* presence on *P. oleae*
35 could open new strategies for the biocontrol of this major pest in olive groves, since the
36 use of an already adapted species increases the guarantee of success of a biocontrol
37 approach. The identification of antagonistic fungi able to control agents that cause
38 major olive diseases, such as *Verticillium dahliae*, will benefit future biological control
39 approaches for limiting this increasingly spreading pathogen.

40

41 **Keywords:** Olive tree; *Prays oleae*; fungal diversity; moth life cycle generation

42

43 **Introduction**

44 The olive tree is an important crop for Mediterranean basin countries including
45 Portugal. Extensive areas occupied by olive groves in Trás-os-Montes region (Northeast
46 of Portugal), not only have a significant economic impact, but also exhibit a social,
47 environmental and landscape significance. The olive moth, *Prays oleae* Bern., is one of
48 the major pests on these olive orchards, being responsible for high losses in the olive
49 yield as much as 40% [33]. This lepidopteran presents three generations per year that
50 damage the olive tree in different organs. The antophagous generation, occurring from
51 April to June, causes damages to the olive tree flowers; the carpophagous generation,
52 which usually appears from July to September, attacks the fruits leading to their
53 premature fall; and the phyllophagous generation, present from October to March of the
54 following year, damages the leaves [33]. Due to the growing awareness of detrimental
55 effects of pesticides to the ecosystems [6], agricultural practices in these orchards have
56 been changing to biological agriculture. Therefore, the search for methods to control
57 pests and diseases has acquired a new motivation. One of the promising methods to
58 control pests can be the use of entomopathogenic fungi, which are known by their
59 ability to infect and kill several insect species [25].

60 Entomopathogenic fungi comprise a large group of pathogens that includes
61 approximately 700 species in almost 85 genera [5]. Due to their large host range,
62 entomopathogenic fungi could be potentially useful as control agents against different
63 insect orders. When compared to conventional chemical pesticides, the use of insect
64 pathogens presents many advantages, such as the safety for humans and other non-target
65 organisms, environmental reduction of pesticides residues and a smaller effect on the
66 natural biodiversity [15]. However, pest management using entomopathogenic fungi has
67 been difficult to achieve. The ecological and environmental variations within agro-

68 ecosystems turn the formulation and application of this kind of biocontrol agent difficult
69 to manage [42]. The isolation of native fungi could provide a collection of isolates, for
70 the development of potential control agents already adapted and suited to a particular
71 habitat. In this work, the fungal diversity encountered on *P. oleae* population was
72 evaluated in olive groves from Trás-os-Montes and will be discussed taking into
73 account the olive moth generation where it appears. From this survey, potential
74 entomopathogenic fungi for future biocontrol strategies will be selected.

75

76 **Methods**

77 Study area

78 Larvae (mostly in the fourth and fifth instars of development) and pupae of *Prays oleae*
79 were collected in six olive groves located in Mirandela – Bragança region, Northeast of
80 Portugal (GPS coordinates: 41°34'03.77''N; 7°05'39.21''W; 41°33'53.29N,
81 7°05'40.23''W; 41°33'52.51''N, 7°05'30.59W; 41°33'33.11N, 7°05'35.62W;
82 41°33'08.02''N, 7°07'24.87''W; 41°32'35.20''N, 7°07'26.27''W). This region
83 comprises an olive tree growing area around 65.000 ha, corresponding to 22% of the
84 olive tree covered area in Portugal. It contains 20% of Portuguese olive trees, which
85 altogether are responsible for 39% of national olive oil production [31]. The topography
86 of this region is mountainous, with altitudes ranging between 300 and 500 m. The
87 prevailing climate is Mediterranean, with cold and rainy winters and long, hot and dry
88 summers. The average annual rainfall ranged from 600 to 800 mm, occurring mainly
89 between October and February, and the annual mean temperature ranged from 9 to
90 20°C. For this study, the selected orchards have been managed through organic [11] or
91 integrated production guidelines [18] and display high levels of olive moth infestations,
92 as observed by delta traps baited with pheromone. Although two different management

93 systems are referred, the studied olive groves are very similar, with low-external inputs.
94 The groves area ranged from 0.4 to 1.5 ha. Trees are of medium size with a planting
95 density of 7 x 7 m. Pruning was made every 2-3 years. No irrigation was done, and only
96 copper based products were used in the beginning of autumn for fungal diseases control
97 and no other pesticides were used in the last five years. The differences between the two
98 management systems are related with the products allowed in soil fertilization. The
99 most important cultivars were Cobrançosa and Verdeal Transmontana.

100

101 Experimental design and collection of plant material

102 The collection of larvae and pupae was performed in 15 randomly chosen trees in each
103 of the selected orchards for the three annual generations of *P. oleae*. Sampling dates
104 were variable according to the life stages of the pest. For the phyllophagous generation
105 the collection was conducted between 17th and 27th March of 2009, for the antophagous
106 generation between 24th May and 5th June and for the carpophagous generation between
107 1st and 3rd of September of the same year. For the first two generations, sampling of
108 leaves (phyllophagous generation) and flower clusters (antophagous generation), in
109 which larvae and pupae were present, was performed. For the carpophagous generation,
110 the collection of whole fruits was randomly performed, due to the presence of larvae
111 and pupae within olives. The plant material (leaves and flower clusters) and olives were
112 collected individually to sterile bags, and processed within a few hours after sampling.

113

114 Isolation of fungi from mycosed larvae and pupae of *P. oleae*

115 In the laboratory, the plant material was further examined for the presence of olive moth
116 larvae and pupae. Since carpophagous larvae develop inside the olive stone, the stone
117 was smoothly broken to expose larvae. Whenever present, the larvae and pupae were

118 individually placed into sterile tubes containing a food source (leaves for the
119 phyllophagous generation, flower buds for the antophagous generation, and olive stones
120 for the carpophagous generation). The tubes were sealed and maintained at $25 \pm 2^\circ\text{C}$,
121 under 16/8 hours light/dark regime, for an average period of 10 days. Larvae and pupae
122 were daily observed, in order to detect their death or evaluate the insect emergence.
123 Each time a fungal agent was growing on the surface of dead larvae or pupae, the fungal
124 specimen was isolated by inoculating Potato Dextrose Agar (PDA) medium,
125 supplemented with 0.01% (w/v) chloramphenicol (Oxoid). Pure cultures of each isolate
126 were deposited in the culture collection of the Polytechnic Institute of Bragança (School
127 of Agriculture).

128

129 Extraction of fungal DNA

130 Fungal isolates were inoculated onto PDA medium and maintained at $25 \pm 2^\circ\text{C}$ in the
131 dark for 1-2 weeks. The colony morphology, spore size and shape were used for the first
132 identification and to group strains. Spores were collected from each isolate and used for
133 DNA extraction. Isolation of genomic DNA was performed by transferring spores into a
134 microtube containing 500 μL of Lysis buffer (200 mM Tris-HCl pH 8.0, 250 mM NaCl,
135 25 mM EDTA pH 8.0 and 0.5% SDS) and sterile glass spheres. Tubes were vortexed
136 for 5 minutes to disrupt cells. After addition of 250 μL of cold 3M NaOAc pH 5.5, the
137 mixture was gently homogenized by inversion and incubated for 10 minutes at -20°C .
138 Following centrifugation at 10.500 rpm (4°C) for 10 minutes, the supernatant was
139 collected to another microtube and one volume of isopropanol (-20°C) was added. This
140 mixture was slowly homogenized and incubated at -20°C for one hour. The DNA
141 precipitate was collected by centrifugation at 10.500 rpm (4°C) for 10 minutes and the
142 pellet washed with cold 70% ethanol. The DNA pellet was air-dried for 20 min at room

143 temperature and re-suspended in 50 μ L of ultra pure water. DNA was stored at -20°C
144 until use.

145

146 Molecular identification of fungal isolates

147 Molecular identification was achieved by amplification of the internal transcribed
148 spacer region (ITS), using the universal primers *ITS1* and *ITS4* [44]. PCR reactions

149 (50 μ l) comprised 50 ng of genomic DNA, 0.2 μ M of each primer (*ITS1* and *ITS4*), 1x
150 GoTaq® Flexi buffer (Promega), 2 mM MgCl_2 (Promega), 0.2 μ M dNTP Mix

151 (Fermentas) and 1 U GoTaq® DNA polymerase (Promega). Amplifications were
152 carried out in the thermocycler Biometra UNO II (Thermoblock, Biotron) using a

153 temperature gradient protocol as follows: initial denaturation at 94°C for 3 min,
154 followed by 35 cycles of 0.5 min at 94°C , 0.5 min at 55°C , 1 min at 72°C , and a final 10

155 min extension at 72°C . PCR amplification products were analysed by electrophoresis
156 and those reactions that amplified a single PCR product were selected for purification

157 using the JETquick PCR product purification kit (Genomed). Amplified fragments were
158 sequenced using both *ITS1* and *ITS4* primers at the sequencing services of STAB Vida

159 (Oeiras, Portugal). DNA sequences were analysed with DNASTAR v.2.58 software,
160 and fungal identification was performed using the NCBI database

161 (<http://www.ncbi.nlm.nih.gov>) and BLAST algorithm.

162 The ecological classification of identified species was only based on the impact they can
163 have on olive orchards regarding their potential role on pests and diseases control. Even

164 though many of the identified fungi are also considered general saprophytes, they were
165 only classified into three ecological roles: phytopathogenic, antagonistic and

166 entomopathogenic. The classification given to each fungal species was based on
167 previously described characteristics.

168

169 Data Analysis

170 For each *P. oleae* generation, the species richness, *Simpson (D)* and *Shannon-Wiener*
171 *(H)* diversity indexes, total and relative abundances were estimated. Calculations of
172 *Simpson* and *Shannon-Wiener* diversity indexes were done using the software *Species*
173 *Diversity and Richness* (v. 3.0). Total abundance (N) was estimated as the number of
174 isolates per fungal taxa, whereas the proportion of isolates from each fungal taxa in
175 relation to the total number of fungal isolates was considered as the relative abundance
176 of a certain taxa. Principal component analysis (PCA) was applied to access the
177 relationship between fungal taxa and the three *P. oleae* generations. PCA was
178 performed using the SPSS software, version 17.0 (SPSS, Inc.). It was applied as an
179 unsupervised approach for reducing the number of variables (43, corresponding to the
180 number of identified fungal species) to a smaller number of new derived variables
181 (principal component or factors) that adequately summarize the original information.
182 This analysis will define which fungal species are correlated with each olive moth
183 generation. PCA analysis also allowed the recognition of patterns in the data by plotting
184 them in a multidimensional space, using the new derived variables as dimensions (factor
185 scores). The aim of the PCA is to produce components suitable to be used as predictors
186 or response variables in subsequent analysis. The number of factors to keep in data
187 treatment was evaluated by the Scree plot, taking into account the eigenvalues and the
188 internal consistency by means of α Cronbach's value [21; 34].

189

190 **Results**

191 Diversity and abundance of fungi encountered in *P. oleae*

192 From the 3828 larvae and pupae (2552 larvae and 1276 pupae), collected in all
193 generations of *P. oleae*, an insect emergence percentage of 38.6% was observed (Table
194 1). Of the total number of dead larvae and pupae (1477), 40.6% (599) exhibited the
195 growth of a fungal agent on their surface (Table 1). From those, the isolation of fungi
196 allowed the identification of 43 species, belonging to 24 genera and 14 families (Table
197 2). The families comprising more diversity were Pleosporaceae and Quambalariaceae (8
198 species each) and Mucoraceae and Nectriaceae (5 species each). These families
199 accounted for as much as 62% of the total identified species. The greatest number of
200 taxa belonged to the genera *Penicillium* (7), *Arthrinium*, *Mucor*, *Fusarium* and
201 *Alternaria* (all with 4 taxa). Concerning abundance, 166 different isolates were obtained
202 in this work (Table 2). The most common taxa were *Beauveria bassiana* (N=50) and
203 *Fusarium oxysporum* (N=25), representing together 45.2% of the total identified
204 isolates.

205 The number of fungal taxa identified in the present study varied between the three
206 generations of *P. oleae* (Table 2). In the phyllophagous generation, where a total of 70
207 fungal isolates were obtained, 16 different species were identified, belonging to 13
208 genera and 9 families. The most representative family was Quambalariaceae,
209 comprising 25% of the identified species in this generation, being the genus *Penicillium*
210 the most represented (3 species). In the antophagous generation, 52 isolates were
211 obtained belonging to 13 species, 8 genera and 8 families. The families Nectriaceae and
212 Apiosporaceae include the majority of the identified taxa (54%). The genus that
213 included more taxa (4) was *Fusarium* (Nectriaceae), which contained 31% of the fungal
214 species surveyed in this generation. Concerning the carpophagous generation, 44 fungal
215 isolates were obtained being identified 21 species, belonging to 14 genera and 10
216 families. Pleosporaceae was the family representing more species (33%), and the genera

217 *Alternaria* (Pleosporaceae) and *Penicillium* (Quambalariaceae), both with 4 species
218 each, comprised 38% of the total fungal taxa found in this generation of *P. oleae*.
219 The most abundant species also differed between generations. For the phyllophagous
220 generation, the most abundant one was *B. bassiana* (N=49), which represented 70% of
221 the relative abundance in this generation. For the antophagous generation, *F. oxysporum*
222 was the most abundant (N=23), corresponding to a relative abundance of 44.2%. In the
223 carpophagous generation, several taxa presented comparable relative abundances, being
224 *P. pinophilum* the most frequent (N=7, corresponding to a relative abundance of
225 15.9%), followed by *Alternaria sp.1* and *A. ustus*, both with N=6 corresponding to
226 13.6% of relative abundance.

227 In order to provide more information about fungal community composition in each
228 *P. oleae* generation, the *Simpson (D)* and *Shannon-Wiener (H)* diversity indexes were
229 determined (Table 3). In the present work, both diversity indexes differed between the
230 three *P. oleae* generations. As verified for species richness, the highest fungal diversity
231 was verified in the carpophagous generation ($D=0.09$ and $H=2.70$). In fact, when using
232 larvae or pupae from this generation, the highest number of unique species (17) was
233 obtained. The number of exclusive species identified from phyllophagous and
234 antophagous generations was only 11 and 9, respectively. The lowest value of species
235 diversity ($D=0.5$ and $H=1.39$) was detected in the phyllophagous generation.

236 When performing a principal component analysis (PCA) using the total fungal taxa,
237 according to the moth generation from which they were isolated, it was possible to
238 separate the samples in eight groups (Fig 1). The phyllophagous generation is clearly
239 related to groups 4 and 7, the antophagous generation is linked to groups 3 and 6, and
240 the carpophagous generation is associated to groups 1 and 2. The two remaining groups
241 (5 and 8) do not appear associated to a specific generation. Group 5 is mainly correlated

242 to the antophagous generation, although presenting some relation to the phyllophagous
243 generation. Group 8 is equally correlated to both phyllophagous and carpophagous
244 generations.

245

246 Fungal ecological roles

247 Taking into account the potential use for the biological control of pests and diseases,
248 45% of the isolates found in this work have been described in the literature as
249 presenting the useful features of antagonism and/or entomopathogenicity (Fig 2A). The
250 remaining 55% of the isolates presented phytopathogenic features, displaying also in
251 large extent (41%) antagonistic and/or entomopathogenic characteristics. When
252 comparing the number of identified species, the majority (75%) has been described as
253 phytopathogenic fungi (Fig 2B). While the mentioned attribute was present alone in
254 28% of the identified species, the remaining 47% also displayed antagonistic and/or
255 entomopathogenic characteristics. From those species described as non-
256 phytopathogenic (23%), the majority are defined as antagonistic (17%).

257 The ecological roles of the identified fungi differed between generations. In the
258 phyllophagous generation, the majority (87%) of the fungal isolates have been
259 described as antagonistic and entomopathogenic (Fig 2A). Included in the
260 phytopathogenic fungi (23%), 13% of fungal isolates also display antagonistic and/or
261 entomopathogenic characteristics. This relation is reversed when considering the
262 number of identified fungal species. The number of phytopathogenic taxa was higher
263 (77%) than those displaying only antagonistic and/or entomopathogenic features (23%)
264 (Fig 2B).

265 In the antophagous generation, a large fraction of isolates (73%) has been described as
266 displaying phytopathogenic characteristics, of which 71% also displays antagonistic

267 and/or entomopathogenic features (Fig 2A). The remaining 27% has been described as
268 antagonistic fungi. Concerning the carpophagous generation, the higher fraction of
269 fungal isolates displays phytopathogenic characteristics (77%) (Fig 2A). Included in
270 these, 50% of fungal isolates also exhibits antagonistic and/or entomopathogenic
271 features. Fungal isolates displaying only antagonistic and/or entomopathogenic features
272 were also found but in a lower proportion (20%). When considering the number of
273 identified fungal taxa, the same trend was observed for the antophagous and
274 carpophagous generations. In both, the phytopathogenic fungi comprised the majority of
275 identified taxa, but most of them also present antagonistic and/or entomopathogenic
276 features. It was in the carpophagous generation that the lowest amount of non-
277 phytopathogenic species with antagonistic and/or entomopathogenic properties were
278 found (12%). This was the only generation where a fungal taxa (*Lichtheimia ramosa*)
279 with no described ecological role was isolated.

280

281 **Discussion**

282 In the present work, the diversity of fungi isolated from dead larvae and pupae of a
283 major olive pest (*P. oleae*) was evaluated for the first time. Following molecular
284 identification, it was possible to identify 43 fungal species, belonging to 24 genera and
285 14 families. The identification of such a high number of fungal taxa described as
286 presenting antagonistic, entomopathogenic or phytopathogenic features was only
287 possible because the studied olive groves were maintained under organic or integrated
288 production guidelines. If sampling had been performed on a conventional orchard the
289 expected fungal diversity would have been lower. Organic and integrated production
290 management creates a healthier and safer environment with higher biological diversity
291 [19]. Similar diversity levels (46 fungal species and 27 genera) were obtained when

292 studying insect-associated fungi isolated from soil samples of different field crops
293 (wheat/maize, corn and soybean) and orchards (peach and apple) [38].

294

295 Fungal diversity and abundance

296 Concerning genera diversity, three main patterns were detected: (i) species-rich and
297 highly abundant genera, e.g. *Fusarium*, *Alternaria* and *Penicillium* (about 35% of the
298 taxonomic diversity and 42% of the total of isolates), (ii) species-rich genera, but
299 displaying low abundance, e.g. *Mucor* and *Arthrimum* (about 19% of the taxonomic
300 diversity and 5% of the total of isolates); and (iii) species-poor but highly abundant
301 genera, e.g. *Beauveria* and *Trichoderma* (about 5% of the taxonomic diversity and 35%
302 of the total of isolates). Among the 43 species recorded during the present study, the
303 genera *Alternaria*, *Arthrimum*, *Fusarium*, *Mucor* and *Penicillium* were the most
304 represented in terms of taxa number. These genera include some of the most ubiquitous
305 fungal species in nature, which have been found in soils, plants and agricultural
306 communities. The wide presence of such microorganisms has been related to the broad
307 ecological roles they play, either as saprophyts, phytopathogens or biocontrol agents
308 [8].

309 The genera that contributed most to species diversity depended on which *P. oleae*
310 generation was used for fungal isolation. While in phyllophagous generation
311 *Penicillium* represented the genus with highest taxonomic diversity (18%), in
312 antophagous generation that genus was *Fusarium* (31%) and in carpophagous
313 generation were *Alternaria* and *Penicillium* (representing together 39%). The
314 composition of fungal community was also inferred by the *Simpson (D)* and *Shannon-*
315 *Wiener (H)* diversity indexes, which offer valuable information about rarity and
316 frequency of species in a community. Both indexes provide more information than

317 simply species richness, since they also take the relative abundances of different species
318 into account. While the carpophagous generation presented the highest fungal diversity
319 (displaying the lowest D and highest H values), the phyllophagous generation displayed
320 the lowest diversity (presenting the highest D and lowest H values). This difference
321 between moth generations could be related to two main factors: climatic conditions
322 during larvae and pupae collection and the moth life cycle.

323 Climate conditions, in particular relative humidity and temperature, are known to affect
324 both conidia dispersion and germination [39]. The collection of larvae and pupae from
325 phyllophagous generation (March) matched with the time of year where the temperature
326 is low, decreasing the ability of spore germination and thus reducing fungal diversity.
327 Furthermore, the moth life cycle also seems to strongly affect the fungal diversity
328 observed in this generation, as a large part of phyllophagous moth development occurs
329 in the leaves as miner larvae. Therefore, the reduced fungal diversity observed in this
330 generation can be associated to the low chances of larvae contact with fungal spores.

331 The high occurrence of the entomopathogenic *B. bassiana* in the phyllophagous
332 generation (70% of the isolates) could be related with an epizootic outbreak that might
333 have occurred during the time of larvae and pupae sampling. These epizootics are
334 described as being dependent on host population dynamics, the number and viability of
335 infective stages in the pathogen population, infection efficiency and development, in
336 addition to a complex set of environmental factors and timing [25]. Although dispersal
337 of *B. bassiana* conidia by larvae could have been limited, due to their small
338 displacement in trees, the wind and rain may have caused the spread of those infectious
339 structures. Also, the high number of infected *P. oleae* individuals could have functioned
340 as sources of infective conidia, thus contributing for the abundance of *B. bassiana* in
341 this generation.

342 In antophagous and carpophagous generations, the most abundant taxa were *Fusarium*
343 *oxysporum* (44% of the total isolates) and *Penicillium pinophilum* (15.9% of the total
344 isolates), respectively. As referred, these two taxa are very frequent in nature, being
345 present in almost all environments. Furthermore, the amount of spores of *Fusarium*
346 species is known to increase from April to July [39] and those from *Penicillium* species
347 rise between August and October [23]. These periods are coincident with the sampling
348 dates of larvae and pupae from antophagous (May-June) and carpophagous (September)
349 generations, explaining in some degree the high abundance of *Fusarium* and
350 *Penicillium* genera in antophagous and carpophagous generations, respectively.

351 The association of climate and life cycle negatively affect the fungal diversity when
352 using antophagous larvae and pupae. In the region where the study was conducted,
353 temperature rapidly increases during May and June, leading to an extremely fast larvae
354 development. Therefore, larvae and pupae stay a short period exposed to the
355 surrounding environment, reducing the possibility of fungal infection. In contrast,
356 carpophagous larvae are exposed to fungal spores for a longer period. After ecloding
357 from the egg, larvae must travel from the oviposition site in the fruits to the petiole and
358 bore down into the stone, where they usually feed for several weeks. After completing
359 their development, larvae re-emerge from the fruit. The chance of infection is still
360 enhanced by the extension of carpophagous generation which, together with the
361 favourable climatic conditions, allows the development of more fungal taxa.
362 Accordingly, it was in this moth generation that higher species diversity was found.

363

364 Fungal ecological roles

365 The relation between the amount of dead larvae and pupae displaying fungal growth and
366 the ecological role of the identified fungal taxa provides interesting data. The higher

367 amount of infected dead larvae and pupae was observed in the phyllophagous
368 generation (61.5%). It was also from this generation that the most isolates exhibiting
369 entomopathogenic features (76%) were obtained. As the presence of fungal growth on
370 dead larvae and pupae decreases (from 61.5% to 32.7% and finally 23.0%, in the
371 phyllophagous, antophagous and carpophagous generations, respectively), the
372 abundance of fungi exhibiting entomopathogenic characteristics also decreases (76%,
373 50% and 30%, respectively). These results seem to suggest that the infection with
374 entomopathogenic fungi could have led to larvae and pupae death. Most of the fungal
375 isolates (86%) and species (70%) identified in this work has been described as
376 antagonistic and/or entomopathogenic, although some of them have also been
377 considered as displaying phytopathogenic features. Because only the non-
378 phytopathogenic fungi displaying antagonistic and/or entomopathogenic features could
379 be explored for limiting fungal diseases and/or pests, the percentage of identified fungal
380 species that might have a future application as biocontrol agents is reduced to 23%.

381 From identified taxa in this work, *B. bassiana* seems to be the most conspicuous among
382 the entomopathogenic/antagonistic species. The natural occurrence of this fungus in
383 over 700 insect hosts from almost all taxonomic orders is well documented [25].
384 However, there are still some reservations about the host range of this fungus. Some
385 authors claim that *B. bassiana* is a “species complex”, referring that different isolates
386 have a restricted host, while others point out that this fungus has no host specificity
387 [29]. Accordingly, several pests are susceptible to the entomopathogenic aptitude of this
388 fungus, like *Alphitobius diaperinus* [36], whiteflies *Bemisia tabaci* and *Trialeurodes*
389 *vaporariorum* [32], *Capnodis tenebrionis* [20], *Lutzomyia longipalpis* [2],
390 *Callosobruchus maculatus* [26], and *Tetranychus urticae* [10]. Some studies also
391 indicate the presence of this fungus associated to several lepidopterans [2, 3, 7, 13, 22,

392 32, 36]. In addition, natural occurrence of mycoses caused by *B. bassiana* is reported in
393 lepidopteran pests such as *Helicoverpa armigera* (Hubner) and *Spodoptera litura* (Fab.)
394 [9], *Argyresthia conjugella* [41], and *Plutella xylostella* [37]. As far as we know this is
395 the first report of *B. bassiana* related to *P. oleae*, where a large amount of isolates were
396 obtained from larvae and pupae of the phyllophagous generation. The presence of this
397 fungus in olive grove soils has already been reported [20, 32], as well as its ability to
398 control pests in this crop, such as *Bactrocera oleae* [14, 17]. All these evidences may
399 suggest that this fungus could be effective in controlling *P. oleae* in olive orchards,
400 although experiments to evaluate its infecting ability towards *P. oleae* have to be
401 performed. The natural occurrence of this species in the studied olive groves guarantees
402 an already adapted and suited strain to be used as a control agent in this particular
403 ecosystem.

404 Another potential entomopathogen identified in the present study that could be able to
405 control *P. oleae* larvae is *Cordyceps sinensis*. Although no literature is available for its
406 ability to infect *P. oleae*, the capacity of infecting other lepidopterans larvae was
407 already described, such as those of *Hepialus armoricanus* [30]. Nevertheless, these
408 fungal taxa described as entomopathogenic cannot be definitely linked to the cause of
409 death of *P. oleae* larvae and pupae. Assays confirming the infection ability and
410 virulence of such fungi must be performed, in order to confirm their entomopathogenic
411 potential.

412 The identification of fungi that could limit the growth of other infectious fungi by their
413 antagonistic properties may also be important for designing future biocontrol strategies
414 for restricting fungal diseases in olive groves. The most abundant taxa with antagonistic
415 characteristics identified in the present study were *Penicillium pinophilum* and
416 *Trichoderma gamsii*. *P. pinophilum* is one of the most important antagonists of

417 *Rhizoctonia solani*, a fungal pathogen of tobacco [1]. Although scarce information is
418 available about *T. gamsii*, antagonistic properties of *Trichoderma* species against a great
419 number of fungal species have already been described, including *Verticillium dahliae*
420 that causes one of the most severe diseases affecting olive (Verticillium wilt) [27, 43].
421 The identification of several isolates with antagonistic features (44%), some of them
422 against fungi that cause olive diseases, opens up the possibility of further research on
423 those antagonistic species for limiting the occurrence of such phytopathogenic fungi in
424 olive orchards.

425 In this work, besides the identification of potential biocontrol agents for limiting pests
426 and fungal diseases in olive groves, many other fungi were identified that could play a
427 role in olive grove ecosystems. One of the most common genus found in this work was
428 *Alternaria* that comprise species that have already been reported to cause spoilage of
429 olives [35] and cause a disease on olive shoots grown under greenhouse conditions [4].
430 The most frequent *Alternaria* species isolated in the present work was *A. tenuissima*,
431 which has been associated to late blight of pistachio and black point of small-grain
432 cereals [16], among other crops. One of the most abundant fungal taxa identified in this
433 study was *Fusarium oxysporum* that exhibits antagonistic, entomopathogenic and
434 phytopathogenic properties. This species has been mainly described as phytopathogenic,
435 causing vascular wilts or rot and crown rots in a large number of crops, including tree
436 crops [12]. This species also presents antagonistic features against *Colletotrichum*
437 *gloeosporioides* and *Pestalotia psidii* [28], and most important, against *Verticillium*
438 *dahliae* [24]. Furthermore, *F. oxysporum* has also been described as an opportunistic
439 insect-pathogen [38]. Accordingly, it has been isolated from several insect hosts, such
440 as those from Homoptera and Coleoptera orders, being able to parasite the greenhouse
441 whitefly, *Trialeurodes vaporariorum* [40].

442 However, the identification of entomopathogenic fungi from dead *P. oleae* larvae and
443 pupae cannot be unequivocally associated to their ability to infect this lepidopteran,
444 since dead larvae and pupae could just become increasingly susceptible to fungi.

445

446 Conclusion

447 As far as we know, the present work describes for the first time the assessment of fungal
448 diversity directly obtained from mycosed dead *P. oleae* larvae and pupae.

449 The strategy used for obtaining fungal isolates (collection of larvae and pupae from the
450 field and isolation of fungi from cadavers) allowed the identification of 43 fungal
451 species, displaying several ecological roles. The diversity and abundance of fungal
452 species differed when using larvae or pupae from different moth generation. Higher
453 fungal diversity was found in the carpophagous generation, followed by antophagous
454 and phyllophagous generations. Although the identified taxa could not be unequivocally
455 associated with the cause of moth death, almost 37% of identified taxa presented
456 entomopathogenic properties.

457 The identification of entomopathogenic and antagonist fungi in olive orchards provided
458 a pool of biocontrol agents that could be used in the future for controlling pests and
459 fungal diseases. The first report of *Beauveria bassiana* presence on *P. oleae* could open
460 new strategies for the biocontrol of this major pest in olive groves. This fungus was
461 isolated with high incidence from phyllophagous generation larvae and pupae. Future
462 studies will be directed to screening the entomopathogens isolates for virulence to adult
463 *P. oleae*. As the environmental conditions influence the performance of a given strain of
464 fungus, the use of already adapted species to a particular ecosystem increases the
465 guarantee of success of a biocontrol approach. The occurrence of antagonistic fungi able
466 to control one of the major fungus attacking olive tree (*Verticillium dahliae*) may as

467 well be investigated, in order to control this pathogen that is spreading throughout olive
468 orchards.

469

470 **Acknowledgements**

471 This work was supported by Science and Technology Foundation (Fundação para a
472 Ciência e Tecnologia – FCT) project PTDC/AGR-AAM/102600/2008
473 “Entomopathogenic fungi associated to olive pests: isolation, characterization and
474 selection for biological control”. The first author is grateful to the Science and
475 Technology Foundation for the PhD grant SFRH/BD/44265/2008.

476

477 **References**

- 478 1. Alagesaboopathi C (1994) Biological control of damping-off disease of cotton
479 seedling. *Curr Sci* 66:864-868.
- 480 2. Amóra S, Bevilaqua C, Feijó F, Silva M, Pereira R, Silva S, Alves N, Freire F,
481 Oliveira D (2009) Evaluation of the fungus *Beauveria bassiana*
482 (Deuteromycotina: Hyphomycetes), a potential biological control agent of
483 *Lutzomyia longipalpis* (Diptera, Psychodidae). *Biol Control* 50:329-335.
- 484 3. Aquino de Muro M, Elliot S, Moore D, Parker B, Skinner M, Reid W, El
485 Bouhssni M 2005 Molecular characterisation of *Beauveria bassiana* isolates
486 obtained from overwintering sites of Sunn Pests (*Eurygaster* and *Aelia* species).
487 *Mycol Res* 109:294-306.
- 488 4. Bourbos V, Skoundridakis M, Metzidakis I (1999) *Alternaria alternata*: a new
489 disease of leafy cuttings of olive shoots. In: Metzidakis I, Voyiatzis D (eds),
490 Proceedings 3rd International ISHS Symposium on Olive Growing, *Acta*
491 *Horticulturae* 474:585-587.
- 492 5. Charnley A, Collins S (2007) Entomopathogenic fungi and their role in pest
493 Control. In: Kubicek C, Druzhinina I (eds) *The Mycota, Volume 4 -*
494 *Environmental and Microbial Relationships* 2nd edn. Springer-Verlag Berlin
495 Heidelberg, pp 159-187.
- 496 6. Cuthbertson A, Walters K (2005) Pathogenicity of the entomopathogenic
497 fungus, *Lecanicillium muscarium*, against the sweetpotato whitefly *Bemisia*
498 *tabaci* under laboratory and glasshouse conditions. *Mycopathologia* 160:315-
499 319.

- 500 7. Dalzoto P, Glienke-Blanco C, Kava-Cordeiro V, Ribeiro J, Kitajima E, Azevedo
501 J (2006) Horizontal transfer and hypovirulence associated with double-stranded
502 RNA in *Beauveria bassiana*. Mycol Res 110:1475-1481.
- 503 8. De Lucca A (2007) Harmful fungi in both Agriculture and Medicine. Rev
504 Iberoamer Micol 24:3-13.
- 505 9. Devi K, Sridevi V, Mohan C, Padmavathi J (2005) Effect of high temperature
506 and water stress on in vitro germination and growth in isolates of the
507 entomopathogenic fungus *Beauveria bassiana* (Bals.) Vuillemin. J Invertebr
508 Pathol 88:181-189.
- 509 10. Eken C, Hayat R (2009) Preliminary evaluation of *Cladosporium*
510 *cladosporioides* (Fresen.) de Vries in laboratory conditions, as a potential
511 candidate for biocontrol of *Tetranychus urticae* Koch. World J Microbiol
512 Biotechnol 25:489-492.
- 513 11. European Union. Council Regulation (EC) No 834/2007 of 28 June 2007
- 514 12. Fravel D, Olivain C, Alabouvette C (2002) *Fusarium oxysporum* and its
515 biocontrol. New Phytol 157:493-502.
- 516 13. Fuguet R, Vey A (2004) Comparative analysis of the production of insecticidal
517 and melanizing macromolecules by strains of *Beauveria* spp.: *in vivo* studies. J
518 Invertebr Pathol 85:152-167.
- 519 14. Konstantopoulou M, Mazomenos B (2005) Evaluation of *Beauveria bassiana*
520 and *B. brongniartii* strains and four wild-type fungal species against adults of
521 *Bactrocera oleae* and *Ceratitis capitata*. BioControl 50:293-305.
- 522 15. Lacey L, Frutos R, Kaya H, Vail P (2001) Insect pathogens as biological control
523 agents: Do They Have a Future? Biol Control 21:230-248.

- 524 16. Logrieco A, Bottalico A, Mulé G, Moretti M, Perrone, G (2003) Epidemiology
525 of toxigenic fungi and their associated mycotoxins for some Mediterranean
526 crops. Eur J Plant Pathol 109:645-667.
- 527 17. Mahmoud M (2009) Pathogenicity of three commercial products of
528 entomopathogenic fungi, *Beauveria bassiana*, *Metarhizium anisopilae* and
529 *Lecanicillium lecanii* against adults of olive fly, *Bactrocera oleae* (Gmelin)
530 (Diptera: Tephritidae) in the laboratory. Plant Protect Sci 45:98-102.
- 531 18. Malavolta C, Delrio G, Boller EF (2002) Guidelines for Integrated Production of
532 Olives. IOBC Technical Guideline III, 1st Edition, 25 (4), pp. 1-8.
- 533 19. Mander U, Mikk M, Külvik M (1999) Ecological and low intensity agriculture
534 as contributors to landscape and biological diversity. Landsc Urban Plann
535 46:169-177.
- 536 20. Marannino P, Santiago-Álvarez C, Lillo E, Quesada-Moraga E (2006) A new
537 bioassay method reveals pathogenicity of *Metarhizium anisopliae* and *Beauveria*
538 *bassiana* against early stages of *Capnodis tenebrionis* (Coleoptera; Buprestidae).
539 J Invertebr Pathol 93:210-213.
- 540 21. Maroco J (2003) Análise Estatística, com utilização do SPSS. Edições Sílabo,
541 Lisboa, Portugal, 822 pp.
- 542 22. Maurer P, Couteaudieri Y, Girard P, Bridge P, Riba G (1997) Genetic diversity
543 of *Beauveria bassiana* and relatedness to host insect range. Mycol Res 101:159-
544 164.
- 545 23. Medrela-Kuder E (2010) Seasonal variations in the occurrence of culturable
546 airborne fungi in outdoor and indoor air in Craców. Int Biodeterior Biodegrad
547 52:203-205.

- 548 24. Mercado-Blanco J, Rodríguez-Jurado D, Hervás A, Jiménez-Díaz R (2004)
549 Suppression of *Verticillium* wilt in olive planting stocks by root-associated
550 fluorescent *Pseudomonas* spp. *Biol Control* 30:474-486.
- 551 25. Meyling N, Eilenberg J (2007) Ecology of the entomopathogenic fungi
552 *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems:
553 Potential for conservation biological control. *Biol Control* 43:145-155.
- 554 26. Murad A, Laumann R, Mehta A, Noronha E, Franco O (2007) Screening and
555 secretomic analysis of entomopathogenic *Beauveria bassiana* isolates in
556 response to cowpea weevil (*Callosobruchus maculatus*) exoskeleton. *Comp*
557 *Biochem Physiol C* 145:333-338.
- 558 27. Ordentlich A, Nachmias I (1990) Integrated control of *Verticillium dahliae* in
559 potato by *Trichoderma harzianum* and captan. *Crop Prot* 9:363-366.
- 560 28. Pandey R, Arora D, Dubey R (1993) Antagonistic interactions between fungal
561 pathogens and phylloplane fungi of guava. *Mycopathologia* 124:31-39.
- 562 29. Pathan A, Devi K, Vogel H, Reineke A (2007) Analysis of differential gene
563 expression in the generalist entomopathogenic fungus *Beauveria bassiana*
564 (Bals.) Vuillemin grown on different insect cuticular extracts and synthetic
565 medium through cDNA-AFLPs. *Fungal Genet Biol* 44:1231-1241.
- 566 30. Paterson R (2008) *Cordyceps* – A traditional Chinese medicine and another
567 fungal therapeutic biofactory? *Phytochemistry* 69:1469-1495.
- 568 31. Pereira J, Bento A, Cabanas E, Torres L, Herz A, Hassan A (2004) Ants as
569 Predators of the Egg Parasitoid *Trichogramma cacoeciae* (Hymenoptera:
570 Trichogrammatidae) Applied for Biological Control of the Olive Moth, *Prays*
571 *oleae* (Lepidoptera: Plutellidae) in Portugal. *Biocontrol Sci Technol* 14:653-664.

- 572 32. Quesada-Moraga E, Maranhao E, Valverde-García P, Santiago-Álvarez C (2006)
573 Selection of *Beauveria bassiana* isolates for control of the whiteflies *Bemisia*
574 *tabaci* and *Trialeurodes vaporariorum* on the basis of their virulence, thermal
575 requirements, and toxicogenic activity. Biol Control 36:274-287.
- 576 33. Ramos P, Campos M, Ramos J (1998) Long-term study on the evaluation of
577 yield and economic losses caused by *Prays oleae* Bern. in the olive crop of
578 Granada (southern Spain). Crop Prot 17: 645-647.
- 579 34. Rencher AC (1995) Methods of Multivariate Analysis. John Willey, New York,
580 USA, 708 pp.
- 581 35. Roussos S, Zaouia N, Salih G, Tantaoui-Elaraki A, Lamrani K, Cheheb M,
582 Hassouni H, Verh F, Perraud-Gaime I, Augur C, Ismaili-Alaoui M (2006)
583 Characterization of filamentous fungi isolated from Moroccan olive and olive
584 cake: Toxinogenic potential of *Aspergillus* strains. Mol Nutr Food Res 50:500-
585 506.
- 586 36. Santoro P, Neves P, Alexandre T, Sartori D, Alves L, Fungaro M (2008)
587 Selection of *Beauveria bassiana* isolates to control *Alphitobius diaperinus*. J
588 Invertebr Pathol 97:83-90.
- 589 37. Silva V, Barros R, Marques E, Torres J (2003) Suscetibilidade de *Plutella*
590 *xylostella* (L.) (Lepidoptera: Plutellidae) aos Fungos *Beauveria bassiana* (Bals.)
591 Vuill. e *Metarhizium anisopliae* (Metsch.) Sorok. Neotrop Entomol 32:653-658.
- 592 38. Sun B, Yu H, Chen A, Liu X (2008) Insect-associated fungi in soils of field
593 crops and orchards. Crop Prot 27:1421-1426.
- 594 39. Topbaş M, Tosun, U, Çan G, Kaklikkaya N, Aydin F (2006) Identification and
595 seasonal distribution of airborne fungi in urban outdoor air in an Eastern Black
596 Sea Turkish town. Turk J Med Sci 31:31-36.

- 597 40. Torres-Barragán A, Anaya A, Alatorre R, Toriello C (2004) Entomopathogenic
598 fungi from 'El Eden' Ecological Reserve, Quintana Roo, Mexico.
599 Mycopathologia 158:61-71.
- 600 41. Vänninen I, Hokkanen H (1997) Efficacy of Entomopathogenic fungi and
601 Nematodes against *Argyresthia conjugella* (Lep: Yponomeutidae).
602 Entomophaga 42:377-385.
- 603 42. Vega F, Goettel M, Blackwell M, Chandler D, Jackson M, Keller S, Koike M,
604 Maniania N, Monzón A, Ownley B, Pell J, Rangel D, Roy H (2009) Fungal
605 entomopathogens: new insights on their ecology. Fungal Ecol 2:149-159.
- 606 43. Verma M, Brar S, Tyagi R, Surampalli R, Valéro J (2007) Antagonistic fungi,
607 *Trichoderma* spp.: Panoply of biological control. Biochem Eng J 37:1-20.
- 608 44. White Tj, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing
609 of fungal ribosomal RNA genes for phylogenetics. In: Innis M, Gelfand D,
610 Shinsky J, White Tj (eds), PCR Protocols: A Guide to Methods and
611 Applications, Academic Press, San Diego, pp. 315-322.
- 612
- 613

614 **Figure captions**

615

616 **Figure 1** Principal component analysis (PCA) of the identified fungal taxa, according to
617 the moth generation from which they were isolated (represented as closed circles; AG -
618 antophagous generation; PG - phyllophagous generation; CG - carpophagous
619 generation). Each number corresponds to a fungal group defined by the coordinates
620 generated by SPSS. Fungal groups are defined as follows: **1** - *C. funicola*, *S. solani*, *P.*
621 *italicum*, *S. vesicarium*, *A. tenuissima*, *T. flavus*, *Alternaria sp.1*, *Penicillium sp.1*, *M.*
622 *hawaiiensis*, *A. ustus*, *Embellisia sp.*, *A. arborescens*, *S. mori*, *M. circinelloides*, *T.*
623 *angustata*, *L. ramosa*, *Penicillium sp.2*, *Alternaria sp.2*; **2** - *P. pinophilum*. **3** -
624 *Arthriniun sp.1*, *Fusarium sp.1*, *T. gamsii*, *F. solani*, *Arthriniun sp.2*, *M. hiemalis*, *B.*
625 *dothidea*, *F. equiseti*, *Arthriniun. sp.3*; **4** - *M. racemosus*, *P. biourgeianum*, *A.*
626 *phaeospermum*, *P. commune*, *P. echinulatum*, *Gibberella sp.*, *E. nigrum*, *C. sinensis*, *B.*
627 *ochroleuca*, *Q. cyanescens*, *M. fragilis*; **5** - *T. roseum*; **6** - *F. oxysporum*; **7** - *B.*
628 *bassiana*. **8** - *C. cladosporioides*.

629

630 **Figure 2** Percentage of fungal isolates (a) and fungal species (b) presenting different
631 ecological roles. Isolates were obtained from dead larvae and pupae of the three
632 generations of *P. oleae* (PG – phyllophagous generation; AG – antophagous generation;
633 CG – carpophagous generation). After molecular identification, fungi were grouped
634 according to their described ecological role (A – antagonistic, E – entomopathogenic, P
635 – phytopathogenic, U – unknown).

636

637 **Table 1** – Larvae and pupae collected from each generation of *P. oleae*. The insect
638 emergence percentage was determined for each moth generation. The percentage of
639 cadavers exhibiting fungal growth on their surface was determined in relation to the
640 total of dead larvae and pupae.

Generation	N° of collected larvae and pupae	Insect emergence	Cadavers presenting surface fungal growth
Phyllophagous	1246	30.1%	61.5%
Antophagous	1745	53.9%	32.7%
Carpophagous	837	19.5%	23.0%
Total	3828	38.6%	40.6%

641

642

643 **Table 2** - Total abundance (N) and relative percentage (%) of each fungal taxa isolated from dead *P. oleae* larvae and pupae in relation to the
 644 total number of identified fungi. Values are presented for all the three moth generations. The ecological role (ER) of each fungus is also
 645 presented (P – phytopathogenic, A – antagonistic, E – entomopathogenic).

646

Family, genera and species	ER	Phyllophagous		Antophagous		Carpophagous		Total	
		N	%	N	%	N	%	N	%
Amphisphaeriaceae									
<i>Truncatella</i>									
<i>T. angustata</i> (Pers.) S. Hughes	P	0	0.0	0	0.0	1	2.3	1	0.6
Apiosporaceae									
<i>Arthrinium</i>									
<i>A. phaeospermum</i> (Corda) M.B. Ellis	A	1	1.4	0	0.0	0	0.0	1	0.6
<i>Arthrinium sp1</i>	A	0	0.0	1	1.9	0	0.0	1	0.6
<i>Arthrinium sp2</i>	A	0	0.0	1	1.9	0	0.0	1	0.6
<i>Arthrinium sp3</i>	A	0	0.0	1	1.9	0	0.0	1	0.6
Bionectriaceae									
<i>Bionectria</i>									
<i>B. ochroleuca</i> (Schwein.) Schroers & Samuels	P, A	2	2.9	0	0.0	0	0.0	2	1.2
Botryosphaeriaceae									
<i>Botryosphaeria</i>									
<i>B. dothidea</i> (Moug.) Ces. & De Not.	P	0	0.0	1	1.9	0	0.0	1	0.6
<i>Microdiplodia</i>									
<i>M. hawaiiensis</i> Crous	P, A	0	0.0	0	0.0	1	2.3	1	0.6
Chaetomiaceae									
<i>Chaetomium</i>									
<i>C. funicola</i> Cooke	P, A	0	0.0	0	0.0	1	2.3	1	0.6

Cordycipitaceae									
<i>Beauveria</i>									
<i>B. bassiana</i> (Bals.-Criv.) Vuill	A, E	49	70.0	0	0.0	1	2.3	50	30.1
<i>Cordyceps</i>									
<i>C. sinensis</i> (Berk.) Sacc.	E	1	1.4	0	0.0	0	0.0	1	0.6
Davidiellaceae									
<i>Cladosporium</i>									
<i>C. cladosporioides</i> (Fresen.) G.A. de Vries	P, A, E	1	1.4	0	0.0	1	2.3	2	1.2
Hypocreaceae									
<i>Trichoderma</i>									
<i>T. gamsii</i> Samuels & Druzhin.	A	0	0.0	8	15.4	0	0.0	8	4.8
Incertae sedis									
<i>Septogloeum</i>									
<i>S. mori</i> (Lév.) Briosi & Cavara	P	0	0.0	0	0.0	1	2.3	1	0.6
<i>Trichothecium</i>									
<i>T. roseum</i> (Pers.) Link	A	1	1.4	2	3.8	0	0.0	3	1.8
Mucoraceae									
<i>Lichtheimia</i>									
<i>L. ramosa</i> (Zopf) Vuill.	-	0	0.0	0	0.0	1	2.3	1	0.6
<i>Mucor</i>									
<i>M. circinelloides</i> Tiegh.	P, A, E	0	0.0	0	0.0	3	6.8	3	1.8
<i>M. fragilis</i> Bainier	P	1	1.4	0	0.0	0	0.0	1	0.6
<i>M. hiemalis</i> Wehmer	P, E	0	0.0	1	1.9	0	0.0	1	0.6
<i>M. racemosus</i> Bull.	P	1	1.4	0	0.0	0	0.0	1	0.6
Nectriaceae									
<i>Fusarium</i>									
<i>F. equiseti</i> (Corda) Sacc.	P, A,	0	0.0	8	15.4	0	0.0	8	4.8
<i>F. oxysporum</i> Schltdl.	P, A, E	2	2.9	23	44.2	0	0.0	25	15.1
<i>F. solani</i> (Mart.) Sacc.	P, E	0	0.0	1	1.9	0	0.0	1	0.6
<i>Fusarium sp. 1</i>	P, A, E	0	0.0	1	1.9	0	0.0	1	0.6

<i>Gibberella</i>									
<i>Gibberella sp. 1</i>	P	1	1.4	0	0.0	0	0.0	1	0.6
Pleosporaceae									
<i>Alternaria</i>									
<i>Alternaria arborescens</i> E. G. Simmons	P	0	0.0	0	0.0	1	2.3	1	0.6
<i>Alternaria sp.1</i>	P, A	3	4.3	3	5.8	6	13.6	12	7.2
<i>Alternaria sp.2</i>	P, A	0	0.0	0	0.0	2	4.5	2	1.2
<i>Alternaria tenuissima</i> (Kunze) Wiltshire	P	0	0.0	0	0.0	5	11.4	5	3.0
<i>Embellisia</i>									
<i>Embellisia sp.</i>	P	0	0.0	0	0.0	1	2.3	1	0.6
<i>Epicoccum</i>									
<i>E. nigrum</i> Link	P, A	1	1.4	0	0.0	0	0.0	1	0.6
<i>Stemphylium</i>									
<i>S. solani</i> G. F. Weber	P	0	0.0	0	0.0	1	2.3	1	0.6
<i>S. vesicarium</i> (Wallr.) E.G. Simmons	P	0	0.0	0	0.0	1	2.3	1	0.6
Quambalariaceae									
<i>Quambalaria</i>									
<i>Q. cyanescens</i> (de Hoog & G.A. de Vries) Z.W. Beer, Begerow & R. Bauer	P	3	4.3	0	0.0	0	0.0	3	1.8
<i>Penicillium</i>									
<i>P. biourgeianum</i> K.M. Zalesky	P	1	1.4	0	0.0	0	0.0	1	0.6
<i>P. commune</i> Thom	P	1	1.4	0	0.0	0	0.0	1	0.6
<i>P. echinulatum</i> Fassat.	P	1	1.4	0	0.0	0	0.0	1	0.6
<i>P. italicum</i> Wehmer	P	0	0.0	0	0.0	1	2.3	1	0.6
<i>P. pinophilum</i> Thom	A	0	0.0	1	1.9	7	15.9	8	4.8
<i>Penicillium sp. 1</i>	P, A, E	0	0.0	0	0.0	1	2.3	1	0.6
<i>Penicillium sp. 2</i>	P, A, E	0	0.0	0	0.0	1	2.3	1	0.6
Trichocomaceae									
<i>Aspergillus</i>									
<i>A. ustus</i> (Bainier) Thom & Church	P, A, E	0	0.0	0	0.0	6	13.6	6	3.6

<i>Talaromyces</i>										
	<i>T. flavus</i> (Klöcker) Stolk & Samson	A	0	0.0	0	0.0	1	2.3	1	0.6
	Total		70	100.0	52	100.0	44	100.0	166	100.0

647

648

649

650

651

652 **Table 3** – *Simpson (D)* and *Shannon-Wiener (H)* diversity indexes of fungal taxa

653 identified in the three generations of *P. oleae*.

Generation	Diversity indexes	
	<i>D</i>	<i>H</i>
Phyllophagous	0.50	1.39
Antophagous	0.25	1.68
Carpophagous	0.09	2.70

654

655