

1  
2  
3 **Title:** *The bed nucleus of stria terminalis and the amygdala as targets of antenatal*  
4 *glucocorticoids: implications for fear and anxiety responses*  
5  
6  
7  
8

9  
10 **Authors:** Mário Oliveira MD<sup>1,2</sup>, Ana-João Rodrigues PhD<sup>1,2</sup>, Pedro Leão MD,PhD<sup>1,2</sup>, Diana  
11 Cardona BSc<sup>1,2</sup>, José Miguel Pêgo MD,PhD<sup>1,2</sup>, Nuno Sousa MD,PhD<sup>1,2</sup>  
12  
13  
14  
15

16  
17 **Affiliations:**  
18

19 <sup>1</sup> Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of  
20 Minho (Braga, Portugal)  
21  
22

23 <sup>2</sup> ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal  
24  
25  
26  
27

28 **Corresponding author:**  
29

30 Nuno Sousa  
31 Life and Health Sciences Research Institute (ICVS), School of Health Sciences  
32 University of Minho  
33 Campus de Gualtar  
34 4710-057 Braga  
35 Portugal  
36 Telephone: +351 253 604 806  
37 Fax: +351 253 604 820  
38 E-mail: njcsousa@eceaude.uminho.pt  
39  
40  
41  
42  
43  
44

45 **Disclosure/conflict of interest:** The authors have no conflicts of interest to declare.  
46  
47  
48  
49

50 **Acknowledgments:** This study was supported by a grant from the Institute for the Study of  
51 Affective Neuroscience (ISAN). AJR is supported by a Fundação para a Ciência e Tecnologia  
52 (FCT) grant.  
53  
54  
55  
56  
57  
58  
59  
60

**ABSTRACT**

*Rationale:* Several human and experimental studies have shown that early life adverse events can shape physical and mental health in adulthood. Stress or elevated levels of glucocorticoids (GCs) during critical periods of development seem to contribute for the appearance of neuropsychiatric conditions such as anxiety and depression, albeit the underlying mechanisms remain to be fully elucidated.

*Objectives:* The aim of the present study was to determine the long-term effect of prenatal exposure to dexamethasone- DEX (synthetic GC widely used in clinics) in fear and anxious behavior and identify the neurochemical, morphological and molecular correlates.

*Results:* Prenatal exposure to DEX triggers a hyperanxious phenotype and altered fear behavior in adulthood. These behavioral traits were correlated with increased volume of the bed nucleus of the stria terminalis (BNST), particularly the anteromedial subdivision which presented increased dendritic length; in parallel, we found an increased expression of synapsin and NCAM in the BNST of these animals. Remarkably, DEX effects were opposite in the amygdala, as this region presented reduced volume due to significant dendritic atrophy. Albeit no differences were found in dopamine and its metabolite levels in the BNST, this neurotransmitter was substantially reduced in the amygdala, which also presented an up-regulation of dopamine receptor 2.

*Conclusions:* Altogether our results show that *in utero* DEX exposure can modulate anxiety and fear behavior in parallel with significant morphological, neurochemical and molecular changes; importantly, GCs seem to differentially affect distinct brain regions involved in this type of behaviors.

**KEYWORDS:** anxiety; fear; amygdala; BNST; stereology; neurodevelopment; corticosteroids

## INTRODUCTION

Since the initial evidence on its benefit in the prevention of respiratory distress syndrome in preterm infants (Liggins and Howie, 1972), antenatal glucocorticoids (GC) have gained wide clinical use in the enhancement of fetal lung maturation in pregnancies at risk of preterm delivery (Crane et al., 2003; Crowley, 1995; NIH, 1995). In fact, accumulating evidence shows a substantial decrease in neonatal morbidity and mortality, which strongly supported the recommendations for its clinical use (Hofmeyr, 2009; NIH, 1995). Additionally, antenatal GCs have been use to treat other conditions such as congenital adrenal hyperplasia (Speiser et al., 2010).

GCs act predominantly through intracellular receptors which control the transcription of GC-responsive target genes (Matthews, 2001); Sousa et al., 2008). While endogenous GCs (corticosterone in rats and cortisone in humans), at basal levels, display higher affinity to the mineralocorticoid receptors (MR), leaving glucocorticoid receptors (GR) receptors largely unoccupied (De Kloet et al., 1998), synthetic GCs such as dexamethasone (DEX) can readily cross the placenta and bind almost exclusively to GR (Reul et al., 1987). GRs can also be activated when endogenous GC secretion is elevated and this chronic activation is noxious for both the periphery and the brain (Sapolsky, 1999). One relevant finding is that despite both being efficacious in preventing respiratory distress syndrome (Crowley, 2000), synthetic GCs effects seem to be more deleterious than endogenous GCs (Oliveira et al., 2006).

While the use of antenatal GCs triggers long-term gains in several clinical conditions, evidence is also revealing the detrimental effects of antenatal exposure to these hormones; in fact, it is now established a correlation between GC exposure in early life with several metabolic (Barker, 1995; Lindsay et al., 1996) and emotional and mood disorders in both animal models and humans in adulthood (Mesquita et al., 2009; Weinstock, 2001). These so-called “programming effects” of GCs may result from endocrine, morphological, neurochemical and molecular changes in specific brain regions; as examples, we have shown that antenatal exposure to DEX significantly changes the mesolimbic circuit (Leao et al., 2007) while others have shown deleterious effects in the prefrontal cortex (Diaz Heijtz et al., 2010), amygdala and hypothalamus (Nagano et al., 2008). Importantly, previous studies from our lab have shown that antenatal exposure to GCs induces a hyperemotional phenotype in

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

adulthood (Oliveira et al., 2006). Given the established the role of the amygdala and bed nucleus of stria terminalis (BNST) in fear and anxiety (Davis, 1992a, b, 1998, 2006), herein we decided to perform an extensive morphological and molecular analysis of these brain regions in adult animals exposed to antenatal DEX and to search for correlations with fear and anxiety behaviors.

## MATERIALS AND METHODS

### Animals and Treatments

Experiments were conducted in accordance with European Union regulations (Directive 86/609/EEC) and National Institutes of Health guidelines on animal care and experimentation.

Fourteen pregnant Wistar rats (Charles River Laboratories, Barcelona, Spain) were individually housed under standard laboratory conditions (room temperature 22°C; 12/12h light/dark cycle; food and water *ad libitum*). Subcutaneous injections of dexamethasone (DEX, 1 mg/kg; N=7) or vehicle (CONT, 1 mL/kg; N=7) were administered on embryonic (ED) 18 and 19 of pregnancy, as previously described (Oliveira et al., 2006; Oliveira et al., 2011). Weaning was carried out at postnatal day 21; male offspring was randomly distributed in groups of two animals per cage according to treatment. Ten males from 4 to 5 different litters of each treatment were tested for behavior at 3 months age.

### Elevated plus maze

The elevated plus maze test, used to assess the degree of anxiousness, was performed on a black polypropylene plus-shaped platform elevated 72.4 cm above the floor (ENV-560; MedAssociates Inc, St Albans, VT, USA). The maze consisted of two open arms (50.8 X 10.2 cm) and two closed arms (50.8 X 10.2 X 40.6 cm). Animals were tested over a period of 5 minutes, under bright white light. The number of entries into each of the arms and the time spent therein were recorded using a system of infrared photobeams. Data were processed to obtain the ratio of time spent in the open arms versus total time and the number of entries into each arm of the maze.

### Acoustic startle as a function of stimulus intensity

Acoustic startle reflex was measured in a startle response system (SR-LAB, San Diego Instruments, San Diego, CA, USA). The apparatus consisted of a non-restrictive Plexiglas cylinder (inner diameter 8.8 cm, length 22.2 cm), mounted on a Plexiglas platform and placed in a ventilated, sound-attenuated chamber. A piezoelectric element detected cylinder

1  
2  
3 movements. Background white noise (intensity 63 dB) was used to minimize the impact of  
4 external acoustic stimuli.  
5

6  
7 Animals were habituated to the apparatus (5 minutes daily) for 2 days before testing. In the  
8 trial day, following 5 minutes acclimatization to the chamber, rats were presented 5 baseline  
9 startle stimuli (50 ms pulse of white noise at 120 dB) at a 30 s inter-stimulus interval, in  
10 order to become familiarized with the startle stimuli. Then, 60 startle stimuli were randomly  
11 presented (50 ms duration and variable intensity between 70 and 120 dB at 10 dB  
12 increments). Startle magnitude was assessed at 1 ms intervals, during the 200 ms period  
13 following stimulus. Chambers were cleaned between tests (70% ethanol and water) in order  
14 to remove olfactory cues.  
15  
16  
17  
18  
19  
20  
21

#### 22 23 24 25 Prepulse inhibition

26  
27 The test lasted 20 minutes, following a 5 minutes acclimatization period. A white background  
28 noise (70 dB) was provided. Following five introductory 120 dB startle trials (noise lasting 40  
29 ms), a total of 35 test trials were pseudo-randomly delivered as follows: a) five trials with  
30 background noise only, b) 10 startle trials of 120 dB and c) five prepulses of each of four  
31 different intensities preceding a startle trial. Prepulse intensities of 2, 4, 8 and 16 dB above  
32 the background noise level lasted 20 ms and preceded the 120 dB startle presentation in 100  
33 ms. Intertrial intervals ranged from 10 to 20 seconds. The average startle response (AVG)  
34 was assessed in the 100 ms period following the onset of the startle stimulus presentation.  
35 The mean AVG for the ten 120dB startle trials was employed in the formula used to assess  
36 the percentage of reduction in AVG compared to startle trials alone: % PPI = 100 X [1-(AVG  
37 at prepulse plus startle trial)/(AVG at startle trial)].  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

#### 50 51 Fear-potentiated startle

52  
53 A stainless steel grid was adapted to the floor of the testing chamber, through which an  
54 electric current could be passed under software control. Following a 5 minutes  
55 acclimatization period, 20 light-shock pairings were presented at 30 seconds intervals. The  
56  
57  
58  
59  
60

1  
2  
3 shock (0.6 mA) was presented during the last 500 ms of the 5 s light pulse (3 watt  
4 incandescent light bulb). Following conditioning, animals were returned to their home cages.  
5  
6

7  
8 On the testing day, the steel grid used during conditioning, was maintained. After 5 minutes  
9 acclimatization, 20 baseline startle stimuli were presented. Startles were measured. Then,  
10 animals were randomly presented 10 startle stimuli (intensity 120 dB, duration 50 ms), five  
11 of them during the last 50 ms of the delivery of a 5 s luminous conditioned stimulus (CS).  
12  
13  
14

#### 15 16 17 18 Histological procedures 19

20 Following behavioral assessment, 8 male animals (derived from 4 to 5 different litters) of  
21 each group were placed under deep pentobarbital anesthesia and transcardially perfused  
22 with either 4% paraformaldehyde solution (N=4) or 0.9% saline (N=4).  
23  
24  
25

26 Brains from the former set of subjects were embedded in glycolmethacrylate (Tecnovit  
27 7100; Heraeus Kulzer, Werheim, Germany) and 30  $\mu$ m coronal sections were obtained by  
28 microtome (Cerqueira et al., 2005). These were placed on a gelatinized slide, stained with  
29 Giemsa, mounted with Entellan (Merck, Darmstadt, Germany) and coverslipped. Shrinkage  
30 factor was calculated according to previous studies (Madeira et al., 1990).  
31  
32  
33  
34

35  
36 Brains from animals perfused with 0.9% saline were processed for Golgi-Cox staining as  
37 previously described (Gibb and Kolb, 1998); details are given in supplementary material.  
38  
39  
40  
41  
42

#### 43 Regional boundaries 44

45 We analyzed stereological parameters in three regions of the amygdaloid complex, namely  
46 the basolateral anterior (BLa), lateral (La) and central (CeA), as previously outlined (Paxinos  
47 and Watson, 2005). CeA and La nuclei were differentiated using prior definitions (De Omlos  
48 et al., 2004) and the BLa division was distinguished from surrounding areas on the basis of  
49 cell size and staining intensity (De Omlos et al., 2004; Krettek and Price, 1978). In accordance  
50 with a recent revision of anatomical and projection studies of the BNST (Dong et al., 2001),  
51 three divisions were considered, the anteromedial (BNSTam), anterolateral (BNSTal) and  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 posterior (BNSTp). In order to avoid observer bias and to permit comparison between  
4 groups, the same observer drew all regional boundaries.  
5  
6  
7  
8

#### 9 10 Stereological procedures

11  
12 Volume and cell number estimations were obtained using Stereoinvestigator® software  
13 (MicroBrightField, Williston, VT, USA) and a motorized microscope (Axioplan 2, Carl Zeiss,  
14 Hamburg, Germany) attached to a camera (DXC-390, Sony Corporation, Tokyo, Japan). The  
15 Cavalieri's principle (Gundersen et al., 1988) was applied to evaluate the volume of each  
16 region. Average cell numbers were estimated using the optical fractionator method (West et  
17 al., 1991). Coefficients of error were computed according to previously published formulas  
18 for cell numbers (Gundersen et al., 1999) and volume estimates (Gundersen and Jensen,  
19 1987). Detailed methods are provided in supplementary material.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

#### 30 Dendritic tree analysis

31  
32 BLA pyramidal-like, CeA multipolar and BNSTam bipolar neurons were chosen as described  
33 elsewhere (McDonald, 1982a, b, 1983). For each selected neuron, all branches of the  
34 dendritic tree were reconstructed at 600x magnification using a motorized microscope  
35 (Axioplan 2, Carl Zeiss), with oil-immersion objectives, and attached to a camera (DXC-390,  
36 Sony Corporation) and NeuroLucida software (MicroBrightField). A 3-D analysis of the  
37 reconstructed neurons was performed using NeuroExplorer software (MicroBrightField).  
38 Dendritic branches were sampled in order to estimate spine density; spines in the selected  
39 segments were classified in thin, wide, ramified and mushroom categories (Harris et al.,  
40 1992). Thin spines were considered immature, while the other spine types were classified as  
41 mature. A total of 20 neurons/group/area were drawn (total 120 neurons).  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

#### 53 Brain catecholamines

54  
55  
56 A different set of males was sacrificed by decapitation (n=6 animals derived from 3 different  
57 litters per group). After brain snap freezing, macrodissection of the amygdala and bed  
58  
59  
60



1  
2  
3 nucleus of stria terminalis was rapidly performed under a stereomicroscope (Model SZX7,  
4 Olympus America Inc., Center Valley, PA, USA). Whole brains were placed upside down and,  
5 using delicate forceps (Dumont #7 forceps, Fine Science Tools USA Inc., Foster City, CA, USA),  
6 the areas of interest were dissected according to stereological coordinates (Paxinos and  
7 Watson, 2005). Samples were frozen in liquid nitrogen (overnight at -20°C) after adding  
8 perchloric acid 0.2M; then, samples were briefly sonicated, centrifuged and 50 µl aliquots of  
9 the supernatant injected on a high performance liquid chromatography (HPLC) combined  
10 with electrochemical detection system. A mobile phase of 0.7 M aqueous potassium  
11 phosphate (monobasic) (pH 3.0) in 10% methanol, 1-heptanesulfonic acid (222 mg/l) and Na-  
12 EDTA (40 mg/l) was used.  
13  
14  
15  
16  
17  
18  
19

20  
21 Levels of dopamine, 3,4-dihydroxyphenylacetic acid (DOPAC) and 4-hydroxy-3-  
22 methoxyphenylacetic acid (homovanillic acid, HVA) were determined using a Gilson  
23 instrument (Gilson Inc., Middleton, WI, USA), fitted with an analytical column (Supelco  
24 Supelcosil LC-18 3 M; 7.5 cm x 4.6 mm; flow rate: 1.0–1.5 ml/min; Supelco, Bellefonte, PA,  
25 USA). A standard curve was then obtained and data presented as concentration (nanogram  
26 per milligram of tissue protein).  
27  
28  
29  
30  
31  
32  
33  
34

### 35 Molecular correlates

36  
37  
38 In order to establish molecular correlates, the mRNA expression of dopamine D1 and D2  
39 receptors (Drd1 and Drd2, respectively) was assessed in the BNST and amygdala. Moreover,  
40 we evaluated the expression of several synaptic plasticity-related genes, including synapsin  
41 (Syn), brain-derived neurotrophic factor (BDNF) and neural cell adhesion molecule (NCAM).  
42 The amygdala and bed nucleus of stria terminalis of an additional set of animals (N=8  
43 animals derived from 4 different litters/group) were isolated as described above. For Real-  
44 Time PCR analysis, total RNA was isolated from frozen areas using Trizol (Invitrogen) and  
45 DNase treated (Fermentas), according to manufacturer. Two µg of RNA were converted into  
46 cDNA using the iSCRIPT kit (Biorad). RT-PCR was performed using SyberGreen (Qiagen) and  
47 the Biorad q-PCR CFX96 apparatus. HPRT was used as a housekeeping gene. We used relative  
48 quantification to determine the fold change difference between control and DEX animals,  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 using the  $\Delta\Delta$ CT method as described before (Pfaffl, 2001). Primer sequences available in  
4 Supplementary Material.  
5  
6

#### 7 Statistical analysis

8

9  
10 Results are presented as mean  $\pm$  SEM. A repeated measures test was used to analyze data  
11 from acoustic startle, prepulse inhibition and fear potentiated startle; Greenhouse-Geisser  
12 and Huynh-Feldt's corrections were applied for acoustic and fear potentiated startle data,  
13 respectively. The comparison of means between groups was performed using the Student's  
14 t-test for the remaining variables. Statistical significance was considered for  $P < 0.05$ .  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RESULTS

### *Brief antenatal exposure to DEX triggers hyperanxiety and impairs fear conditioning*

Animals exposed to either DEX 1mg/kg or vehicle (controls) at embryonic day 18 and 19 were tested at 3 months of age. Antenatal exposure to DEX resulted in a significant reduction on the ratio of time spent in open arm when compared to controls ( $t=3.636$ ,  $P=0.004$ ; Figure 1A); these results confirm our previous findings (Oliveira et al., 2006). Furthermore, while the number of open arm entries was also decreased following DEX exposure ( $t=2.424$ ,  $P=0.026$ ; Figure 1B), there were no significant differences between groups in the number of closed arm entries ( $t=0.340$ ,  $P=0.738$ ), thus showing that the increased anxiety behavior should not be attributed to locomotory differences.

In the acoustic startle test, startle amplitudes for DEX exposed subjects increased more rapidly as a function of stimulus intensity when compared to controls ( $F_{(3,47)}=4.036$ ,  $P=0.016$ ; Table 1). When comparing responsiveness to individual noise intensities between groups, DEX exposed rats displayed significantly increased startle amplitudes at 70 ( $t=5.121$ ,  $P<0.001$ ), 80 ( $t=4.252$ ,  $P=0.001$ ), 90 ( $t=3.537$ ,  $P=0.006$ ), 100 ( $t=3.045$ ,  $P=0.012$ ) and 110 dB ( $t=2.452$ ,  $P=0.025$ ).

Interestingly, the acoustic startle response to inhibitory prepulses was not affected by exposure to antenatal DEX, as shown by the analysis of treatment x prepulse intensity interaction ( $F_{(3,54)}=3.858$ ,  $P=0.994$ ; Figure 2). Moreover, comparison between groups failed to reveal significant differences at any prepulse intensity. The fact that all groups had similar percentages of reduction in the average of startle response, supports the absence of sensorimotor deficits.

Startle amplitude varied as a function of the treatment X stimulus interaction in the fear potentiated startle ( $F_{(1,18)}=8.379$ ;  $P=0.01$ ; Table 2). While controls displayed enhanced startle amplitude after being presented a conditioned stimulus startle, DEX exposure resulted in an

1  
2  
3 impaired fear conditioning in the acoustic startle, as shown by the significantly decreased  
4 ratio of conditioned stimulus/basal startle responsiveness in the DEX group ( $t=-2.602$ ,  
5  $P=0.018$ ; Figure 3).  
6  
7

8  
9  
10  
11 *Antenatal-DEX exposure promotes dendritic remodeling in the BNST and in the amygdala*

12  
13  
14 To further understand the behavioral findings, we have performed a detailed morphological  
15 characterization of the BNST and amygdala brain regions, known to be involved in fear and  
16 anxiety, respectively (Walker and Davis, 1997).  
17  
18

19  
20 Volumetric determinations reveal that antenatal exposure to DEX resulted in an increase in  
21 BNST total volume ( $t=3.841$ ,  $P=0.009$ ; data not shown). This effect was largely due to  
22 increased volumes in anteromedial division of the BNST of DEX-exposed animals ( $t=5.887$ ,  
23  $P=.001$ ; Figure 4A); conversely, the volumes of BNSTal and BNSTp were not influenced by  
24 such antenatal exposure ( $t=-0.484$ ,  $P=.646$  and  $t=0.718$ ,  $P=.500$ , respectively). DEX exposure  
25 during pregnancy did not affect the total number of cells in the BNST ( $t=1.595$ ,  $P=0.162$ ) nor  
26 in its divisions (BNSTam:  $t=1.475$ ,  $P=0.191$ ; BNSTal:  $t=-1.506$ ,  $P=0.183$ ; BNSTpost:  $t=0.946$ ,  
27  $P=0.381$ ; Figure 4B).  
28  
29

30  
31 The volumetric increase in the BNST was further scrutinized through a 3-D morphological  
32 analysis of dendritic arborizations of neurons in the BNSTam; the neurons in this brain region  
33 were previously described as cells with ovoid soma, moderate polarized dendritic branching,  
34 with sparse spines (McDonald, 1983). Data revealed that antenatal exposure to DEX resulted  
35 in a significant increase of total dendritic length when compared to controls ( $t=4.498$ ,  
36  $P<0.001$ ; Table 3). Spine densities were not affected by antenatal DEX treatment ( $t=-1.779$ ,  
37  $P=0.085$ ), even when a separate analysis of mature ( $t=-1.825$ ,  $P=.076$ ) and immature forms  
38 ( $t=-0.534$ ,  $P=.0596$ ) was performed.  
39  
40

41  
42 In contrast to the BNST, antenatal exposure to DEX resulted in a decreased volume of BLa  
43 and CeA divisions of the amygdala (BLa  $t=-3.564$ ,  $P=0.02$ ; CeA  $t=-3.072$ ,  $P=0.027$ ) but not in  
44 the La division ( $t=-0.561$ ,  $P=0.605$ ) (Figure 4C). No effect was observed on the estimated  
45 number of cells in the BLa, CeA and La divisions of the amygdala ( $t=1.457$ ,  $P=0.196$ ;  $t=-1.383$ ,  
46  $P=.216$ ;  $t=-0.725$ ,  $P=0.496$ , respectively; Figure 4D).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The analysis of the dendritic arborizations of pyramidal-like neurons in the BLA region of the  
4 amygdala, which represent the predominant cell-type in the area (70%; type I) (McDonald,  
5 1982b; Sah et al., 2003), revealed a reduction in total and basal dendritic lengths following  
6 DEX treatment ( $t=-2.471$ ,  $P=0.018$  and  $t=-3.252$ ,  $P=0.002$ , respectively; Table 3); however,  
7 apical dendritic length was not affected ( $t=-0.832$ ,  $P=0.411$ ) by antenatal DEX-exposure. The  
8 density of spines in these neurons was not affected by antenatal exposure to DEX ( $t=0.165$ ,  
9  $P=0.870$ ), even when assessing mature and immature spines separately ( $t=-0.122$ ,  $P=0.903$   
10 and  $t=0.598$ ,  $P=0.555$ , respectively).

11  
12 Treatment also affected the CeA predominant cell type, previously called medium spiny  
13 neurons (by comparison with neurons in the striatum) (Sah et al., 2003). These neurons have  
14 an ovoid or fusiform soma and three to five nonspiny primary dendrites from which  
15 moderately spiny, sparsely branching secondary and tertiary dendrites arise. These neurons  
16 in the DEX progeny displayed a significant dendritic atrophy when compared to controls ( $t=-$   
17  $3.491$ ,  $P=0.002$ ; Table 3). No effect was found on the densities of total, mature and  
18 immature spines ( $t=0.435$ ,  $P=0.666$ ,  $t=0.023$ ,  $P=0.982$  and  $t=1.332$ ,  $P=0.191$ , respectively).

### 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 *Neurochemical and molecular correlates*

34  
35 To complement the behavioral and morphometric analysis, we decided to measure  
36 dopamine levels and its metabolites. Antenatal DEX exposure had no effect on the  
37 concentration of dopamine in the BNST, nor its turnover (dopamine  $t=0.026$   $P=0.980$ ;  
38 dopamine turnover  $t=0.291$ ,  $P=0.777$ ; Figure 5A-B).

39  
40 In contrast, the neurochemical analysis in the amygdala revealed a decrease in dopamine  
41 concentration following antenatal exposure to DEX ( $t=5.006$ ,  $P=0.001$ ; Figure 5C). Moreover,  
42 dopamine turnover was significantly increased in these subjects ( $t=-4.405$ ,  $P=0.001$ ; Figure  
43 5D).

44  
45 At a molecular level, and given the impact of antenatal DEX-exposure on dendritic  
46 arborization, we assessed the expression of the synaptic plasticity-related genes in the BNST  
47 and amygdala, namely a synaptic gene (synapsin (Syn)), a neurotrophin (brain-derived  
48 neurotrophic factor (BDNF)) and a cell adhesion molecule (neural cell adhesion molecule  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (NCAM)), all known to be involved in synaptic/dendritic plasticity. Moreover, as the  
4 dopaminergic innervation of the BLA is considered to facilitate amygdala dependent  
5 functions (Asan, 1997), and since a hypodopaminergic status was previously associated with  
6 DEX-exposure (Leao et al., 2007; Rodrigues et al., 2010), the expression of dopamine D1 and  
7 D2 receptors (Drd1 and Drd2, respectively) was also evaluated.  
8  
9

10  
11  
12 In the BNST, DEX exposed animals displayed increased expression of synapsin ( $t=-2.418$ ,  
13  $P=0.030$ ) and NCAM ( $t=-2.032$ ,  $P=0.045$ ) (Table 4). No significant differences between groups  
14 were found in the expression of the remaining genes analysed (Drd1  $t=-0.663$ ,  $P=0.514$ ; Drd2  
15  $t=-0.881$ ,  $P=0.386$ ; BDNF  $t=-0.214$ ,  $P=0.832$ ).  
16  
17  
18

19  
20 In the amygdala, Drd2 mRNA levels were significantly increased in the DEX exposed subjects  
21 ( $t=-3.392$ ,  $P=0.008$ ), while Drd1 were not affected ( $t=0.370$ ,  $P=0.717$ ) (Table 4). The  
22 expression levels of the synaptic plasticity-related genes, namely synapsin, BDNF and NCAM  
23 were not significantly different between groups ( $t=0.546$ ,  $P=0.594$ ,  $t=-1.457$ ,  $P=0.167$  and  $t=-$   
24  $0.376$ ,  $P=0.713$ , respectively).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

*In utero* and early postnatal environment, namely the GC milieu, is crucial for neurodevelopment not only through its activational effects, but also through important programming effects. Increasing evidence suggests that early life exposure to GC triggers undesired metabolic, cardiovascular, neuroendocrine and behavioral phenotypes in adulthood (Mesquita et al., 2009). Both synthetic (such as DEX) and naturally occurring GCs are able to exert a programming effect in the brain; however, it is important to remember that their effects can be quite distinct (Oliveira et al., 2006) as endogenous GCs preferentially bind to MR, whereas DEX binds almost exclusive to GR (Reul et al., 1987), the receptors that mediate most of the detrimental effects of GCs. Although it has been argued that synthetic GCs have a limited capacity to reach the brain in mice (De Kloet et al., 1975), its deleterious effects in the central nervous system are undeniable (Cerqueira et al., 2005; Cerqueira et al., 2007; Leao et al., 2007; McArthur et al., 2005; Yu et al., 2010), specially in rats, and may arise from the use of higher dosages which can indeed pass the barriers and activate brain GC receptors (Miller et al., 1992; Reul et al., 1987). Moreover, in the context of the present study it is important to highlight the fact that the exposure to DEX has occurred during a period in which the brain barriers are still immature (Saunders et al., 2000)."

The data herein presented extends our previous observations that antenatal exposure to DEX results in anxiety-like behavior in adult animals (Oliveira et al., 2006). In fact, while we confirmed a hyperanxious phenotype in the elevated plus maze, we also found an increased acoustic startle response (ASR) in DEX-exposed rats. The startle, a fast protective response from the organism to a sudden and intense stimulus (Koch, 1999), is considered an indicator of anxiety-like behavior. The ASR results from the activation of a trisynaptic circuit, where auditory inputs from several brainstem nuclei (dorsal cochlear nucleus, cochlear root nucleus, ventral cochlear nucleus, lateral superior olive and ventrolateral tegmental nucleus) are conveyed to a main sensorimotor interface, the caudal pontine reticular nucleus, which then projects to spinal motor neurons to trigger the motor response (Koch, 1999; Lee et al., 1996). Importantly, it was demonstrated that a direct projection from the BNST to the caudal pontine reticular nucleus, but also an indirect one via the CeA, are responsible for the modulatory response of stressors upon the ASR (Davis et al., 1997).

1  
2  
3 The BNST is one of the relay stations that conveys inputs from stress-sensitive areas of the  
4 cortex and limbic system both to the HPA axis but also to brain stem nuclei implicated in  
5 several emotional behaviors (Herman and Cullinan, 1997). Here, we show a hypertrophy of  
6 the BNST, due to the enlargement of its anteromedial region. Interestingly, this region  
7 presented alterations in the expression levels of NCAM, a molecule important for neuronal  
8 plasticity (Bisaz and Sandi, 2010; Nacher et al., 2002). Augmented levels of synapsin, a  
9 protein involved in the regulation of neurotransmitter release (Rosahl et al., 1995), also  
10 suggest increased synaptic activity of this brain region in DEX-exposed animals, which is in  
11 line with the increased dendritic arborization of these animals. Such plastic changes are  
12 paralleled with an increased startle response, thus supporting the idea of an overactivation  
13 of this brain subregion. This increased activity of the BNSTam is also in accordance with our  
14 previous observation of a hyperresponsive HPA axis in these subjects (Oliveira et al., 2006).  
15 Importantly, the activational increase of circulating corticosteroids might further promote  
16 the activation of the BNST as it is known that these hormones are implicated in the  
17 neuroanatomical changes observed in this brain region (Pego et al., 2010).

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30 Several projections to the caudal pontine reticular nucleus are considered responsible for  
31 the increase in startle amplitude induced by fear conditioning to a previous neutral stimulus.  
32 These include projections from the CeA, either direct or via mesencephalic reticular  
33 formation and deep layers of superior colliculus, but also indirect projections from the  
34 medial amygdala via ventromedial hypothalamus and ventral periaqueductal gray (Davis,  
35 2006). Previous data supports that while baseline startle amplitudes are not affected by  
36 interference with these connections, the response to fear conditioning is. Interestingly, we  
37 here show that antenatal DEX exposure impairs fear conditioning in adulthood, which is  
38 consistent with deficits in amygdalar function and memory consolidation for emotionally  
39 arousing experiences. These results, together with previous data showing that early life  
40 stress (neonatal isolation) resulted in impairment of context-induced fear conditioning in  
41 adult male rats (Kosten et al., 2006), confirms that fear conditioning can be modulated by  
42 adverse early life events. The role of different divisions of the amygdala in fear behavior, in  
43 particular the CeA and BLA, has been extensively studied. Several forms of CeA lesions have  
44 been correlated with disruption of fear-potentiated startle (Campeau and Davis, 1995;  
45 Hitchcock and Davis, 1987; Walker and Davis, 1997). Electrolytic lesions of the CeA  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 completely blocked the expression of fear-potentiated startle in rats (Kim and Davis, 1993).  
4 Conversely, inactivation of BNST did not disrupt fear-potentiated startle (Gewirtz et al.,  
5 1998; Walker and Davis, 1997); however, more recently, it has been suggested that BNST lat  
6 tonically inhibits fear-potentiated startle, probably through projections to the CeA (Meloni  
7 et al., 2006). Thus, the present observations of decreased volumes in the CeA and BLA in  
8 DEX-exposed rats are likely to be implicated in the changes in fear conditioning displayed by  
9 these animals; once again, the volumetric decreases in these amygdalar divisions result  
10 largely from dendritic atrophy-which fits previous observations of the effects of chronic  
11 unpredictable stress in the BLA (but not the CeA) (Vyas et al., 2003; Vyas et al., 2002). It is  
12 relevant to stress at this point that others have shown that a single course of  
13 betamethasone at postnatal day one failed to affect volumes of cerebral cortex, corpus  
14 callosum, hippocampus, dentate gyrus or amygdala (Yossuck et al., 2006), however, several  
15 technical differences may explain this discrepancy. The time of the exposure and the nature  
16 of insults seems to be critical issues to consider, as several windows of vulnerability to the  
17 programming effects of different stimuli seem to occur. Indeed, stress effects in neuronal  
18 morphology of specific brain regions are not necessarily equal across life and depend on the  
19 type of stress. While stress effects in the BNST morphology seem more similar, since  
20 different types of stress induce a general hypertrophy of dendrites in the neurons of this  
21 brain region (Pego et al., 2008; Vyas et al., 2003), the effects of stress in the amygdala are  
22 less concordant to our present observation of dendritic atrophy. Whereas chronic juvenile  
23 stress leads to a general hypertrophy of amygdalar dendrites (Eiland et al., 2011), others  
24 failed to find any significant change in dendritic structure after stress exposure in adulthood  
25 (Pego et al., 2008) and others have shown remarkably divergent changes in amygdalar  
26 neurons following different types of stress (Vyas et al., 2003; Vyas et al., 2002).

27  
28 Remarkably, such morphological effects in the amygdala were accompanied by changes in  
29 dopamine but not other catecholamines levels. Dopamine, arising mainly from the VTA,  
30 plays a facilitative role in the function of the amygdala (Asan, 1997). *In vitro* studies also  
31 suggest that the excitability of amygdalar, namely BLA, neurons is modulated by dopamine  
32 (Kroner et al., 2005). In addition, behavioral data showed that VTA lesion results in blocked  
33 fear-potentiated startle (Borowski and Kokkinidis, 1996). At least in part, the facilitation of  
34 affective behaviors by DA may be explained by actions at the cellular level on BLA neurons. It  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 has been suggested that DA receptor activation could simultaneously facilitate the BLA  
4 output in response to strong inputs that cause spike firing, while suppressing weaker inputs  
5 via activation of GABAergic interneurons (Kroner et al., 2005). Thus, the present observation  
6 of a hypodopaminergic status in the amygdala seems to be of relevance to the deficits in  
7 fear memory. In fact, it confirms observations in models of dopamine deficient mice (Fadok  
8 et al., 2010), in which the restoration of dopamine in the BLA was shown to be required for  
9 the formation of fear-related memory (Fadok et al., 2010). Dopamine levels in the amygdala  
10 are also determinant for sensorimotor gating, as it was shown that NAcc and amygdalar  
11 infusions of this neurotransmitter significantly impair PPI (Swerdlow et al., 1992);  
12 surprisingly, we did not find any differences in PPI in DEX-exposed animals, which is also in  
13 accordance with another previous study (Hauser et al., 2006).  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 Herein, the decreased amygdalar levels of dopamine are associated with increased  
24 expression of D2 receptor mRNA in DEX exposed subjects; such up-regulation could be seen  
25 as a compensation mechanism. Remarkably, no differences were found in D1 receptor  
26 expression, suggesting that D2 receptor has a prominent role in fear behavior. Indeed, while  
27 D1 receptors may participate in recognition of danger, D2 receptors seem to have a role in  
28 setting up adaptive responses to adverse stimuli (de la Mora et al., 2010). Moreover, D2 (but  
29 not D1) antagonists injection in the BLA impairs fear-potentiated startle probably due to  
30 reduced dopaminergic tone (de Oliveira et al., 2011), which is in agreement with the  
31 amygdalar hypodopaminergic status seen in DEX-exposed animals. These findings are  
32 consistent with recent data showing that D2 receptor pathway connecting the VTA and BLA  
33 modulates conditioned fear (de Oliveira et al., 2011).  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The present findings unravel the impact of antenatal exposure to DEX in the brain regions  
44 implicated in fear and anxiety behaviors. We show that subjects exposed to GCs during  
45 neurodevelopment present marked neuroanatomical, neurochemical and molecular  
46 programming changes in the BNST and amygdala, and are more vulnerable to anxiety and  
47 fear pathology in adulthood. Thus, our findings further support the concerns raised on the  
48 potential deleterious effects of antenatal exposure to synthetic corticosteroids (Weinstock  
49 et al., 2005; Talge et al., 2007; Mesquita et al., 2009) and call for the need of a parsimonious  
50 use of these drugs.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

- 1  
2  
3  
4  
5 Asan, E. (1997). Ultrastructural features of tyrosine-hydroxylase-immunoreactive afferents and their targets in the rat  
6 amygdala. *Cell Tissue Res* 288, 449-469.
- 7  
8 Barker, D.J. (1995). Fetal origins of coronary heart disease. *BMJ (Clinical research ed)* 311, 171-174.
- 9 Bisaz, R., and Sandi, C. (2010). The role of NCAM in auditory fear conditioning and its modulation by stress: a focus on the  
10 amygdala. *Genes Brain Behav* 9, 353-364.
- 11 Borowski, T.B., and Kokkinidis, L. (1996). Contribution of ventral tegmental area dopamine neurons to expression of  
12 conditional fear: effects of electrical stimulation, excitotoxin lesions, and quinpirole infusion on potentiated startle in rats.  
13 *Behavioral neuroscience* 110, 1349-1364.
- 14 Campeau, S., and Davis, M. (1995). Involvement of the central nucleus and basolateral complex of the amygdala in fear  
15 conditioning measured with fear-potentiated startle in rats trained concurrently with auditory and visual conditioned  
16 stimuli. *J Neurosci* 15, 2301-2311.
- 17 Cerqueira, J.J., Pego, J.M., Taipa, R., Bessa, J.M., Almeida, O.F., and Sousa, N. (2005). Morphological correlates of  
18 corticosteroid-induced changes in prefrontal cortex-dependent behaviors. *J Neurosci* 25, 7792-7800.
- 19 Cerqueira, J.J., Taipa, R., Uylings, H.B., Almeida, O.F., and Sousa, N. (2007). Specific configuration of dendritic degeneration  
20 in pyramidal neurons of the medial prefrontal cortex induced by differing corticosteroid regimens. *Cereb Cortex* 17, 1998-  
21 2006.
- 22 Crane, J., Armonson, A., Brunner, M., De La Ronde, S., Farine, D., Keenan-Lindsay, L., Leduc, L., Schneider, C., and Van Aerde, J.  
23 (2003). Antenatal corticosteroid therapy for fetal maturation. *J Obstet Gynaecol Can* 25, 45-52.
- 24 Crowley, P. (2000). Prophylactic corticosteroids for preterm birth. *Cochrane Database Syst Rev*, CD000065.
- 25 Crowley, P.A. (1995). Antenatal corticosteroid therapy: a meta-analysis of the randomized trials, 1972 to 1994. *American*  
26 *journal of obstetrics and gynecology* 173, 322-335.
- 27  
28 Davis, M. (1992a). The role of the amygdala in fear and anxiety. *Annu Rev Neurosci* 15, 353-375.
- 29 Davis, M. (1992b). The role of the amygdala in fear-potentiated startle: implications for animal models of anxiety. *Trends*  
30 *Pharmacol Sci* 13, 35-41.
- 31 Davis, M. (1998). Are different parts of the extended amygdala involved in fear versus anxiety? *Biological psychiatry* 44,  
32 1239-1247.
- 33 Davis, M. (2006). Neural systems involved in fear and anxiety measured with fear-potentiated startle. *The American*  
34 *psychologist* 61, 741-756.
- 35 Davis, M., Walker, D.L., and Lee, Y. (1997). Amygdala and bed nucleus of the stria terminalis: differential roles in fear and  
36 anxiety measured with the acoustic startle reflex. *Philos Trans R Soc Lond B Biol Sci* 352, 1675-1687.
- 37 De Kloet, E.R., Vreugdenhil, E., Oitzl, M.S., and Joels, M. (1998). Brain corticosteroid receptor balance in health and disease.  
38 *Endocrine reviews* 19, 269-301.
- 39 De Kloet, R., Wallach, G., and McEwen, B.S. (1975). Differences in corticosterone and dexamethasone binding to rat brain  
40 and pituitary. *Endocrinology* 96, 598-609.
- 41 de la Mora, M.P., Gallegos-Cari, A., Arizmendi-Garcia, Y., Marcellino, D., and Fuxe, K. (2010). Role of dopamine receptor  
42 mechanisms in the amygdaloid modulation of fear and anxiety: Structural and functional analysis. *Prog Neurobiol* 90, 198-  
43 216.
- 44 de Oliveira, A.R., Reimer, A.E., de Macedo, C.E., de Carvalho, M.C., Silva, M.A., and Brandao, M.L. (2011). Conditioned fear is  
45 modulated by D2 receptor pathway connecting the ventral tegmental area and basolateral amygdala. *Neurobiol Learn Mem*  
46 95, 37-45.
- 47 De Omlos, J.S., Beltramino, C.A., and Alheid, G. (2004). Amygdala and extended amygdala of the rat: a cytoarchitectonical,  
48 fibroarchitectonical and chemoarchitectonical survey. In *The Rat Nervous System*, G. Paxinos, ed. (Amsterdam, Elsevier).
- 49 Diaz Heijtz, R., Fuchs, E., Feldon, J., Pryce, C.R., and Forsberg, H. (2010). Effects of antenatal dexamethasone treatment on  
50 glucocorticoid receptor and calcyon gene expression in the prefrontal cortex of neonatal and adult common marmoset  
51 monkeys. *Behav Brain Funct* 6, 18.
- 52 Dong, H.W., Petrovich, G.D., and Swanson, L.W. (2001). Topography of projections from amygdala to bed nuclei of the stria  
53 terminalis. *Brain Res Brain Res Rev* 38, 192-246.
- 54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Eiland, L., Ramroop, J., Hill, M.N., Manley, J., and McEwen, B.S. (2011). Chronic juvenile stress produces corticolimbic  
4 dendritic architectural remodeling and modulates emotional behavior in male and female rats. *Psychoneuroendocrinology*.  
5  
6 Fadok, J.P., Darvas, M., Dickerson, T.M., and Palmiter, R.D. (2010). Long-term memory for pavlovian fear conditioning  
7 requires dopamine in the nucleus accumbens and basolateral amygdala. *PLoS One* 5, e12751.  
8  
9 Gewirtz, J.C., McNish, K.A., and Davis, M. (1998). Lesions of the bed nucleus of the stria terminalis block sensitization of the  
10 acoustic startle reflex produced by repeated stress, but not fear-potentiated startle. *Progress in neuro-psychopharmacology*  
11 & biological psychiatry 22, 625-648.  
12  
13 Gibb, R., and Kolb, B. (1998). A method for vibratome sectioning of Golgi-Cox stained whole rat brain. *J Neurosci Methods*  
14 79, 1-4.  
15  
16 Gundersen, H.J., Bendtsen, T.F., Korbo, L., Marcussen, N., Moller, A., Nielsen, K., Nyengaard, J.R., Pakkenberg, B., Sorensen,  
17 F.B., Vesterby, A., *et al.* (1988). Some new, simple and efficient stereological methods and their use in pathological research  
18 and diagnosis. *APMIS* 96, 379-394.  
19  
20 Gundersen, H.J., and Jensen, E.B. (1987). The efficiency of systematic sampling in stereology and its prediction. *J Microsc*  
21 147, 229-263.  
22  
23 Gundersen, H.J., Jensen, E.B., Kieu, K., and Nielsen, J. (1999). The efficiency of systematic sampling in stereology--  
24 reconsidered. *J Microsc* 193, 199-211.  
25  
26 Harris, K.M., Jensen, F.E., and Tsao, B. (1992). Three-dimensional structure of dendritic spines and synapses in rat  
27 hippocampus (CA1) at postnatal day 15 and adult ages: implications for the maturation of synaptic physiology and long-  
28 term potentiation. *J Neurosci* 12, 2685-2705.  
29  
30 Hauser, J., Feldon, J., and Pryce, C.R. (2006). Prenatal dexamethasone exposure, postnatal development, and adulthood  
31 prepulse inhibition and latent inhibition in Wistar rats. *Behav Brain Res* 175, 51-61.  
32  
33 Herman, J.P., and Cullinan, W.E. (1997). Neurocircuitry of stress: central control of the hypothalamo-pituitary-  
34 adrenocortical axis. *Trends Neurosci* 20, 78-84.  
35  
36 Hitchcock, J.M., and Davis, M. (1987). Fear-potentiated startle using an auditory conditioned stimulus: effect of lesions of  
37 the amygdala. *Physiol Behav* 39, 403-408.  
38  
39 Hofmeyr, G.J. (2009). Antenatal corticosteroids for women at risk of preterm birth: RHL commentary. The WHO  
40 Reproductive Health Library; Geneva: World Health Organization.  
41  
42 Kim, M., and Davis, M. (1993). Lack of a temporal gradient of retrograde amnesia in rats with amygdala lesions assessed  
43 with the fear-potentiated startle paradigm. *Behavioral neuroscience* 107, 1088-1092.  
44  
45 Koch, M. (1999). The neurobiology of startle. *Progress in neurobiology* 59, 107-128.  
46  
47 Kosten, T.A., Lee, H.J., and Kim, J.J. (2006). Early life stress impairs fear conditioning in adult male and female rats. *Brain Res*  
48 1087, 142-150.  
49  
50 Krettek, J.E., and Price, J.L. (1978). A description of the amygdaloid complex in the rat and cat with observations on intra-  
51 amygdaloid axonal connections. *The Journal of comparative neurology* 178, 255-280.  
52  
53 Kroner, S., Rosenkranz, J.A., Grace, A.A., and Barrionuevo, G. (2005). Dopamine modulates excitability of basolateral  
54 amygdala neurons in vitro. *Journal of neurophysiology* 93, 1598-1610.  
55  
56 Leao, P., Sousa, J.C., Oliveira, M., Silva, R., Almeida, O.F., and Sousa, N. (2007). Programming effects of antenatal  
57 dexamethasone in the developing mesolimbic pathways. *Synapse* 61, 40-49.  
58  
59 Lee, Y., Lopez, D.E., Meloni, E.G., and Davis, M. (1996). A primary acoustic startle pathway: obligatory role of cochlear root  
60 neurons and the nucleus reticularis pontis caudalis. *J Neurosci* 16, 3775-3789.  
61  
62 Liggins, G.C., and Howie, R.N. (1972). A controlled trial of antepartum glucocorticoid treatment for prevention of the  
63 respiratory distress syndrome in premature infants. *Pediatrics* 50, 515-525.  
64  
65 Lindsay, R.S., Lindsay, R.M., Waddell, B.J., and Seckl, J.R. (1996). Prenatal glucocorticoid exposure leads to offspring  
66 hyperglycaemia in the rat: studies with the 11 beta-hydroxysteroid dehydrogenase inhibitor carbenoxolone. *Diabetologia*  
67 39, 1299-1305.  
68  
69 Madeira, M.D., Pereira, A., Cadete-Leite, A., and Paula-Barbosa, M.M. (1990). Estimates of volumes and pyramidal cell  
70 numbers in the prelimbic subarea of the prefrontal cortex in experimental hypothyroid rats. *J Anat* 171, 41-56.  
71  
72 Matthews, S.G. (2001). Antenatal glucocorticoids and the developing brain: mechanisms of action. *Semin Neonatol* 6, 309-  
73 317.  
74  
75 McArthur, S., McHale, E., Dalley, J.W., Buckingham, J.C., and Gillies, G.E. (2005). Altered mesencephalic dopaminergic  
76 populations in adulthood as a consequence of brief perinatal glucocorticoid exposure. *J Neuroendocrinol* 17, 475-482.

- 1  
2  
3 McDonald, A.J. (1982a). Cytoarchitecture of the central amygdaloid nucleus of the rat. *The Journal of comparative neurology* 208, 401-418.
- 4  
5 McDonald, A.J. (1982b). Neurons of the lateral and basolateral amygdaloid nuclei: a Golgi study in the rat. *The Journal of comparative neurology* 212, 293-312.
- 6  
7 McDonald, A.J. (1983). Neurons of the bed nucleus of the stria terminalis: a golgi study in the rat. *Brain Res Bull* 10, 111-120.
- 8  
9  
10 Meloni, E.G., Jackson, A., Gerety, L.P., Cohen, B.M., and Carlezon, W.A., Jr. (2006). Role of the bed nucleus of the stria terminalis (BST) in the expression of conditioned fear. *Ann N Y Acad Sci* 1071, 538-541.
- 11  
12 Mesquita, A.R., Wegerich, Y., Patchev, A.V., Oliveira, M., Leao, P., Sousa, N., and Almeida, O.F. (2009). Glucocorticoids and neuro- and behavioural development. *Seminars in fetal & neonatal medicine* 14, 130-135.
- 13  
14 Miller, A.H., Spencer, R.L., Pulera, M., Kang, S., McEwen, B.S., and Stein, M. (1992). Adrenal steroid receptor activation in rat brain and pituitary following dexamethasone: implications for the dexamethasone suppression test. *Biol Psychiatry* 32, 850-869.
- 15  
16  
17 Nacher, J., Lanuza, E., and McEwen, B.S. (2002). Distribution of PSA-NCAM expression in the amygdala of the adult rat. *Neuroscience* 113, 479-484.
- 18  
19 Nagano, M., Ozawa, H., and Suzuki, H. (2008). Prenatal dexamethasone exposure affects anxiety-like behaviour and neuroendocrine systems in an age-dependent manner. *Neurosci Res* 60, 364-371.
- 20  
21 NIH (1995). Effect of corticosteroids for fetal maturation on perinatal outcomes. NIH Consensus Development Panel on the Effect of Corticosteroids for Fetal Maturation on Perinatal Outcomes. *Jama* 273, 413-418.
- 22  
23  
24 Oliveira, M., Bessa, J.M., Mesquita, A., Tavares, H., Carvalho, A., Silva, R., Pego, J.M., Cerqueira, J.J., Palha, J.A., Almeida, O.F., *et al.* (2006). Induction of a hyperanxious state by antenatal dexamethasone: a case for less detrimental natural corticosteroids. *Biol Psychiatry* 59, 844-852.
- 25  
26  
27 Oliveira, M., Leao, P., Rodrigues, A.J., Pego, J.M., Cerqueira, J.J., and Sousa, N. (2011). Programming Effects of Antenatal Corticosteroids Exposure in Male Sexual Behavior. *J Sex Med*.
- 28  
29 Paxinos, G., and Watson, C. (2005). *The rat brain in stereotaxic coordinates* (San Diego, Academic Press).
- 30  
31 Pego, J.M., Morgado, P., Pinto, L.G., Cerqueira, J.J., Almeida, O.F., and Sousa, N. (2008). Dissociation of the morphological correlates of stress-induced anxiety and fear. *Eur J Neurosci* 27, 1503-1516.
- 32  
33 Pego, J.M., Sousa, J.C., Almeida, O.F., and Sousa, N. (2010). Stress and the neuroendocrinology of anxiety disorders. *Curr Top Behav Neurosci* 2, 97-117.
- 34  
35 Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research* 29, e45.
- 36  
37 Reul, J.M., van den Bosch, F.R., and de Kloet, E.R. (1987). Relative occupation of type-I and type-II corticosteroid receptors in rat brain following stress and dexamethasone treatment: functional implications. *J Endocrinol* 115, 459-467.
- 38  
39 Rodrigues, A.J., Leao, P., Oliveira, M., Pego, J.M., Franky, A., Morgado, P., Mesquita, A., Araujo, D., Almeida, O.F., and Sousa, N. (2010). Vulnerability to addiction can be modulated by dopamine. Program No 497-12, 2010 Neuroscience Meeting Planner, San Diego, CA: Society for Neuroscience *Online*.
- 40  
41 Rosahl, T.W., Spillane, D., Missler, M., Herz, J., Selig, D.K., Wolff, J.R., Hammer, R.E., Malenka, R.C., and Sudhof, T.C. (1995). Essential functions of synapsins I and II in synaptic vesicle regulation. *Nature* 375, 488-493.
- 42  
43 Sah, P., Faber, E.S., Lopez De Armentia, M., and Power, J. (2003). The amygdaloid complex: anatomy and physiology. *Physiol Rev* 83, 803-834.
- 44  
45 Sapolsky, R.M. (1999). Glucocorticoids, stress, and their adverse neurological effects: relevance to aging. *Exp Gerontol* 34, 721-732.
- 46  
47 Saunders, N.R., Knott, G.W., and Dziegielewska, K.M. (2000). Barriers in the immature brain. *Cell Mol Neurobiol* 20, 29-40.
- 48  
49 Speiser, P.W., Azziz, R., Baskin, L.S., Ghizzoni, L., Hensle, T.W., Merke, D.P., Meyer-Bahlburg, H.F., Miller, W.L., Montori, V.M., Oberfield, S.E., *et al.* (2010). Congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency: an Endocrine Society clinical practice guideline. *The Journal of clinical endocrinology and metabolism* 95, 4133-4160.
- 50  
51 Swerdlow, N.R., Caine, S.B., and Geyer, M.A. (1992). Regionally selective effects of intracerebral dopamine infusion on sensorimotor gating of the startle reflex in rats. *Psychopharmacology (Berl)* 108, 189-195.
- 52  
53 Vyas, A., Bernal, S., and Chattarji, S. (2003). Effects of chronic stress on dendritic arborization in the central and extended amygdala. *Brain Res* 965, 290-294.
- 54  
55  
56  
57  
58  
59  
60

1  
2  
3 Vyas, A., Mitra, R., Shankaranarayana Rao, B.S., and Chattarji, S. (2002). Chronic stress induces contrasting patterns of  
4 dendritic remodeling in hippocampal and amygdaloid neurons. *J Neurosci* 22, 6810-6818.

5 Walker, D.L., and Davis, M. (1997). Double dissociation between the involvement of the bed nucleus of the stria terminalis  
6 and the central nucleus of the amygdala in startle increases produced by conditioned versus unconditioned fear. *J Neurosci*  
7 17, 9375-9383.

8 Weinstock, M. (2001). Alterations induced by gestational stress in brain morphology and behaviour of the offspring.  
9 *Progress in neurobiology* 65, 427-451.

10 West, M.J., Slomianka, L., and Gundersen, H.J. (1991). Unbiased stereological estimation of the total number of neurons in  
11 the subdivisions of the rat hippocampus using the optical fractionator. *Anat Rec* 231, 482-497.

12 Yossuck, P., Kraszpulski, M., and Salm, A.K. (2006). Perinatal corticosteroid effect on amygdala and hippocampus volume  
13 during brain development in the rat model. *Early human development* 82, 267-272.

14 Yu, S., Patchev, A.V., Wu, Y., Lu, J., Holsboer, F., Zhang, J.Z., Sousa, N., and Almeida, O.F. (2010). Depletion of the neural  
15 precursor cell pool by glucocorticoids. *Ann Neurol* 67, 21-30.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## TABLES

Noise (dB)	Controls	DEX
<b>70 dB</b>	30,51 ± 3,03	<b>97,50 ± 15,07 *</b>
<b>80 dB</b>	32,50 ± 4,72	<b>291,17 ± 86,56 *</b>
<b>90 dB</b>	145,19 ± 21,39	<b>561,26 ± 115,66 *</b>
<b>100 dB</b>	1203,54 ± 168,88	<b>3126,46 ± 608,44 *</b>
<b>110 dB</b>	4645,30 ± 461,93	<b>6221,83 ± 447,09 *</b>
<b>120 dB</b>	6358,69 ± 257,17	7101,62 ± 519,47

Table 1 – Acoustic startle data. Startle amplitude (arbitrary units) in response to acoustic stimulus. Data presented as mean ± SEM. \* different from controls, P<.05.

	Controls	DEX
<b>Startle</b>	5290.89 ± 658.66	5252.08 ± 716.75
<b>CS + Startle</b>	6229.04 ± 425.83	<b>4554.13 ± 577.61*</b>

Table 2 – Fear-potentiated acoustic startle data. Startle amplitude (arbitrary units) in response to acoustic stimulus with or without previous conditioned stimulus (CS). Data presented as mean ± SEM. \* different from controls, P<.05.



	Controls	DEX
<b>BNSTam (bipolar)</b>		
Dendritic length ( $\mu\text{m}$ )	395 $\pm$ 17	<b>511 <math>\pm</math> 19*</b>
Density of spines (n/ $\mu\text{m}$ )	0.58 $\pm$ 0.042	0.49 $\pm$ 0.027
Mature	0.47 $\pm$ 0.035	0.39 $\pm$ 0.026
Immature	0.10 $\pm$ 0.015	0.09 $\pm$ 0.016
<b>BLa (pyramidal)</b>		
Dendritic length ( $\mu\text{m}$ )	1969 $\pm$ 128	<b>1506 <math>\pm</math> 137*</b>
Basal dendrites	1088 $\pm$ 87	<b>721 <math>\pm</math> 72*</b>
Apical dendrite	881 $\pm$ 80	785 $\pm$ 83
Density of spines (n/ $\mu\text{m}$ )	0.95 $\pm$ 0.046	0.95 $\pm$ 0.027
Mature spines	0.71 $\pm$ 0.037	0.70 $\pm$ 0.026
Immature spines	0.24 $\pm$ 0.022	0.26 $\pm$ 0.012
<b>CeA</b>		
Dendritic length ( $\mu\text{m}$ )	818 $\pm$ 38	<b>670 <math>\pm</math> 19*</b>
Density of spines (n/ $\mu\text{m}$ )	0.45 $\pm$ 0.043	0.47 $\pm$ 0.037
Mature	0.35 $\pm$ 0.039	0.36 $\pm$ 0.032
Immature	0.09 $\pm$ 0.009	0.12 $\pm$ 0.015

Table 3 – Dendritic length and spine density of bipolar neurons of anteromedial region of the bed nucleus of stria terminalis (BNSTam), pyramidal-like neurons of basolateral amygdaloid nucleus (BLa) and multipolar neurons of the central nucleus of the amygdala (CeA) and Data presented as mean  $\pm$  SEM. \* different from controls,  $P < .05$ .

Treatment	Drd1	Drd2	Syn	BDNF	NCAM
<b>BNST</b>					
Controls	0.950 ± 0.12	1.070 ± 0.12	0.918 ± 0.05	1.127 ± 0.14	0.987 ± 0.04
DEX	1.229 ± 0.21	1.226 ± 0.13	<b>1.200 ± 0.11*</b>	1.196 ± 0.27	<b>1.318 ± 0.25*</b>
<b>Amygdala</b>					
Controls	1.066 ± 0.13	0.927 ± 0.07	1.167 ± 0.19	0.920 ± 0.04	1.043 ± 0.10
DEX	0.990 ± 0.16	<b>2.494 ± 0.62*</b>	1.060 ± 0.04	1.102 ± 0.12	1.115 ± 0.17

Table 4 - Gene expression analysis (measured by real-time PCR) in amygdala and bed nucleus of stria terminalis (BNST). The mRNA expression levels are presented as the fold change increase in relation to the respective control. Data presented as mean ± SEM. \* different from controls, P<.05. Dopamine D1 receptor (Drd1), dopamine D2 receptor (Drd2), synapsin (Syn), brain-derived neurotrophic factor (BDNF) and neural cell adhesion molecule (NCAM).

**LEGENDS TO FIGURES**

Figure 1 – Elevated plus maze data. (A) Ratio of time spent in open arm over total time. (B) Number of open and closed arm entries. Data presented as mean  $\pm$  SEM. \* different from controls,  $P < .05$ .

Figure 2 – Prepulse inhibition test. Percentages of reduction in average startle response compared with startle trials at prepulse intensities of 2, 4, 8 and 16 dB above background noise level.

Figure 3 – Fear potentiated acoustic startle data. Data presented as the average of ratios between conditioned stimulus/basal startle responsiveness  $\pm$  SEM. \* different from controls,  $P < .05$ .

Figure 4 – (A) Estimated volumes of anterolateral (BNSTal), anteromedial (BNSTam) and posterior (BNSTpost) divisions of the bed nucleus of stria terminalis. Data presented in  $\text{mm}^3$ , as mean  $\pm$  SEM. (B) Estimated total number of cells in the anterolateral (BNSTal), anteromedial (BNSTam) and posterior (BNSTpost) divisions of the bed nucleus of stria terminalis. (C) Estimated volumes of BLA, CeA and La divisions of the amygdala. Data presented in  $\text{mm}^3$ , as mean  $\pm$  SEM. (D) Estimated total number of cells in the BLA, CeA and La divisions of the amygdala. Data presented as mean  $\pm$  SEM; \* different from controls,  $P < .05$

Figure 5 – Neurochemical analysis by HPLC of the bed nucleus of stria terminalis (BNST) (A-B) and the amygdala (C-D). (A, C) Concentration of each neurotransmitter (ng/mg of protein) is presented. 3,4-dihydroxyphenylacetic acid (DOPAC) and 4-hydroxy-3-methoxyphenylacetic acid (homovanillic acid, HVA). (B, D) Dopamine turnover assessed by the ratio between dopamine metabolites and dopamine concentrations. Data presented as mean  $\pm$  SEM.