

## Metadata of the chapter that will be visualized online

Chapter Title	Detection of the Glucocorticoid Receptors in Brain Protein Extracts by SDS-PAGE	
Copyright Year	2014	
Copyright Holder	Springer Science+Business Media New York	
Author	Family Name	<b>Marques</b>
	Particle	
	Given Name	<b>Fernanda</b>
	Suffix	
	Division	Life and Health Sciences Research Institute (ICVS), School of Health Sciences
	Organization	University of Minho
	Address	Campus Gualtar, Braga, 4710-057, Portugal
Author	Family Name	<b>Sousa</b>
	Particle	
	Given Name	<b>João C.</b>
	Suffix	
	Division	Life and Health Sciences Research Institute (ICVS), School of Health Sciences
	Organization	University of Minho
	Address	Campus Gualtar, Braga, 4710-057, Portugal
Author	Family Name	<b>Cerqueira</b>
	Particle	
	Given Name	<b>João J.</b>
	Suffix	
	Division	Life and Health Sciences Research Institute (ICVS), School of Health Sciences
	Organization	University of Minho
	Address	Campus Gualtar, Braga, 4710-057, Portugal
Corresponding Author	Family Name	<b>Sousa</b>
	Particle	
	Given Name	<b>Nuno</b>
	Suffix	

Division Life and Health Sciences Research  
Institute (ICVS), School of Health  
Sciences

Organization University of Minho

Address Campus Gualtar, Braga, 4710-057,  
Portugal

Email njcsousa@ecsau.de.uminho.pt

---

**Abstract**

Glucocorticoids are steroid hormones vital for organ system homeostasis and for the maintenance of essential biological processes. A significant part of these actions are mediated through glucocorticoid receptor (GR) that belongs to the nuclear receptor superfamily. To cover such variety of processes the different glucocorticoids act through different GR isoforms that are originated due to posttranscriptional and posttranslational mechanisms. For this reason when evaluating the levels of GRs we should preferentially determine protein levels instead of gene expression. Here, we describe the detection by Western blotting of the GR ( $\alpha$  and  $\beta$  isoforms) protein, using macrodissected brain tissue.

---

**Key words**  
(separated by “-”)

Glucocorticoid receptor - Brain - Protein levels - Western blot

---

## Detection of the Glucocorticoid Receptors in Brain Protein Extracts by SDS-PAGE 2 3

Fernanda Marques, João C. Sousa, João J. Cerqueira, and Nuno Sousa 4

### Abstract 5

Glucocorticoids are steroid hormones vital for organ system homeostasis and for the maintenance of essential biological processes. A significant part of these actions are mediated through glucocorticoid receptor (GR) that belongs to the nuclear receptor superfamily. To cover such variety of processes the different glucocorticoids act through different GR isoforms that are originated due to posttranscriptional and post-translational mechanisms. For this reason when evaluating the levels of GRs we should preferentially determine protein levels instead of gene expression. Here, we describe the detection by Western blotting of the GR ( $\alpha$  and  $\beta$  isoforms) protein, using macrodissected brain tissue. 6 7 8 9 10 11 12

**Key words** Glucocorticoid receptor, Brain, Protein levels, Western blot 13

---

## 1 Introduction 14

Glucocorticoids are steroid hormones that are regulated in a circadian and stress-associated manner to maintain various functions of the different body systems [1, 2]. These hormones regulate essential biological processes including growth, inflammation, metabolism, apoptosis, and behavior. Most of these actions are mediated at the genome level through the action of two distinct receptors: the high-affinity mineralocorticoid receptor (MR) and the low-affinity glucocorticoid receptor (GR) [1]; herein, we will describe the detection of GR given its ubiquitous distribution throughout the body (including the central nervous system cells), and because of the availability of good antibodies for its identification. In the absence of glucocorticoids, GR is located in the cytoplasm bound to chaperone proteins such as heat shock protein 90 [3]. Glucocorticoids are lipophilic molecules and they can easily cross the cell membrane. Then, intracellularly they bind to GR that undergoes a conformational change, with the consequent GR translocation to the nucleus. In the nucleus the ligand-bound GR recognizes and binds to specific DNA sequences, known as 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32

33 glucocorticoid-response elements (GREs), which are usually  
34 located in the promoter region of target genes [4]. Once it binds  
35 to these DNA regions GR modulates gene expression mainly  
36 through activation and/or repression mechanisms. The diversity  
37 observed in GR signaling comes from the fact that different genes  
38 may contain, or not, GREs, and multiple receptor isoforms can be  
39 generated by alternative splicing and alternative translation initia-  
40 tion [2]. Also of relevance are the multiple posttranslational modi-  
41 fications that GR can suffer such as phosphorylation, acetylation,  
42 ubiquitination, and SUMOylation with small ubiquitin-related  
43 modifier proteins that can alter the function of this transcription  
44 factor [5]. One of the most relevant posttranscriptional alterations  
45 in GR synthesis is the alternative splicing that occurs in exon 9 (GR  
46 is the product of a single gene, NR3C1, that contains nine exons  
47 both in humans and in rodents). This alternative splicing results in  
48 two isoforms of the GR: GR $\alpha$  and GR $\beta$ , which are identical through  
49 amino acid 727 but differ in their C terminal [5]. The GR $\alpha$  iso-  
50 form binds to glucocorticoids, translocates to the nucleus, and  
51 recruits coregulators to exert transcriptional effects. However, the  
52 GR $\beta$  isoform resides constitutively in the nucleus and acts as a nat-  
53 ural dominant negative inhibitor of the GR $\alpha$  isoform [2]. Here we  
54 will describe a Western blot method to quantify GR in rodents  
55 using an antibody that recognizes both isoforms [GR $\alpha$  (more  
56 abundant) and GR $\beta$ ], since it is an affinity-purified polyclonal anti-  
57 body raised against a peptide mapping at the N-terminus of GR of  
58 mouse origin. The first step in the Western blotting procedure is,  
59 after protein extraction and quantification, using mixtures of solu-  
60 bilized proteins under denaturing conditions [in the presence of  
61 detergent, sodium-dodecyl-sulfate (SDS), and reducing agent  
62 ( $\beta$ -mercaptoethanol)] to separate the macromolecules based on  
63 their molecular weights using a polyacrylamide gel electrophoresis  
64 (SDS-PAGE). After electrophoresis, the separated molecules are  
65 transferred or blotted onto a second matrix, generally a nitrocel-  
66 lulose or polyvinylidene-difluoride (PVDF) membrane. Next, the  
67 membrane is blocked to prevent any nonspecific binding of anti-  
68 bodies to the surface of the membrane. After blocking, the mem-  
69 brane is incubated with a primary antibody that will recognize and  
70 bind the antigen. A labeled secondary antibody that is directed  
71 against the primary antibody is then used. Labels include biotin,  
72 fluorescent probes such as fluorescein or rhodamine, and enzyme  
73 conjugates such as horseradish peroxidase or alkaline phosphatase.  
74 An appropriate substrate is then added to the enzyme and together  
75 they produce a detectable product such as a chromogenic precipi-  
76 tate on the membrane for colorimetric detection. The most sensi-  
77 tive detection methods use a chemiluminescent substrate that,  
78 when combined with the enzyme, produces light as a by-product.  
79 The light output can be captured using film, a CCD camera, or a  
80 phosphoimager that is designed for chemiluminescent detection.

Alternatively, fluorescent-tagged antibodies can be used, which are directly detected with the help of a fluorescence imaging system. Whatever system is used, the intensity of the signal should correlate with the abundance of the antigen on the membrane when normalized for a housekeeping protein.

## 2 Materials

All solutions are prepared using ultrapure water (prepared by purifying deionized water to attain a sensitivity of 18 M $\Omega$  cm at 25 °C) and analytical grade reagents. C57BL/6 male mice at 8- to 9-week of age (Charles River) are used and mice brain regions are collected by macrodissection, using a light microscope (SZX7, Olympus).

### 2.1 Protein Extraction and Denaturation Buffers

- Radio-Immunoprecipitation Assay (RIPA) buffer (*see Note 1*):  
50 mM Tris-HCl, pH 8.0, 250 mM sodium chloride (NaCl), 2 mM EDTA, 10 % Glycerol, 1 Complete Protease Inhibitor Cocktail Tablet (Roche Diagnostics), and inhibitor phosphatases 1 and 2 cocktail (Sigma-Aldrich).  
Prepare:
  - 1 M of Tris-HCl, pH 8.0: weight 30.3 g Trizma Base and add 150 mL of water, adjust pH to 8.0, and adjust with water until 250 mL.
  - 3 M NaCl: weight 5.84 g of NaCl and add 100 mL of water.
  - 0.5 M Ethylene-diamine-tetraacetic acid (EDTA), pH 8.0: weight 186.1 g of EDTA and add 700 mL of water, adjust pH to 8.0 with NaOH 10 M (~50 mL), and complete with water until 1 L.
  - Dissolve one tablet of Complete Protease Inhibitor Cocktail in 2 mL of water.

For 20 mL of RIPA buffer, use 1 mL of 1 M Tris-HCl, pH 8.0, 0.2 mL of 3 M NaCl, 40  $\mu$ L of 0.5 M EDTA, 2.3 mL Glycerol (87 %), 500  $\mu$ L proteases inhibitor, 200  $\mu$ L cocktail inhibitor phosphatases 1 and 2. Add water until 20 mL. Store at -20 °C. Before use, add 1 mM phenyl-methanesulfonyl-fluoride (PMSF). A PMSF solution at 100 mM can be prepared by weighting 0.7 g of PMSF in 40 mL ethanol 100 % (*see Note 2*). Add 200  $\mu$ L of this solution to the 20 mL of RIPA solution.
- For quantification: use Quick Start™ Bradford 1 $\times$  Dye Reagent (Bio-rad, Hercules, CA, USA) following the manufacturer instructions.
- SDS denaturing/lysis buffer (5 $\times$ ): 0.3 M Tris-HCl, pH 6.8, 10 % SDS (*see Note 3*), 25 %  $\beta$ -mercaptoethanol, 0.1 % bromophenol blue, 45 % glycerol. For 10 mL use 6 mL of 0.5 M

122 Tris-HCl, pH 6.8 (*see* Subheading 2.2), 1 g of SDS, 2.5 mL  
123 of  $\beta$ -mercaptoethanol, 0.01 g of bromophenol blue, and  
124 4.5 mL glycerol 100 %. Aliquot and store at  $-20^{\circ}\text{C}$ .

## 125 **2.2 SDS**

### 126 **Polyacrylamide Gel**

### 127 **Materials**

### 128 **and Solutions**

- 129 1. Resolving gel buffer: 1.5 M Tris-HCl, pH 8.8, weigh 181.7 g  
130 Tris, and transfer to a 1 L flask. Add water to a volume of  
131 900 mL. Mix and adjust pH with HCl. Complete to 1 L with  
132 water. Store at  $4^{\circ}\text{C}$ .
- 133 2. Stacking gel buffer: 0.5 M Tris-HCl, pH 6.8. Weigh 60.6 g  
134 Tris and prepare a 1 L solution, as in the previous step. Store  
135 at  $4^{\circ}\text{C}$ .
- 136 3. 40 % acrylamide and bis-acrylamide solution 19:1 (Bio-Rad;  
137 *see* Note 4). Store at  $4^{\circ}\text{C}$ .
- 138 4. SDS 10 %: weigh 5 g of SDS to 50 mL of water. Store at room  
139 temperature (*see* Note 5).
- 140 5. *N,N,N,N'*-tetramethyl-ethylenediamine (TEMED; Bio-Rad).  
141 Store at room temperature (*see* Note 6).
- 142 6. Ammonium persulfate (APS) 10 %: weigh 1 g of APS and dis-  
143 solve in 10 mL of water (*see* Note 7).
- 144 7. SDS-PAGE running buffer: 0.025 M Tris-HCl, pH 8.3,  
145 0.192 M glycine, 0.1 % SDS. Prepare 10 $\times$  native buffer  
146 (0.25 M Tris, 1.92 M glycine, 1 % SDS), for that weigh 30 g  
147 Tris, 144 g glycine, and 10 g SDS, mix, and make it to 1 L  
with water (*see* Note 3). Dilute 100 mL of 10 $\times$  native buffer  
to 900 mL with water.
8. Protein markers (Page Ruler<sup>TM</sup> Plus Prestained Protein Ladder,  
Fermentas, Thermo Scientific).

## 148 **2.3 Immunoblotting**

### 149 **Solutions**

- 150 1. Western blot transfer buffer: 0.025 M Tris, 0.192 M glycine,  
151 and 20 % methanol. Prepare 10 $\times$  transfer buffer (0.25 M Tris,  
152 1.92 M glycine). Weigh 30 g Tris and 144 g glycine to make  
153 it to 1 L with water and mix until dissolution of the powder.  
Dilute 100 mL of 10 $\times$  transfer buffer to 700 mL with water  
and add 200 mL of methanol (*see* Note 8).
- 154 2. Phosphate buffered saline (PBS; 10 $\times$ ): 1.5 M NaCl, 0.1 M  
155 Tris-HCl, pH 7.4. Add 80 g sodium chloride (NaCl), 2 g  
156 potassium chloride (KCl), 14.4 g sodium phosphate dibasic  
157 ( $\text{Na}_2\text{HPO}_4$ ), and 2.4 g potassium phosphate monobasic  
158 ( $\text{KH}_2\text{PO}_4$ ), dissolve with 800 mL of water, adjust pH to 7.4,  
159 and fill with water until 1 L.
- 160 3. PBS-0.2 % T: dissolve 2 mL of Tween-20 in 1 L of PBS. Store  
161 at  $4^{\circ}\text{C}$ .
- 162 4. Blocking solution (5 % milk in PBS-T): weigh 5 g of milk in  
163 100 mL of PBS-T and store at  $4^{\circ}\text{C}$ .

	5. Antibody diluent solution (2.5 % milk in PBS-T): weigh 0.25 g of milk in 10 mL of PBS-T and store at 4 °C.	164 165
	6. Clarity™ Western ECL Substrate (Bio-Rad).	166
<b>2.4 Western Blot Antibodies</b>	1. Primary antibody: mouse anti-GR (M-20; Santa Cruz Biotechnology). This antibody is recommended for detection of mouse and rat GR $\alpha$ and GR $\beta$ . Cross-reactivity with human GR is, however, indicated by the manufacturer.	167 168 169 170
	2. Secondary antibody: anti-rabbit IgG-HRP (SC-2301; Santa Cruz Biotechnology).	171 172
<b>2.5 Immunoblotting Materials</b>	1. Mini-PROTEAN Tetra Cell Electrophoresis Module (Bio-Rad).	173
	2. Mini Trans-Blot cell (Bio-Rad).	174
	3. Nitrocellulose membranes (Amersham, GE Healthcare).	175
	4. Whatman no.3 filter.	176
<hr/>		
<b>3 Methods</b>		177
	Unless otherwise specified, carry out all procedures at room temperature.	178 179
<b>3.1 Preparation of the Gel</b>	1. Set the casting frames (clamp two glass plates in the casting frames) on the casting stands ( <i>see Note 9</i> ).	180 181
	2. Prepare the running gel: 12.5 mL of 40 % acrylamide solution, 12.5 mL 1.5 M Tris-HCl, pH 8.8, 500 $\mu$ L SDS 10 %, complete with water until 50 mL. From this mixture and for each mL of solution use 10 $\mu$ L of APS 10 % and 1 $\mu$ L of TEMED for polymerization. This is a 10 % acrylamide gel that will allow a good resolution of the GR protein, which migrates with an apparent molecular weight of 95 kDa. The volume will depend on the protein apparatus that is available in the laboratory.	182 183 184 185 186 187 188 189
	3. Swirl the solution gently but thoroughly.	190
	4. Pipet appropriate amount of separating gel solution (listed above) into the gap between the glass plates.	191 192
	5. To make the top of the separating gel horizontal, fill the gap until the top of the glasses with water (isopropanol can also be used).	193 194 195
	6. Wait for 20–30 min to let it gelate ( <i>see Note 10</i> ).	196
	7. Prepare the stacking gel: 5 mL of 40 % acrylamide solution, 12.6 mL of 0.5 M Tris-HCl, pH 6.8, 500 $\mu$ L SDS 10 % water until 50 mL. From this mixture and for each milliliter of solution use 10 $\mu$ L of APS 10 % and 1 $\mu$ L of TEMED for polymerization. This is a 4 % acrylamide gel.	197 198 199 200 201

- 202 8. Discard the water or the isopropanol.  
203 9. Pipette in stacking gel until it overflows.  
204 10. Insert the well-forming comb without trapping air under the  
205 teeth. Wait for 20–30 min to let it polymerized.  
206 11. Make sure a complete polymerization of the stacking gel  
207 occurred and remove the comb. Take the glass plates out of  
208 the casting frame and set them in the running tank. Put the  
209 running buffer (electrophoresis buffer) into the inner cham-  
210 ber and keep pouring after overflow until the buffer surface  
211 reaches the required level in the outer chamber.

### 212 **3.2 Preparation** 213 **of the Samples**

214 C57BL/6 male mice (8- to 9-week of age) are utilized. Mice are  
215 anesthetized with ketamine hydrochloride (150 mg/kg) plus  
216 medetomidine (0.3 mg/kg). Upon transcardiac perfusion with  
217 saline buffer, brain regions are collected by macrodissection using  
218 a conventional light microscopy. Samples are immediately frozen  
219 in dry ice and stored at  $-80^{\circ}\text{C}$ . Use chemical inhibitors and con-  
220 trolled temperature is strongly suggested during the below  
described steps to minimize the activity of proteases and other  
enzymes that may modify the protein composition of the sample.

- 221 1. Add RIPA buffer to the brain samples (*see Note 11*) and pass  
222 through a 20 G needle until all the tissue is disrupted  
223 (*see Note 12*).  
224 2. Sonicate the sample on ice five times, for 2 s every time  
225 (*see Note 13*).  
226 3. Centrifuge at 13,000 rpm during 10 min at  $4^{\circ}\text{C}$ .  
227 4. Remove the supernatant for a new centrifugation, as in **step 3**.  
228 5. Perform a protein assay to determine the protein concentra-  
229 tion of the supernatant that should be 5–10  $\mu\text{g}/\mu\text{L}$ .  
230 6. Mix 25  $\mu\text{g}$  of total protein of each sample with SDS denaturat-  
231 ing/lysis buffer (loading buffer). The maximum volume that  
232 is possible to load in the well depends on the comb used.  
233 7. Heat them in boiling water for 5–10 min.  
234 8. Load the protein marker to the first well and then load pre-  
235 pared samples into the other wells, making sure not to over-  
236 flow. Then close the chamber with the cover and connect the  
237 anodes.  
238 9. Set an appropriate volt, usually 100 V, and run the  
239 electrophoresis.  
240 10. As for the total running time, since we are analyzing a protein  
241 that is approximately 95 kDa, stop SDS-PAGE running when  
242 a good resolution of the proteins with higher molecular weight  
243 in the protein marker used is visible.

[AU1]




## 3.3 Immunoblot

We use the wet transfer unit (Mini Trans-Blot cell) to perform the transference of the proteins from the gel to the membrane. Using this electrophoretic transfer system, the gel and membrane sandwich is entirely submerged in transfer buffer within a buffer tank. A nonconducting cassette holds the membrane in close contact with the gel and the cassette assembly is placed in the tank between the electrodes, transverse to the electrical field, and submerged under conducting transfer buffer.

1. Place the transfer tank onto a magnetic stir-plate.
2. Add enough transfer buffer to the tank to fill it approximately halfway, add a stirbar, and begin stirring. This will help to maintain uniform conductivity and temperature during electrophoretic transfer.
3. Set up the cooling system for the tank transfer system.
4. Immediately following SDS-PAGE separate the gel plates with the help of a spatula or similar tool. Remove the stacking gel.
5. Insert the gel carefully in a box with deionized water to remove traces of SDS-PAGE running buffer and then let it equilibrate in Western blot transfer buffer.
6. Let the membrane equilibrate for 5 min in western blot transfer buffer.
7. Embed the six sections of Whatman no. 3 filter paper in the Western blot transfer buffer.
8. Soak a fiber pad in transfer buffer and place it on top of the white side of the cassette.
9. Place on the top of the fiber pad three sections of embedded Whatman no. 3 filter paper and remove the bubbles. On top put the membrane without the formation of bubbles (*see Note 14*).
10. Place the gel on the top of the membrane followed by three or more embedded Whatman no. 3 filter paper. Remove the bubbles (*see Note 14*).
11. Soak a fiber pad in transfer buffer and place it on top of the filter paper.
12. Close the cassette and lock it, insert it into the tank. Make sure the black cassette plate faces the black electrode plate.
13. Add transfer buffer to the tank until the tank is filled and place the lid on top, checking that the color-coded cables on the lid are attached to the electrode cards of the same color.
14. Connect the cables to the power supply and set at 100 V/350 mA and run for 1 h.
15. Upon completion of the run, remove the cassettes and disassemble the gel and membrane sandwich. Rinse the membrane briefly in water to ensure that no residual gel pieces stay adherent to the membrane.

[AU2]



**Fig. 1** The figure shows the immunoblot of protein extracted from mouse hippocampus with anti-GR Ab. One 95 kDa band is detected. It corresponds to GR $\alpha$ , the most abundant isoform

- 288 16. Incubate for 5 min in Ponceau S dye (*see Note 15*).
- 289 17. Wash in water until the Ponceau S staining disappears from the
- 290 membrane.
- 291 18. Block the membranes with blocking solution for 1 h.
- 292 19. Wash 3  $\times$  10 min in PBS-T.
- 293 20. Incubate with diluted (1:1,000 in PBS-T) anti-GR antibody
- 294 and incubate overnight at 4 °C with shaking.
- 295 21. Wash 3  $\times$  10 min in PBS-T.
- 296 22. Incubate with secondary antibody (diluted 1:10,000 in PBS-
- 297 T) for 1 h at RT.
- 298 23. Wash 3  $\times$  10 min in PBS-T.
- 299 24. Develop the blot using Clarity™ Western ECL Substrate and
- 300 incubate for 5 min, following manufacture instructions.
- 301 25. Expose the membrane to X-ray film during approximately
- 302 3–5 min (*see Note 16*).
- 303 26. Check the molecular weight of the band putting the film on
- 304 the top of the membrane and checking the position of the
- 305 band relatively to the protein marker (Fig. 1) (*see Note 17*).

## 4 Notes

306

1. RIPA buffer enables efficient cell lysis and protein solubilization while avoiding protein degradation and interference with the proteins' immune reactivity and biological activity. 307-309
2. Store the PMSF solution at  $-20^{\circ}\text{C}$  protected from the light. 310
3. Care should be taken to weight SDS since it is toxic; individual protection mask should be used. 311-312
4. Due to the Acrylamide/Bis solution toxicity we advice to use the commercially available ones. 313-314
5. SDS can precipitate if stored at  $4^{\circ}\text{C}$  or if the room temperature is too low, if this happens dissolve it by warming if needed. 315-316
6. Due to its pungent smell when pipetting TEMED you should use individual protection mask. If stored at  $4^{\circ}\text{C}$ , its pungent smell is reduced. 317-319
7. When aliquots of APS are stored at  $-20^{\circ}\text{C}$  for long periods, the correct gel polymerization might be compromised. 320-321
8. Care should be taken to add methanol since it is toxic; it should be done in a hood. Use only high-quality, analytical grade methanol. Impure methanol can cause increased transfer buffer conductivity and poor transfer. Avoid adding methanol directly to the  $10\times$  buffer, since it precipitates its ingredients. 322-326
9. All glass plates should be clean and dry. 327
10. Various factors affect the properties of the resulting gel: higher concentration of APS and TEMED will lead to a faster polymerization, but on the other hand, to a lower stability and elasticity. The optimal temperature for gelation is  $23-25^{\circ}\text{C}$ . Low temperature will lead to turbid, porous, and inelastic gels. 328-332
11. A suggestion of the RIPA volume that should be used for different brain regions is indicated in Table 1. 333-334

t1.1 **Table 1**  
t1.2 **Volume of RIPA for each mouse brain region to obtain approximately**  
t1.3 **14 ng of protein**

t1.4	<b>Brain region</b>	<b>RIPA volume (<math>\mu\text{L}</math>)</b>
t1.5	Brainstem	400
t1.6	Cerebellum	300
t1.7	Hypothalamus	150
t1.8	Hippocampus	300
t1.9	Cortex	600
t1.10	Substantianigra	30

- 335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349
12. Mechanical cell lysis usually generates heat. Use cooling where required to avoid overheating the sample.
  13. Pause between sonication steps to avoid overheating.
  14. A 10 mL pipette can be used to remove by rolling out the air bubbles from the gel-membrane sandwich prior to placing in the transfer cassette.
  15. It is recommended to perform Ponceau S staining before proceeding for the immune-blot to control the transference efficiency and the efficient running of the samples in the gel. If the efficiency was low or if the protein running in the acrylamide gel was abnormal or if there are bubbles in the membrane, do not proceed with the immunoblot.
  16. Exposure time to the X-ray film will also depend on the expression level of GR in the sample.
  17. Predicted band size: 87 kDa; observed band size: 95 kDa.

## 350 **References**

- 351  
352  
353  
354  
355  
356  
357  
358  
359  
360
1. Oakley RH, Cidlowski JA (2013) The biology of the glucocorticoid receptor: new signaling mechanisms in health and disease. *J Allergy Clin Immunol* 132:1033–1044
  2. Kadmiel M, Cidlowski JA (2013) Glucocorticoid receptor signaling in health and disease. *Trends Pharmacol Sci* 34:518–530
  3. Grad I, Picard D (2007) The glucocorticoid responses are shaped by molecular chaperones. *Mol Cell Endocrinol* 275:2–12
  4. Meijsing SH, et al (2009) DNA binding site sequence directs glucocorticoid receptor structure and activity. *Science* 324:407–410
  5. Oakley RH, Cidlowski JA (2011) Cellular processing of the glucocorticoid receptor gene and protein: new mechanisms for generating tissue-specific actions of glucocorticoids. *J Biol Chem* 286:3177–3184
- 361  
362  
363  
364  
365  
366  
367  
368

# Author Queries

Chapter No.: 20      0002151701

Queries	Details Required	Author's Response
AU1	Please provide g-force value for 13,000 rpm.	
AU2	Please check if edit to the sentence "Let the membrane equilibrate ..." is appropriate.	

Uncorrected Proof