


Cashew Nut Allergy: Clinical Relevance and Allergen Characterisation

Cíntia Mendes¹ · Joana Costa¹  · António A. Vicente² · Maria Beatriz P. P. Oliveira¹ · Isabel Mafra¹

Published online: 1 September 2016
© Springer Science+Business Media New York 2016

Abstract Cashew plant (*Anacardium occidentale* L.) is the most relevant species of the *Anacardium* genus. It presents high economic value since it is widely used in human nutrition and in several industrial applications. Cashew nut is a well-appreciated food (belongs to the tree nut group), being widely consumed as snacks and in processed foods by the majority of world's population. However, cashew nut is also classified as a potent allergenic food known to be responsible for triggering severe and systemic immune reactions (e.g. anaphylaxis) in sensitised/allergic individuals that often demand epinephrine treatment and hospitalisation. So far, three groups of allergenic proteins have been identified and characterised in cashew nut: Ana o 1 and Ana o 2 (cupin superfamily) and Ana o 3 (prolamin superfamily), which are all classified as major allergens. The prevalence of cashew nut allergy seems to be rising in industrialised countries with the increasing consumption of this nut. There is still no cure for cashew nut allergy, as well as for other food allergies; thus, the allergic patients are advised to eliminate it from their diets. Accordingly, when carefully choosing processed foods that are commercially available, the allergic consumers have to rely on proper food labelling. In this sense, the control of labelling compliance is much needed, which has prompted the development of proficient analytical

methods for allergen analysis. In the recent years, significant research advances in cashew nut allergy have been accomplished, which are highlighted and discussed in this review.

Keywords *Anacardium occidentale* L. · Food allergy · Allergens · Prevalence · Threshold levels · Immunotherapy · Clinical relevance · Cashew detection

Introduction

Cashew plant (*Anacardium occidentale* L.) belongs to the Anacardiaceae family, which comprises nine species of the *Anacardium* genus [1]. *A. occidentale* is the most relevant in terms of economic interest, although other trees and shrubs of this genus have begun to gain special attention. Additional data on each species is presented in Table 1, namely their geographical distribution, applications and potential allergenic proteins.

A. occidentale is a tropical evergreen tree that is well known for its fruits and seeds. The cashew fruit (cashew apple) is actually a swollen peduncle (pseudofruit) with sweet flavour and aroma. The pulpy and juicy part of cashew apple is highly popular, being consumed as juices, jams and alcoholic beverages [2, 3]. The cashew nut (seed) grows externally, inside a hard shell, at the end of the peduncle. The hard shell is composed by two layers that form a honeycomb structure containing an oily substance (phenolic cashew nut shell liquid), which presents several industrial applications, namely in plastic, paint, resin and surface coating industries [2]. Inside the hard shell and protected from the oily substance lays the kernel (cashew nut) that is a rich source of protein, carbohydrate and triglyceride oil. Cashew nuts are normally consumed roasted and salted, being worldwide appreciated as snacks.

Cíntia Mendes and Joana Costa contributed equally to this work.

✉ Joana Costa
jbcosta@ff.up.pt

✉ Isabel Mafra
isabel.mafra@ff.up.pt

¹ REQUIMTE-LAQV, Faculdade de Farmácia, Universidade do Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal

² CEB, Centro de Engenharia Biológica, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal

Table 1 Geographic distribution, uses and industrial applications, identified allergens and cross-reactivity of the nine species of *Anacardium* genus [1–5]

<i>Anacardium</i> species	Geographic distribution	Uses	Allergenic proteins	Cross-reactivity
<i>A. occidentale</i> (also known as <i>A. microcarpum</i>)	Endemic from South America but currently grown worldwide (major producers: India, Indonesia, Brazil, China, Mozambique, Tanzania and Vietnam)	Edible fruits and seeds (e.g. snacks, jams, juices), industrial applications (e.g. paints, plastics), traditional medicine (e.g. treatment of malaria, diabetes)	Ana o 1, Ana o 2 and Ana o 3	–
<i>A. excelsum</i> (also known as <i>A. rhinocarpus</i>)	Central and South America (e.g. Venezuela, Colombia, Ecuador, Brazil)	Wood for construction, fruits and seeds are toxic for humans but can be consumed after processing (e.g. roasting)	–	–
<i>A. giganteum</i> (wild cashew)	South America (e.g. Brazil (Amazonia forest), Colombia; Peru, Suriname; Venezuela)	Closely related to <i>A. occidentale</i> . Wood for construction (local important timber—South America). Edible fruits and seeds consumed by local population (e.g. juices, jams)	–	Potential cross-reactivity with <i>A. occidentale</i> since species are genetically related
<i>A. othonianum</i>	South America (Brazil)	Edible fruits and seeds consumed by local population (e.g. juices, jams), traditional medicine (e.g. treatment of infections, diarrhoea)	–	–
<i>A. humile</i> (also known as <i>A. pumilum</i>)	South America (e.g. Brazil-Planalto Central, Bolivia, Paraguay)	Closely related to <i>A. occidentale</i> . Edible fruits and seeds consumed by local population (e.g. juices, jams), traditional medicine (e.g. antibacterial, antihelminthic and antifungal activities)	–	Potential cross-reactivity with <i>A. occidentale</i> since species are genetically related
<i>A. spruceanum</i>	South America (e.g. Brazil-Amazonas, Guyana; Suriname; Venezuela, Peru)	Ornamental purposes (white leaves)	–	–
<i>A. nanum</i>	South America (Brazil-Amazonas)	Edible fruits (e.g. jams, juices)	–	–
<i>A. negense</i>	South America (Brazil-Amazonas)	This species do not produce fruits. The seeds (nuts) are eaten by the locals, normally after roasting	–	–
<i>A. corymbosum</i>	South America (Brazil-Planalto Central)	The seeds can be consumed after roasting	–	–

A. occidentale species originated from South America (e.g. Brazil, Venezuela), but it is currently cultivated in several countries of Africa and Asia (e.g. Vietnam, India, United Republic of Tanzania, Indonesia) [6]. In 2013, Vietnam was the main producer of cashew nut with shell, thus assuring 25 % of its global production. In the same year and in terms of trade, cashew nut ranked second place just behind almond [6]. Included in the tree nuts, cashew nut is also related to several beneficial health effects that are commonly attributed to this group of nuts. Accordingly, cashew nut and its co-products have a long history of worldwide application in traditional medicine for the treatment of different health conditions, such as asthma, diabetes, skin infections and inflammation, amongst others (Table 1) [3]. However, no clinical studies have yet proven its actual medicinal efficacy [7].

Owing to the health food claims attributed by the World Health Organization and the Food and Agriculture Organization (WHO/FAO) to tree nuts [8], the consumption of cashew nut and other tree nuts has been increasing in the past years. However, cashew (nut and fruit) can also represent a concrete health risk for a small but rather significant part of the world's population, namely the food-sensitised/allergic individuals.

With the increasing awareness that food allergies represent an important public health problem, legal measures have been established to protect the life of those individuals. In 1985, the Codex Alimentarius Commission issued a recommendation for mandatory labelling of pre-packaged foods susceptible of containing potentially allergenic ingredients. Accordingly, a list of priority allergenic foods (tree nuts, peanuts, gluten-containing cereals, soybean, fish, eggs, milk and crustaceans) and sulphites was proposed by the Codex Alimentarius Commission [9]. Within the European Union (EU), the first priority list was issued by the Directive 2003/89/EC [10], adding sesame, celery and mustard to the previous list [9], totalising 12 groups of allergenic foods. Since then, the EU has established legislation increasing the priority list of allergens to 14 groups (by adding molluscs and lupine) that are required to be emphasised from the rest of the ingredients enumerated in processed foods, regardless of their quantity [11, 12]. Using the Codex Alimentarius Commission and/or the EU standards as guidelines, many countries have adopted allergen labelling regulatory frameworks [13, 14]. Owing to the fact that tree nuts are one of the eight groups responsible for a great number of allergic reactions, they are included in the list of allergenic foods with mandatory labelling in most countries/regions (e.g. USA, Canada, EU), with exception of Japan and Korea [13, 14].

In the recent years, significant research advances in cashew nut allergy, namely prevalence, diagnosis, immunotherapy, clinical relevance and thresholds, as well as biochemical characterisation of cashew nut allergens, processing and digestibility studies, management and method development

for its detection, have been achieved. The research progresses in cashew nut allergy have been summarised and discussed in this review.

Epidemiology of Cashew Allergy

The definition of the prevalence of food allergies has been estimated on the basis of different indicators. Amongst them, surveys and questionnaires have been widely used to predict the prevalence of several food allergies. However, the retrieved information must be carefully evaluated since the sensitised/allergic individuals might overrate self-allergic symptoms. To overcome these drawbacks, more objective indicators, such as provocation food tests (open food challenges (OFC) and double-blind placebo-controlled food challenges (DBPCFC)) or in vitro (serum IgE) and in vivo (skin prick tests (SPT)) diagnostic tools, to assess food sensitisation are recommended [15]. However, OFC and DBPCFC studies are more difficult to perform since their assessment is highly dependent on the collaboration of the allergic individuals.

So far, there are several studies reporting prevalence data about tree nut allergies, but only a few specify information regarding individual nuts (e.g. cashew nut). Additionally, most of the reports involve small geographical regions (e.g. city, country), which retrains a broad overview about the real prevalence of cashew nut or other food allergies. Literature reporting the prevalence of cashew nut allergy is available for some countries in Europe, the USA and Australia, but no information could be retrieved from emergent economies (Asia or Africa).

Like other nuts, the prevalence of cashew nut allergy varies from region to region, presenting its highest reported incidence in the USA [16]. A voluntary registration involving 5149 participants with peanut and tree nut allergies from 48 states was established in 1997 to evaluate their prevalence in the USA. On the basis of a structured questionnaire that encompassed peanut and different nuts, it was possible to estimate that walnut (34 %) and cashew nut (20 %) allergies presented the highest incidence rate amongst allergic patients [17]. Using more objective indicators, such as SPT and DBPCFC, Fleischer et al. [18] reported high prevalence of walnut and cashew nut allergies, both with 30 % in a USA test population of 101 patients with previous clinical diagnosis of tree nut allergies.

Besides USA, cashew nut allergy also seems to be epidemiologically increasing in other regions over the past years, namely in Northern Europe [19, 20]. A study using clinical records from a hospital in Uppsala County (Sweden) over a 10-year period (2001–2010) evidenced that peanuts and tree nuts accounted for 50 % of the 703 food allergic cases in children (<18 years), being more frequently associated with epinephrine treatment and hospitalisation. Over the same

period, the estimated prevalence of cashew nut allergy was 6 % of food allergic individuals, with reports of increasing frequency and severity of episodes caused by cashew nut ingestion [19]. In a different region of Sweden (Stockholm County), similar data could be retrieved from three paediatric hospitals in 2007. Accordingly, 15 % of the reported food allergies in children were attributed to tree nuts. From those, 5 % were triggered by cashew nut ingestion with most patients experiencing anaphylaxis [20]. In a 3-year follow-up study involving 139 tree nut allergic adults from Utrecht (Netherlands), cashew nut allergy was estimated with a prevalence of 20 % [21]. In France, Moneret-Vautrin [22] estimated an incidence of 10.3 % of cashew nut allergy on a test population of 141 patients, reporting that the number of anaphylaxis cases increased from 3 to 9 % between 2003 and 2007, respectively.

In a 5-year study (1999–2003) based on clinical records from a paediatric hospital in Australia (Melbourne), peanut (18 %) and cashew nut (13 %) were the most common cause for anaphylaxis in children [23]. Similar prevalence of cashew nut allergy (12.6 %) was described by Davoren and Peake [24] using clinical reports of a test population comprising 213 peanut and tree nut allergic children (<18 years) from Brisbane (Australia) during a period of 42 months. Moreover, in patients with diagnosed cashew nut allergy, more than 74 % had experienced severe and life-threatening episodes, such as anaphylaxis [24].

Diagnosis, Therapy and Immunotherapy

Cashew nut allergy is a classical IgE-mediated disorder. The ingestion of cashew nut seems to be the principal sensitisation path for its allergy, although mechanisms associated with poor skin barrier function (atopic dermatitis) and loss of function in the filaggrin gene have been also highlighted as increasing risk factors for the development of cashew nut and other food allergies [25, 26]. Like for most of food allergies, the initial diagnosis of cashew nut allergy is usually built on the basis of a clear definition of a medical history [27, 28]. Accordingly, several aspects, such as the amount of ingested cashew nut, the type of clinical presentations and organs/systems (skin, gastrointestinal or respiratory) affected, the temporal association between ingestion and the appearance of the first observable symptoms and the symptoms after eating similar foods, must be considered [27].

Following a record of medical history, diagnostic testing methods, namely the use of *in vivo* SPT and/or *in vitro* fluorescent-enzyme immunoassays (FEIA-CAP) or radioallergosorbent tests (RAST), are recommended. The SPT are performed by placing a small amount of the allergen into the skin, eliciting a localised allergic reaction in the form of a wheal at the testing site, whilst the FEIA-CAP and RAST

are based on the detection of specific IgE in serum. However, these tests can only be considered as mere indicators of sensitisation rather than truly predictive tools for food allergy. In spite of presenting high sensitivities, negative results of the diagnostic *in vivo* and/or *in vitro* tests cannot be considered for allergy exclusion [29]. More recently, allergen microarrays, such as ImmunoCAP ISAC (diagnostic testing, Thermo Fisher Scientific, Waltham, MA, USA) or the MeDALL allergen-chip based platforms, have been advanced as excellent tools for the simultaneous detection of multiple allergen-specific IgE towards a large number of clinically relevant allergens [30].

Regarding cashew nut, reports suggest that a wheal diameter over 3 mm in SPT [25, 31, 32] and/or a value ≥ 0.35 kUA/L in immunoCAP FEIA test can be indicative of allergy [21, 25, 33]. Mean values of wheal diameter of 7 mm (3–16 mm) and 8.5 mm (6.5–10 mm) were reported by Rancé et al. [25] and Clark et al. [31] for children (<18) with medical history of cashew nut allergy. To increase the accuracy of the SPT, the area of the allergen-induced wheal can be compared with the area of a histamine-induced wheal size (as positive control) to correct for skin histamine sensitivity. The ratio is defined as histamine equivalent prick (HEP) index [34]. Considering an average wheal diameter over 3 mm and HEP index area ≥ 0.4 as baseline for positive sensitization, van der Valk et al. [32] reported a mean value of 3.02 of HEP index (range 0–15.16) in 164 of cashew nut allergic children.

Using the immunoCAP FEIA test, the mean values for cashew nut-specific IgE described by Rancé et al. [25], van der Valk et al. [32], and Maloney et al. [33] were 3.1, 3.72, and 5.5 kUA/L, respectively. Recombinant Ana o 3 has been reported as a highly accurate diagnostic marker for cashew nut allergy, showing higher sensitivity and specificity levels (93.7 and 94.4 %, respectively) over the conventional whole cashew nut extract (95.2 and 58.3 %, respectively) [35]. Additionally, the recombinant Ana o 3 was also considered an excellent diagnostic marker for pistachio allergy with similar sensitivity and specificity levels of cashew nut diagnosis. For rAna o 3, an optimal cutoff point of 0.16 kUA/L was reported as indicative for cashew nut and pistachio allergies [35].

Besides SPT and serum-specific IgE tests, oral provocation tests such as OFC or DBPCFC are considered the “gold standard” tools to identify allergenic foods (e.g. cashew nut) and/or to confirm the allergy [28]. In general, individuals presenting a convincing medical history of cashew nut allergy, SPT of wheal diameter over 3 mm (HEP index ≥ 0.4) and/or serum-specific IgE of 0.35 kUA/L are often purposed for oral provocation tests (DBPCFC or OFC) with increasing doses of cashew nut [32, 34]. The oral provocation tests are normally discontinued and considered positives when objective symptoms occur, when severe subjective symptoms persist (>1 h) or when subjective symptoms re-appear after second dose of the same challenge material [32]. Patients with negative test

challenges are frequently advised to introduce cashew nut at home following a defined schedule with increasing doses of the referred nut [32]. Therefore, the correct clinical diagnosis of cashew nut allergy as well as other food allergies should be made considering data from different indicators, namely the medical history, the results from diagnostic testing tools (SPT and serum-specific IgE tests) and food challenges (Fig. 1).

Currently, most food allergic individuals are strongly recommended to follow a prophylactic measure that consists on the total avoidance of the allergenic food. However, accidental exposure to the allergenic food (e.g. cashew nut) or to cross-reactive ones is frequently occurring, so therapeutic measures must be applied in those circumstances. Consequently, antihistamines are amongst the most frequently prescribed drugs, followed by epinephrine auto-injectors and systemic corticoids [21, 31]. Amongst cashew nut allergic reactions, the treatment with intramuscular epinephrine is often common owing to the severity of the allergic reactions (anaphylaxis) [31]. Treatment with intramuscular epinephrine (into the mid-outer thigh) is normally the first line of intervention for the emergency management of an anaphylactic shock [36]. Accordingly, allergic patients at risk of developing anaphylaxis towards cashew nut or other foods are recommended to carry with them an epinephrine auto-injector to be use in case of emergency. This type of auto-injectors (e.g. EpiPen) is normally prepared to release a dose of intramuscular epinephrine, but depending on different factors (e.g. patients' sex, body habitus), the injection might end up being subcutaneous [37]. Despite the fact that in most of the cases the subcutaneous injection of epinephrine is known to work, the administration of subcutaneous or inhaled adrenaline in the treatment of anaphylaxis is not recommended according to recent guidelines [36].

The administration of inhaled short-acting beta-2 agonists (to relieve symptoms of bronchoconstriction) along with oxygen and fluids can be used as a second line of intervention. Glucocorticosteroids and H1 and H2 antihistamines represent the third line of treatment for anaphylaxis. Intravenous or oral glucocorticosteroids are thought to prevent protracted anaphylaxis symptoms, especially in patients with concomitant asthma. The administration of H1 and H2 antihistamines is only recommended for the relief of cutaneous symptoms of anaphylaxis (angioedema, urticaria) [36].

More recently, some strategies have been applied aiming at inducing tolerance to different allergens. Immunotherapy has been faced as a potential tool to induce immunomodulation and a desensitisation state in allergic patients, through daily exposure to increasing doses of the offending food. So far, some immunotherapies based on different routes of administration (subcutaneous, epicutaneous, sublingual and oral) have been described for the most common allergenic foods, namely egg, peanut or milk [38–41].

Regarding cashew nut immunotherapy, only few studies could be found in literature, being mostly carried out in animal models (mice). Kulis et al. [42] reported that cashew nut immunotherapy was effective in protecting mice from allergic reactions to cashew nut and had a favourable impact on the walnut challenge in cashew nut/walnut-sensitised mice, leading to a significant decrease of the allergic symptoms. The authors also observed that immunotherapy led to increased cashew nut-specific IgG1 with decreased Th2-type cytokine production, thus suggesting a potential protective role in vivo [42]. In a subsequent study, Kulis et al. [43] proposed the use of pepsinised cashew nut proteins, which elicit weaker allergic reactions, as highly effective immunotherapy reagents for treating cashew nut allergy in mice. Like for immunotherapy




Fig. 1 Diagnosis of food (cashew nut) allergy

with native cashew nut proteins, in immunotherapy with pepsinised proteins, the levels of IgG1 and IgG2a were increased and the production of Th2-type cytokines decreased [43], thus proposing the use of hypoallergenic cashew nut proteins in allergy treatment.

Information on clinical studies involving the administration of cashew nut to human-allergic patients as immunotherapy is very limited. One of the reports described the application of oral immunotherapy to a 22-year-old man with cashew nut, walnut and peanut allergies, using a daily dose of 0.5 g of total nut (0.17 g of each offending nut) during 8 months. Levels of nut-specific IgE increased in the first month but decreased after 3 and 6 months. The clinical symptoms also decreased in intensity and frequency along the 8-month treatment [44]. The results evidenced the successful use of immunotherapy to induce some tolerance to cashew nut, walnut and peanut allergens upon potential accidental exposure to those nuts.

In a different study that evaluated the ability of omalizumab (anti-IgE) to induce desensitisation to multiple allergens, cashew nut was used as one of the test foods [45, 46]. Results from phase 1 trials seem to indicate that rush oral immunotherapy to multiple foods with 16 weeks of treatment with omalizumab could allow a fast desensitisation in subjects with multiple food allergies. However, the study is still at a preliminary stage and further understanding about the safety and efficacy parameters of multi-oral immunotherapy experimental treatments with and without omalizumab is much needed [45].

Clinical Threshold Levels for Cashew Nut Allergy

As part of food allergy diagnosis, individuals with medical history and/or positive results *in vitro/in vivo* diagnostic tests are often challenged to enter OFC or DBPCFC. From these trials, clinical threshold levels can be characterised for allergenic foods at individual basis [47]. So far, several studies conducted on a variety of allergenic foods evidenced that individual thresholds or lowest observed adverse effect level (LOAEL) vary within food allergic population. Considering that population distribution of minimum eliciting doses (ED) is estimated to be affected by host- (e.g. genetic predisposition, environmental factors), food- (e.g. food processing habits) and population-associated factors (e.g. demographics, geography), the definition of global threshold levels for allergenic foods is difficult to achieve [48]. Nonetheless, the information retrieved from individual threshold levels is foreseen as valuable data for the definition of population threshold levels, thus providing guidance for regulatory and public health agencies, as well as for allergic patients and food industry [47, 49].

Based on OFC and DBPCFC studies, the threshold levels of some allergenic foods (peanut, egg or milk) have been advanced in recent years. However, the determination of

cashew nut threshold doses is still at a preliminary stage. In a test population of 31 cashew nut allergic children (<18 years), the minimum dose eliciting objective allergic symptoms in 5 % (ED₀₅) of the patients corresponded to 7.41 mg of cashew nut protein. When considering any type of symptoms, the ED₀₅ was 23 times lower, being 0.32 mg of cashew nut protein [50].

In a different study using DBPCFC on a test population of 179 children (2–17 years), 1 mg of cashew nut protein was advanced as the minimal eliciting dose for triggering an allergic reaction [32]. Following a revision of allergen reference doses, statistical dose distribution models (log-normal, log-logistic, Weibull) were applied to individual threshold doses for each allergenic food aiming at defining reference doses at population level [49, 51]. According to the present data of ED₀₅ for cashew nut, a reference dose of 2.0 mg of protein has been provisionally proposed for this allergenic food [49, 51]. However, until further information regarding individual threshold levels in cashew nut allergic adults is available, Crevel et al. [48] recommend considering the reference dose of 0.1 mg of protein as reference dose for all tree nuts.

Allergen Characterisation and Clinical Relevance

From the synopsis of allergology and protein evolution, food allergens have been classified as belonging to a few families of proteins, according to their structures and biochemical functions [52]. So far, almost 65 % of the identified plant food allergens are included in four families of proteins (cupin superfamily, prolamin superfamily, Bet v 1-related proteins and profilins) [53].

From an immunological point of view, food allergens can be further categorised as class I or class II, according to their subjacent pattern of sensitisation. Class I food allergens are normally considered as presenting high stability to thermal denaturation, acidic environment and gastrointestinal proteases. Owing to these features, class I food allergens are more likely to function as sensitizer agents through the gastrointestinal tract, being usually responsible for eliciting moderate to severe allergic reactions [54]. Seed storage proteins of the cupin and prolamin superfamilies, namely vicilins, legumins and 2S albumins, are examples of class I food allergens [55, 56]. Class II food allergens present less thermal and digestive stability, thus suggesting that they cannot function as primary sensitizers. As consequence, the route of sensitisation occurs via inhaled allergens that lead to the production of IgE. Class II food allergens are most commonly related to mild clinical manifestations, although moderate or severe symptoms might also occur as consequence of adverse immunological responses to these allergens [54]. In the specific case of cashew nut, the identified allergens are seed storage

proteins (vicilins, legumins and 2S albumins), which are classified as class I food allergens.

In terms of clinical presentation, the reported symptoms of cashew nut allergy are commonly classified as severe and potentially life-threatening, occurring with complex manifestations that often involve more than one target organ/system. Accordingly, most of the allergic reactions to cashew nut evidence skin lesions followed by respiratory and/or gastrointestinal symptoms [57]. Mild clinical symptoms such as oral allergy syndrome are not associated with cashew nut allergy. Within the sphere of clinical manifestations, urticaria and angioedema (cutaneous), abdominal pain and vomiting (gastrointestinal), wheeze, rhino-conjunctivitis, laryngeal oedema and anaphylaxis are amongst the most classical symptoms related to cashew nut allergy [19, 21, 24, 25, 31, 58]. Anaphylaxis is one of the most severe and complex clinical presentation of an allergic reaction since it normally involves several systems (cutaneous, respiratory and cardiovascular). Anaphylaxis is a common clinical presentation of cashew nut allergy similarly to peanut and other nut allergies [24, 57, 58].

So far, three allergenic groups of proteins have been identified and characterised in cashew nut, which have been already included in the World Health Organization/International Union of Immunological Societies (WHO/IUIS) official list of allergens [59]. From those, two allergens belong to the cupin (Ana o 1 and Ana o 2) and one to the prolamin (Ana o 3) superfamilies of proteins.

Cupin Superfamily

The cupin superfamily presents a wide diversity of proteins, which can be found in bacteria and eukaryotes, including plants and animals [60]. Members of this superfamily are characterised by two short conserved consensus sequence motifs and a beta-barrel structural core domain, which is the source of its designation “cupin” [61]. Vicilins and legumins are 7S- and 11S-type globulins, respectively, with two core domains (bicupins), being considered the major seed storage proteins in several nuts and legumes [61]. Like in other nuts, allergenic vicilins and legumins were also identified in cashew nut.

Ana o 1—Vicilin

Vicilins present a trimeric structure with a total molecular weight of 150–190 kDa, being characterised by three subunits of 40 to 80 kDa each [62]. These proteins are composed of N- and C-terminal domains that are structurally stabilised by non-covalent hydrophobic interactions, hydrogen bonds and van der Waals interactions. Vicilins are glycosylated, presenting one or two N-linked glycosylation sites at the C-terminal domain, which seem to have a significant stabilising effect on protein structure [61].

Ana o 1 are the allergenic vicilins identified and characterised in cashew nut [59, 63], encompassing two isoforms (proteins with more than 90 % of sequence identity) [64]: Ana o 1.0101 and Ana o 1.0102 with 538 and 536 amino acids (aa), encoded by mRNA sequences of 1810 and 1660 bp, respectively (Table 2). Using competent *Escherichia coli* BL21 (DE3) cells, Wang et al. [65] were able to express recombinant Ana o 1.0101 and Ana o 1.0102 with molecular sizes of 65 and 55 kDa, respectively, after digestion with thrombin to separate the 43-kDa maltose-binding protein from the expressed fusion proteins. Sequence identity and similarity with other vicilins (oil palm, macadamia nut and walnut) were in the ranges of 33–40 and 54–59 %, respectively. The IgE-binding capacity of these recombinant proteins was further evaluated by immunoblotting with sera from cashew nut allergic patients. As a result, 10 out of the 20 sera from cashew nut allergic patients (50 %) were IgE-reactive to the recombinant proteins, thus suggesting that Ana o 1 is a major allergen in cashew nut [65]. In the same study, the authors also sequenced possible epitopes for Ana o 1 and evaluated their IgE reactivity with three sets of pooled sera from different cashew nut allergic patients. From the 11 peptides covering the entire length of the protein, three were strongly immunoreactive, indicating that they are most likely immunodominant epitopes [65]. Using the same procedure to obtain a recombinant Ana o 1, Guan et al. [66] demonstrated that two of the linear epitope-contributing peptide segments, which were previously reported by Wang et al. [65], participate in a conformational epitope. Additionally, the authors also revealed three new peptide segments that are believed to function as independent linear epitopes when complexed with some antibodies and/or together, contributing to one or more conformational epitopes [66].

More recently, Reitsma et al. [67] were able to extract and purify native Ana o 1, Ana o 2 and Ana o 3 allergens from defatted cashew nut flour. Using non-reducing denaturing conditions by SDS-PAGE and further confirmed by matrix-assisted laser desorption/ionisation-tandem mass spectrometry (MALDI-MS/MS), Ana o 1 was identified as a single protein of 50 kDa or as a dimer of 100 kDa. The predicted glycosylation of Ana o 1 was also confirmed by glycoprotein staining. Besides some cross-reactivity between Ana o 1 and Ana o 2 (both globulins), minor unexpected cross-reactivity was observed between Ana o 1 and Ana o 3 allergens [67].

Using molecular modelling, Barre et al. [68] proposed three-dimensional (3D) models for allergenic vicilins of peanut (Ara h 1), walnut (Jug r 2), hazelnut (Cor a 11) and cashew nut (Ana o 1). 3D models built for the Jug r 2, Cor a 11 and Ana o 1 monomers exhibit an overall cupin fold very similar to that of beta-conglycinin (soybean), by presenting a cupin motif made of two tandemly arrayed modules. When comparing linear epitopes, Ana o 1 exhibited homology with Ara h 1 and beta-conglycinin, but the highest epitope homology was observed with vicilins from other nuts, namely Cor a 11

Table 2 Identification of cashew nut allergens according to their biochemical classification, biological function, clinical relevance and respective accession numbers

Allergen	Isoallergens	Isoforms	MW (kDa)	Length (aa)	Protein superfamily	Biochemical classification	Clinical relevance	Nucleotide (NCBI)	Protein (NCBI)	Protein (UniProt)
Ana o 1	Ana o 1.01	Ana o 1.0101 Ana o 1.0102	61.8 61.6	538 536	Cupin	7S vicilin	Major allergen. Induces severe and potentially life-threatening symptoms, namely anaphylaxis	AF395894 AF395893	AAM73730 AAM73729	Q8L5L5 Q8L5L6
Ana o 2	Ana o 2.01	Ana o 2.0101	52.0	457	Cupin	11S legumin	Major allergen. Induces severe and potentially life-threatening symptoms, namely anaphylaxis	AF453947	AAN76862	Q8GZP6
Ana o 3	Ana o 3.01	Ana o 3.0101	16.3	138	Prolamin	2S albumin	Major allergen. Induces severe and potentially life-threatening symptoms, namely anaphylaxis	AY081853	AAL91665	Q8H2B8

(hazelnut) and Jug r 2 (walnut) [68]. In in vivo cross-reactivity assays, cashew nut-sensitised mice reacted to 1 mg of walnut almost as strongly as to the same quantity of cashew nut, often experiencing severe anaphylactic reactions.

Clinical symptoms were weaker in cashew nut-sensitised mice upon challenging with 1 mg of peanut than with equal amounts of cashew nut or walnut, although symptoms increased in intensity depending on the dose of peanut used for the challenge. As a result, Jug r 2 and Ara h 1 were the cross-reactive allergens implicated in the allergic responses triggered in cashew nut-sensitised mice [69]. Contrarily to a previous report stating that there was no cross-reactivity between cashew nut and peanut allergens [70], Ana o 1 was also found to be IgE-reactive with human sera of peanut-allergic patients, thus suggesting cross-reactivity between Ara h 1 and Ana o 1 allergens [68]. Strong cross-reactivity of cashew nut with pistachio (both members of the Anacardiaceae family) has been previously described in cashew nut and/or pistachio-allergic patients [71–73]. Confirming these clinical evidences, a high overall identity (80 %) and similarity (90 %) were found between the primary sequences of Ana o 1 and Pis v 3 (pistachio), with Pis v 3 sharing high homology in two of the Ana o 1 immunodominant peptides [74, 75]. In some cashew nut/pistachio allergic patients, cashew nut seems to act as the primary sensitizer agent in both cashew nut and pistachio allergies [75].

Ana o 2—Legumin

The legumins or 11S globulins are a different class of functional proteins that belong to the cupin superfamily. Legumins are multimeric proteins that exist as a mixture of trimers and hexamers of 50–60 kDa linked by non-covalent interactions [76]. Each monomer results from the expression of multiple genes and is synthesised as a single polypeptide, which is then post-translationally cleaved into two polypeptides—one acidic (30–40 kDa) and one basic (~20 kDa)—linked by a disulphide bond. Contrarily to vicilins, legumins are normally non-glycosylated proteins. Legumins are seed storage proteins, representing more than 50 % of protein fraction in several nuts and legumes [76].

Ana o 2 is the allergenic legumin identified and characterised in cashew nut [59, 63], comprising one isoform: Ana o 2.0101 with 457 aa and a molecular size of approximately 55 kDa, which is encoded by a mRNA sequence of 1671 bp (Table 2). Upon characterisation of the soluble protein fraction of cashew nut, Teuber et al. [77] identified the heavy chain of a legumin that present high IgE reactivity (>50 %) with sera from cashew nut allergic patients. Accordingly, the protein was designated as Ana o 2 and classified as a major allergen in cashew nut [77]. Native Ana o 2 was isolated presenting bands of 33, 53 and 120 kDa, which were estimated to correspond to a heavy chain polypeptide, a

monomer and a dimer of legumin, respectively [78]. Using non-reducing denaturing conditions by SDS-PAGE and MALDI-MS/MS, 53-, 30- and 21-kDa bands of Ana o 2 were confirmed as the complete protein (monomer), large subunit and small subunit, respectively [67].

Like for the Ana o 1 allergen [65], the recombinant Ana o 2 was also expressed by competent *E. coli* BL21 (DE3) cells as a 93-kDa fusion protein with a 43-kDa maltose-binding protein [78]. Sequence identity and similarity were in ranges of 42–58 and 58–74 %, respectively, with different allergenic 11S globulins, namely Cor a 9 (hazelnut), Jug r 4 (walnut), Ara h 3 (peanut), Ses i 7 (sesame) and Gly m 6 (soybean) amongst others [78, 79]. The IgE-binding capacity of recombinant Ana o 2 was further evaluated by immunoblotting with sera from 21 cashew nut allergic patients with clear history of life-threatening adverse responses to cashew nut. The sera from 13 patients were IgE-reactive to recombinant Ana o 2, which represents 62 % of the test population [78], thus confirming its previous classification as a major allergen of cashew nut [77]. In the same study, Ana o 2 was sequenced to determine possible IgE epitopes against pooled sera from different cashew nut allergic patients. From a total of seven epitopes that were strongly IgE-reactive (potential immunodominant epitopes), six epitopes were located in the acidic polypeptide [78], suggesting that this region might be predominantly related to allergenicity.

Linear IgE-binding epitopes were mapped from 3D models of Ana o 2 (cashew nut), Jug r 4 (walnut), Cor a 9 (hazelnut) and Ara h 3 (peanut) obtained by homology modelling [80]. Based on the 3D model of Ana o 2, three linear epitopes were identified on its surface, most of those containing positively or negatively charged residues, which were predominantly located on the acidic face of Ana o 2 trimer. Epitope no. 13 of Ana o 2 overlaps with one epitope (epitope no. 3) of Ara h 3 (peanut). Both epitopes form a strongly charged area on the molecular surface of the corresponding allergens and exhibit a rather conserved overall conformation, despite the moderate degree of amino acid sequence identity [80]. These common structural features seem to indicate that linear IgE-binding epitope no. 13 of Ana o 2 corresponds to a consensual epitope that also occurs in other nut (Ara h 3, Cor a 9 and Jug r 4) and plant (G2-glycinin) legumins [79, 80]. Additionally, Ana o 2 displays some exposed residues belonging to linear epitopes that are predicted to participate in conformational epitopes. This is the case of some exposed residues of epitope no. 13, no. 5 and no. 9 that are estimated to be part of a conformational epitope. Likewise, some residues from epitope no. 21 and no. 22 are also predicted to participate in a unique conformational epitope located at one extremity of the Ana o 2 protomer [80]. More recently, Robotham et al. [81] confirmed the first Ana o 2 conformational epitope by reactivity with monoclonal antibody 2B5, which is located on the large subunit and it is structurally dependent upon association with the small subunit.

With respect to this conformational epitope, both native and recombinant Ana o 2 reacted similarly upon denaturation despite the structural differences between forms [81, 82].

Lately, Kshirsagar et al. [83] estimated the native secondary structure of anacardein (Ana o 2) with 49 % of beta-sheets, 23.7 % of alpha-helices and 27.4 % of random coil (absence of regular secondary structure). Different denaturing conditions (guanidine-HCl, urea, sodium dodecyl sulphate, beta-mercaptoethanol and heat—100 °C, 5 min) were used to assess the effect of structural changes on the immunoreactivity of anacardein (Ana o 2). As a result, denaturing treatments with sodium dodecyl sulphate or beta-mercaptoethanol induced conformational changes, causing a significant reduction in the immunoreactivity of anacardein when tested with an antibody that target conformational epitopes [83]. The addition of oleic acid to Ana o 2 at a concentration of 5 mM and overnight incubation at 37 °C induced a 35 % reduction of the IgE-binding capacity of Ana o 2. This fact seems to suggest that the oleic acid can reduce the allergenic properties of cashew nut by binding to its major protein Ana o 2 [84].

Prolamin Superfamily

The prolamin superfamily is one of the most relevant families of proteins since it encompasses a great number of allergenic members. Common features, such as the presence of eight conserved residues of cysteine, high content in proline and glutamine residues (source of its denomination), low molecular weight, similar conformational structures rich in alpha-helices and great stability to thermal processing/proteolysis, are shared by the members of this superfamily [60]. Contrarily to legumes and nuts, prolamins are the main storage proteins in most cereal grains (except in rice and oat) and members of the grass family [60, 85]. The members of this superfamily include the 2S albumin seed storage proteins, the non-specific lipid transfer proteins and the cereal seed inhibitors of alpha-amylase and trypsin [62].

Ana o 3—2S Albumin

2S albumins represent a family of water-soluble proteins at low-salt concentrations, presenting a primary structure with high content in arginine, glutamine, asparagine and cysteine residues, sharing the same common features of prolamin members [86]. They are small globular proteins with 12–15 kDa that undergo sequence modifications after synthesis. Mature 2S albumins are composed by heterodimers, consisting of a large (8–10 kDa) and a small (3–4 kDa) subunits linked by two conserved inter-chain disulphide bonds [62, 87, 88]. These proteins have two additional intra-chain disulphide bonds, which stabilise and compact their structure. Besides intervening in seed

germination most likely as nitrogen and sulphur donors, some 2S albumins are also known to act as antifungal agents, as serine protease inhibitors and as calmodulin antagonists [89, 90]. They are considered an important family of proteins since they are responsible for most plant food allergies [91].

Ana o 3 is the allergenic 2S albumin identified and characterised in cashew nut [59, 63]. So far, one isoform, Ana o 3.0101, has been identified in cashew nut presenting 138 aa and a molecular size of approximately 14 kDa, which is encoded by a mRNA sequence of 585 bp (Table 2). Besides identifying a vicilin (Ana o 1) and a legumin (Ana o 2) in the soluble protein fraction of cashew nut, Teuber et al. [77] also isolated three small proteins with molecular sizes below 14 kDa (estimated to correspond to a large subunit) that were classified as 2S albumins. Owing to their high immunoreactivity with sera from cashew nut allergic patients (>73 %), these proteins were classified as major allergens, being designated as Ana o 3 [77]. Using competent *E. coli* BL21 (DE3) cells, Robotham et al. [92] identified a recombinant Ana o 3 after the purification of the fusion protein (57 kDa) with a 43-kDa maltose-binding protein. The primary structure of Ana o 3 was evaluated with known sequences of other nut 2S albumins, revealing high degree of identity and similarity with Pis v 1 (pistachio), Jug n 1 (black walnut), Jug r 1 (English walnut) and Car i 1 (pecan nut), amongst others [92, 93]. Ana o 3 evidenced strong IgE reactivity with sera from 21 out of 26 patients with clear medical history of moderate to severe allergic reactions upon cashew nut ingestion [92], thus confirming its previous classification as major allergen [77].

More recently, Reitsma et al. [67] evaluated Ana o 3 by SDS-PAGE in denaturing reducing conditions, allowing detecting three distinct bands at 10, 8 and 6 kDa. Native and denatured reduced Ana o 3 triplet was further analysed by liquid chromatography-mass spectrometry (LC-MS). The bands of 10 and 8 kDa were identified as the large subunit presenting 8.2–8.4 and 8.1–8.2 kDa, respectively. The band of 6 kDa was recognised as the small subunit of 3.7–4.5 kDa [67]. Contrarily to Robotham et al. [92], who described that all three bands were isoforms of the large subunit of Ana o 3 with very similar N-terminal sequences, Reitsma et al. [67] evidenced that the small and the large subunits of Ana o 3 presented N- and C-terminal micro-heterogeneity and C-terminal micro-heterogeneity, respectively. This feature is also common to other allergenic 2S albumins, namely in Ber e 1 (Brazil nut) and Ses i 1 (sesame) [94, 95]. After sequencing, epitope mapping of Ana o 3 was performed using the sera of 4 cashew nut allergic patients in a total of 42 overlapping peptides, covering the entire length of the protein. From those, 12 peptides were IgE-reactive to the serum of at least one cashew nut allergic patient, with the most strongly reactive peptides being located at the large subunit of Ana o 3 [67].

Effect of Processing and Digestibility on Cashew Nut Allergens

Presently, there is a wide variety of technologies available for food processing aiming at improving the organoleptic attributes and/or shelf-life of the product, which is mostly achieved by removing/inactivating microbial activity and by altering the properties of different food components [96]. Accordingly, proteins involved in sensitising/eliciting allergic reactions may undergo extensive modification during food processing and/or may be present within complex structures in food. Due to the diversified nature of the allergenic proteins (e.g. albumins, vicilins, legumins, profilins, nsLTP), their immunoreactivity is associated with the presence of conformational and/or linear epitopes [97, 98]. If on one side some thermal treatments are known to reduce the immunoreactivity of proteins, such as profilins and vicilins, the same processes cause no effect on proteins, such as 2S albumins or legumins [98]. On the other side, the formation of novel IgE-binding sites as a consequence of allergen unfolding (thermal treatment) or chemical/enzymatic modifications can increase their immunoreactivity [97].

In the specific case of cashew nut, few reports describing the effect of different food processing treatments on its immunoreactivity are available in the literature [70, 99, 100]. de Leon et al. [70] assessed the effect of roasting (thermal treatment) on the immunoreactivity of cashew nut, verifying that roasting at 180 °C for 15 min did not affect the IgE-binding capacity of cashew nut allergens. Su et al. [99] evaluated the effect of other processing treatments, namely the application of gamma irradiation or thermal conditions and the combination of both, on the immunoreactivity of cashew nut allergens. From different thermal conditions that included autoclaving, blanching, frying, microwave and roasting, varying the time and temperature in each of the procedures, only roasting at 160 °C for 30 min reduced IgE-binding of Ana o 2 with rabbit anti-cashew nut polyclonal antibodies. IgE reactivity was also strongly reduced in cashew nut subjected to gamma irradiation at 25 kGy followed by autoclaving at 121 °C during 30 min [99].

Venkatachalam et al. [100] also tested the effect of different thermal (autoclaving, blanching, microwave heating and dry roasting—varying time/temperature) and pH (1, 3, 5, 7, 9, 11 and 13) conditions as well as gamma irradiation (1–25 kGy) on the immunoreactivity of cashew nut allergens (Ana o 1, Ana o 2 and Ana o 3). Two monoclonal antibodies (mAb 4B7 and mAb 1H2) were used to evaluate the reactivity of Ana o 1. In samples submitted to thermal treatments, the monoclonal antibody (mAb 4B7) did not detect any band at ~50 kDa, although the immunoreactivity was not considerably affected by gamma irradiation. Contrarily, using the monoclonal antibody (mAb 1H2), Ana o 1 was detected in all samples, although the intensity of the IgE-binding was decreased in the

sample autoclaved for 20 min. Therefore, Ana o 1 is most likely to lose its conformational structure upon heat treatments, although preserving linear epitopes that remain IgE-reactive. Ana o 1 seems to be stable to pH alterations since immunoreactivity was only affected in the sample submitted to pH 13. However, the significance of Ana o 1 instability at this level of pH is minimal, considering that most methods for cashew nut processing do not use pH 13 [100].

The immunoreactivity of Ana o 2 after processing simulations was also assessed by one polyclonal (pAb) and two monoclonal (mAb 4C3 and mAb 4H9) antibodies. Extreme processing conditions (e.g. autoclaving for 30 min or pH 1) seemed to enable minor reductions on the IgE reactivity by pAb-based ELISA (enzyme-linked immunosorbent assay). The immunoreactivity of Ana o 2 using mAb 4H9 or mAb 4C3 was very similar to the results from pAb, with minor IgE-binding reduction at extreme pH conditions (pH 1–3 or 13). Regardless the type of assay or antibody used, Ana o 2 seems to preserve its structural integrity, which is normally expected in this family of proteins (legumins) [100].

The monoclonal antibody (mAb D2) used for Ana o 3 recognition did not bind any of its known linear epitopes, suggesting that it probably recognises a conformational epitope or a potential linear motif that has not yet been identified in this allergen. Ana o 3 is stable to a pH range of 1–11 but not detectable at pH 13. Thermal processing, such as autoclaving, blanching or roasting for 10 min, significantly decreased the immunoreactivity of Ana o 3 when tested with mAb D2 by ELISA, which might be related to some structural alterations (e.g. denaturation) of Ana o 3 [100].

In spite of the distinct nature of cashew nut allergens (vicilin, legumin and 2S albumin), the majority of the processing conditions did not affect or have little effect on the allergenicity of Ana o 1, Ana o 2 and Ana o 3. These results seem to confirm their great potential for eliciting adverse immunological responses in sensitised/allergic patients, even after extensive food processing. Moreover, a general increase in the intensity of the bands, which is an indicative of increased immunoreactivity, was observed in samples subjected to roasting at 170 °C for 20 min. These results might also indicate that some processing conditions can contribute to increase, rather than reduce, the immunoreactivity of cashew nut allergens [100].

More recently, Mattison et al. [101] reported that treating cashew nut with sodium sulphite disrupted the conformational structure of Ana o 3, in a temperature-dependent manner. The structural integrity of Ana o 2 was also disrupted in cashew nut extracts treated with high concentrations of sodium sulphite (≥ 50 mM) at high temperatures (≥ 100 °C). In general, the immunoreactivity of Ana o 2 and Ana o 3 was greatly reduced with high concentrations of sodium sulphite (≥ 50 mM) and high temperatures (≥ 100 °C). In the same study, the authors suggest that sulphating agents (e.g. sodium

sulphite) seem to act at multiple levels in the reduction of IgE-binding to allergens. Accordingly, by targeting the disulphide bonds and negatively charging the thiol group, conformational epitopes are destroyed and linear ones with charged thiols are most likely to be repelled from IgE-binding sites. As a result, the incorporation of sodium sulphite or other food-grade reagents with similar redox potential may be useful to reduce or eliminate the allergenicity of some proteins [101].

Based on current knowledge relating to gastrointestinal digestion and food allergies, *in vitro* assays simulating the *in vivo* digestion have been used as predictive tools to assess the allergenic potential of novel food proteins [102]. In spite of the increasing number of works detailing the effect of the simulated gastrointestinal digestion on specific allergenic foods (e.g. milk, egg, crustaceans), only a report could be found targeting cashew nut allergens. To mimic the gastric and intestinal digestion of cashew nut proteins, Mattison et al. [103] used *in vitro* assays to assess the effect of gastric and intestinal fluids on defatted cashew nut flour. Some Ana o 2 peptides were resistant to digestion and remained immunoreactive, whilst Ana o 3 peptides seemed to lose some capacity to bind IgE. However, the study also suggested that Ana o 3 was only partially digested. Ana o 3 is most likely to resist to both gastric and intestinal digestion, which is consistent with the stability of other 2S albumins, thus preserving their immunoreactivity when reaching the lumen [103].

Management and Traceability of Cashew Nut

Food allergies are considered as an important public health problem with major impact on the quality of life of sensitised/allergic individuals [104], representing a current societal challenge. Faced as a multidisciplinary task, the management of food allergies requires the involvement of multiple actors, namely regulatory authorities, different stakeholders (food industry, clinicians and caretakers) and sensitised/allergic consumers [105]. Acting at different levels, all entities share a common feature of protecting the health of sensitised/allergic individuals. The regulatory authorities and food industry operate at population level, being responsible for guaranteeing the protection of public health and for providing safe foods for all intended consumers, respectively. However, the ultimate responsibility lies at the individual level, specifically on the sensitised/allergic consumers, since they are accountable for all preventive measures regarding the total avoidance of the allergenic food (and/or cross-reactive ones) [105]. Therefore, the allergic consumers are fully dependent on the label information of processed foods, whilst carefully selecting their nutrient sources. According to recent legislation, ingredients belonging to the 14 groups of potentially allergenic foods must be highlighted in the label through a typeset (font, style or background colour) that clearly

distinguishes them from the rest of the list of ingredients, independently of their quantity [11, 12].

Currently, food industry practices precautionary labelling (e.g. may contain traces of cashew nut) under global legal directives [11, 12]; therefore, most of the allergic individuals are protected against the consumption of allergenic foods [106]. However, accidental exposure to hidden allergens in foods owing to mislabelling or cross-contaminations during food processing (e.g. shared production lines) constitutes a real risk for allergic consumers [107, 108]. To avoid potential health risks, proper and highly sensitive analytical methodologies represent essential assets to aid the industrial management of allergenic foods and subsequently to facilitate allergen control/monitoring by the regulatory authorities.

Despite some contradictory opinions regarding the best target analyte for allergen detection/quantification in foods, there is a wide spectrum of analytical methods based on proteins and/or DNA analysis for most of the allergenic food commodities. Nonetheless, the lack of available official methods and testing/reference materials for their detection/quantification represents a major shortcoming in food allergen management [107, 108]. The choice of the best method for allergen analysis depends on specific criteria, such as target analyte (proteins

versus DNA), basis of detection (e.g. chemical), cost per run/analysis, setup cost, cross-reactivity, the need for expertise knowledge and possibility for multi-target detection [109]. For cashew nut analysis, there are several commercial and in-house-based methods either targeting allergenic/marker proteins or respective encoding DNA (Fig. 2).

Protein-Based Methods

Considering the advantages and drawbacks associated with all analytical methods, the classical protein-based methods (lateral flow devices (LFD), dipstick tests, ELISA, immunoblotting) are the most commonly used for food allergen analysis. However, other cutting-edge technologies, such as the immunosensors and the MS platforms, have also been developed and applied to allergen detection/quantification in foods.

ELISA and LFD

Based on typical antigen-antibody interaction, ELISA and LFD offer good sensitivity, adequate specificity and simple sample handling. ELISA are considered semi-quantitative or quantitative tools, being the most widely used analytical




Fig. 2 Representative protein- and DNA-based methods for the detection and quantification of cashew nut allergens in foods

technique for routine analysis and screening of allergens in foods [110]. LFD are simplified versions of ELISA [110], which are also quite popular in the food industry, being commonly used to check potential cross-contaminations along the production lines or in the final formulations [107].

Presently, ELISA kits and LFD can be commercially acquired, enabling the detection of a great number of different allergenic foods. Table 3 summaries the available ELISA kits and LFD for the specific detection of cashew nut in foods. These assays can be applied to a wide spectrum of food matrices with alleged sensitivities of 1–2 or 0.2–1 mg/kg of cashew nut in foods by LFD or ELISA, respectively. However, as expected for this type of assays, some cross-reactivity occurs between cashew nut and other plants, namely with Brazil nut, pistachio, hazelnut, peanut and walnut (Table 3).

Besides commercial kits, some in-house developed ELISA have been proposed for the detection of cashew nut in complex and processed food matrices, namely spices, cereals, chocolates, ice creams and cookies (Table 4). ELISA sandwich type is the most commonly applied, although other formats can be used (e.g. direct ELISA, indirect ELISA, competitive ELISA). The reported limits of detection (LOD) for these assays are aligned with commercial kits, ranging from 0.1 to 1.0 mg/kg depending on the type of food matrix [111–116]. ELISA systems can also be performed as a multi-target approach using the combination of different antibodies in single assays. Accordingly, an indirect multi-target ELISA was successfully proposed by Rejeb et al. [115] for the simultaneous detection of peanut and four nuts (cashew nut, almond, hazelnut and Brazil nut) in milk and dark chocolates (Table 4). In the reported study, the multi-target ELISA enabled detecting down to 1 mg/kg of cashew nut in milk and dark chocolates [115].

A different commercial immunoassay based on magnetic beads was performed by Cho et al. [116] for the simultaneous detection of 14 allergenic foods plus gluten. In the described method, each allergenic food was combined with microspheres and targeted by two capture antibodies, detecting raw cashew nut with some cross-reactivity towards hazelnut, pistachio and Brazil nut.

In spite of all the advantages attributed to immunoassays, there are several drawbacks that cannot be ruled out. Both LFD and ELISA are prone to cross-reactivity issues that might occur from matrix effects that lead to false positive results and, upon harsh food processing conditions (heat treatments, glycation, pH alterations, formation of Maillard products, fermentation, partial hydrolysis), false negative results are likely to occur [107].

Immunosensors

Presently, a great attention has been devoted to biosensing technology, being regarded as one of the most promising approaches for fast, reproducible, simple and low-cost multi-

target analysis of food allergens. Along with their potential for automation and high speed of execution, biosensors have been faced as tools with great potential for real-time allergen monitoring at an industrial scale [121]. In concept, biosensors are based on the direct recognition of a biological interaction between a receptor and a target molecule (proteins or DNA) by means of a transducer that produces a measurable signal [110, 122]. This interaction can be registered by different transducers (optical, piezoelectric or electrochemical) that are capable of acquiring the signal and further process it to give a proportional output to the concentration of a specific target [110].

So far, some biosensors using antibodies as biological receptors (also known as immunosensors) have been suggested for food allergen analysis [123]. Rebe Raz et al. [117] developed a sensor chip as a multi-target approach for the detection of 12 allergenic foods, including cashew nut. The antibodies targeting each allergenic food were immobilised onto a sensor chip as a microarray platform, and the signal acquisition was performed by an optical transducer (imaging surface plasmon resonance—iSPR) without requiring labelling or signal amplification. In the proposed microarray, it was possible to detect cashew nut down to 0.4 and 0.9 mg/kg in cookies and chocolates, respectively (Table 4). The sensitivity of the immunosensor was in the same range of the ELISA kits (Table 3) and of the in-house developed ELISA (Table 4), being suggested as excellent alternatives for multi-target allergen detection.

Mass Spectrometry

The continuous and rapid evolution of proteomic approaches has turned them as efficient tools for the characterisation of food-derived proteins. Owing to the impressive increase in the performance and versatility of MS instrumentation, the development of novel analytical strategies for the identification, characterisation and determination of allergens represents a valuable platform for food allergenomics [124–126]. MS platforms are normally coupled to a LC system that counts several advantages, such as high accuracy, sensitivity, specificity and reproducibility, which have greatly contributed for their application in the field of food allergen analysis [126, 127]. In addition, MS technology allows direct detection of the target peptide/protein without requiring an interaction with a biological receptor (e.g. antibody), which eliminates the problem related to cross-reactivity phenomena and enables the unequivocal identification of target analytes [126].

So far, three MS-based applications have been reported for the detection and quantification of cashew nut in foods, which are listed in Table 4. The methods were developed as multi-target approaches, enabling cashew nut identification and discrimination from other nuts (e.g. almond, walnut, hazelnut) in different food matrices (biscuits, chocolates, cookies, cakes and flours) [118–120]. At least, two marker peptides from

Table 3 Protein-based (ELISA, LFD) and DNA-based (real-time PCR) kits commercially available for the detection/quantification of cashew nut in foods

Commercial kit (brand)	Assay type (cat. no.)	Cross-reactivity	Dynamic range	LOD (mg/kg)	LOQ (mg/kg)	Performance time
Lateral Flow Cashew Kernel (r-Biopharm AG, Darmstadt, Germany)	Qualitative LFD (BL610-25)	10 % Brazil nut, 1 % hazelnut, 0.01 % peanut (roasted/boiled), 1 % walnut	–	1	–	10 min (incubation)
AgraStrip® Cashew/Pistachio (Romer Labs Division, Getzersdorf, Austria)	Qualitative LFD (COKAL1310AS)	No information available	2–10,000 mg/kg	2	–	11 min (including extraction)
Cashew Protein Rapid Test (Elution Technologies, Colchester, VT, USA)	Qualitative LFD (CASH-1013)	No cross-reactivity with tested samples	–	2	–	10 min
Reveal for multi-Treenut (Neogen Corporation, Lansing, MI, USA)	Qualitative hand-held device test (detects almond, cashew nut, hazelnut, pecan, pistachio, walnut) (8555)	–	5–10 mg/kg	–	–	10 min
AgraQuant® ELISA Cashew (Romer Labs Division, Getzersdorf, Austria)	Quantitative Sandwich ELISA (COKAL3148)	No information available	2–60 mg/kg	0.2	2	~60 min (total incubation time)
Cashew ELISA Kit (Arigo Biolaboratories Corp, Hsinchu, Taiwan)	Sandwich ELISA (ARG80791)	No cross-reactivity with 52 plant and animal species	2–60 mg/kg	0.2	2	~60 min (total incubation time)
AgraQuant® Plus Cashew (Romer Labs Division, Getzersdorf, Austria)	Quantitative Sandwich ELISA (COKAL3148F)	No information available	1–25 mg/kg	1	1	~30 min
Cashew Check (Bio-Check Ltd, Denbighshire, UK)	Qualitative ELISA (R-6046, R-6047)	No information available	2–50 mg/kg	<0.2	–	~60 min (total incubation time)
Cashew ELISA kit (BioFront Technologies, Tallahassee, FL, USA)	Quantitative ELISA (CA2-EK-48/96)	No cross-reactivity with other species (including all tree nuts)	1–40 mg/kg	0.13	1	10 min extraction 30 min testing time
Cashew Protein ELISA Kit (Elution Technologies, Colchester, VT, USA)	Sandwich ELISA (E-75CSH)	No cross-reactivity with 32 plant species	243–0.9 mg/kg	0.9	0.9	~50 min
Immunolab Cashew ELISA (Immunolab GmbH, Kassel, Germany)	Sandwich ELISA (CAW-E01)	4 % pistachio	2–60 mg/kg	0.2	2	~60 min (total incubation time)
SureFood® ALLERGEN ID Cashew (R-Biopharm AG Darmstadt, Germany)	Qualitative real-time PCR (S3115)	No cross-reactivity with 18 plants (including pistachio)	1–100,000 DNA copies	≤0.4 (in non-processed matrices)	–	~2 h
SPECIALfinder Cashew (Anacardo) (Generon, Modena, Italy)	Qualitative real-time PCR (PAV08A-PAV08A-50)	No cross-reactivity with 124 plant and animal species (including tree nuts)	–	≤5 DNA copies	–	~1.5 h
CashewKit Real-Time PCR (4Lab Diagnostics, Vicomosciano, Italy)	Qualitative real-time PCR (JC-02-1014)	No cross-reactivity with 20 plant and animal species (including tree nuts)	1–1000 cashew nut DNA copies	1 DNA copy	–	~2 h

LOD limit of detection, LOQ limit of quantification

Table 4 Protein-based techniques (ELISA, LC-MS and immunosensor) for cashew nut detection in foods

Method	Antibody/target	Cross-reactivity	Sensitivity level	Food matrices	References
Sandwich ELISA	Rabbit and goat IgG against anacardoin	Minor reactivity with sunflower seeds, pistachio, walnut and pecan nut	LOD, 1 mg/kg	Fortified wheat flours, cereals, spices and chocolate	[111, 112]
Sandwich ELISA	Polyclonal sheep and rabbit antibodies against roasted cashew nut	Cross-reactivity with hazelnut and pistachio from a panel of 99 food/ingredients	LOQ, 1 mg/kg	Commercial foods (Indian snacks) Spiked cookies, chocolate and ice cream	[113]
Sandwich ELISA	Polyclonal sheep, goat and rabbit IgG against raw and roasted defatted cashew nut	Cross-reactivity with hazelnut and pistachio from 101 ingredients (spices, tree nuts, fruits amongst others)	LOD, 0.11 mg/kg, LOQ, 0.46 mg/kg	Spiked sugar cookies, milk chocolate and vanilla ice cream. Commercial foods	[114]
Multi-allergen screening ELISA	Polyclonal rabbit antibodies against cashew nut proteins	No cross-reactivity with 23 plant or animal species (including other nuts)	LOD, 1 mg/kg	Fortified milk and dark chocolates	[115]
Multiplex magnetic beads	Not available	Potential reactivity with almond and pistachio.	LOD, 5 mg/mL	Raw cashew nut. Other raw nuts and reference materials	[116]
Antibody-microarrayed chip using iSPR	Polyclonal antibodies against cashew nut	Strong cross-reactivity with pistachio. Moderate reactivity with hazelnut, Brazil nut and almond	Cookies: LOD, 0.4 mg/kg LOQ, 2.3 mg/kg Chocolate: LOD, 0.9 mg/kg LOQ, 3.1 mg/kg MS ² mode: LOD, 14 mg/kg LOQ, 46 mg/kg MS ³ mode: LOD, 30 mg/kg LOQ, 98 mg/kg	Fortified cookies and chocolates	[117]
LC-ESI-LIT-MS ² vs LC-ESI-LIT-MS ³	Ana o 2 (2 peptides)	No cross-reactivity observed with almond, hazelnut, walnut and peanut		Model biscuits	[118]
LC-ESI-MS	Ana o 2 (2 peptides)	No cross-reactivity observed with almond, hazelnut, walnut and peanut	Biscuits: LOD, 0.5 mg/kg LOQ, 1.6 mg/kg Chocolates: LOD, 15 mg/kg LOQ, 50 mg/kg LOD, 1 mg/kg	Model chocolates and model biscuits	[119]
LC-MS/MS	Ana o 2 (2 peptides) Ana o 3 (2 peptides)	No cross-reactivity with other nuts		Biscuits, cookies, cakes and flours	[120]

LOD limit of detection, LOQ limit of quantification

allergenic proteins (Ana o 2, Ana o 3) of cashew nut were used for its unequivocal identification, showing no cross-reactivity phenomena with peptides from different nuts (Table 4). In general, the limits of detection and quantification (0.5–100 mg/kg) proposed by the MS methods were higher than those reported for ELISA kits (0.2–1 mg/kg) (Table 3) or in-house developed ELISA (0.1–1 mg/kg) (Table 4). Like for other protein-based methods (e.g. ELISA), the sensitivity attained with MS approaches was also affected by the food matrix (Table 4). Accordingly, LOD and LOQ were considerably higher when targeting cashew nut in difficult food matrices (such as chocolates) than in cookies/biscuits [119]. MS-based methods are excellent alternatives for direct allergen detection/quantification, but the high cost of MS platform maintenance/equipment and the need for specialised personnel are main factors to consider.

DNA-Based Methods

Presently, methodologies based on the detection of DNA for food allergen analysis have been regarded with increasing interest. DNA molecules are more likely to preserve their integrity upon harsh food processing conditions, when compared to proteins. Therefore, DNA-based methods have proved to be excellent alternatives to protein-based methods, especially when analysing highly processed foods. In spite of being considered as indirect approaches for allergen detection, DNA-based tools can be easily implemented in routine analyses and considered, at the same time, as confirmatory tools for the identification of allergenic commodities [107, 108, 128]. In recent years, DNA-based methods have emerged, demonstrating their potentialities for food allergen analysis.

Real-Time PCR

Methods based on the polymerase chain reaction (PCR) are amongst the most commonly used DNA approaches for food allergen analysis. With adequate setup cost, reasonable running time and moderate requirements for specialised equipment and personnel, PCR systems can be used for quantification purposes (e.g. real-time PCR). PCR-based techniques are also less prone to cross-reactivity phenomena by carefully selecting the target sequence [107, 108], thus being independent of possible biological effects associated with antibody production.

Presently, some real-time PCR kits can be commercially acquired for food allergen analysis, although at a much smaller scale than the immunoassays (e.g. ELISA or LFD). Table 3 summarises most of the available real-time PCR kits for the specific detection of cashew nut in foods. These kits can be tested in a variety of foods with sensitivities that are strongly dependent on the type of food matrix analysed. Nonetheless, the listed kits alleger sensitivities down to one to five DNA

genomic copies, which can be comparable to the sensitivities stated in ELISA kits (Table 3). As expected, by the high specificity of these kits, no cross-reactivity phenomena are reported for cashew nut.

In addition to the commercial kits, some in-house developed real-time PCR systems have been proposed for the detection of cashew nut in raw and processed foods (Table 5) [129–132]. The spectrum of analysed food samples includes chocolates, biscuits, ice creams, spreads and pesto, amongst others. For quantitative purposes, most of the developed real-time PCR systems targeted single copy genes (Ana o 1, Ana o 2 or Ana o 3) (Table 5), since they are considered the best choices owing to the more constant copy number in different cells from distinct tissues [122]. However, the sensitivities of the systems targeting single copy genes are frequently lower (~100 mg/kg of cashew nut in matrix) than those using multi-copy genes (e.g. internal transcribed spacer) (~0.1 mg/kg of cashew nut in matrix) (Table 5). As for the commercial kits, the reported real-time PCR systems also present high specificity for cashew nut without cross-reactivity with other plant and animal species. Two multiplex real-time PCR have been proposed as multi-target platforms, enabling detecting cashew nut from a wide variety of food matrices with high specificity and sensitivity (Table 5) [133, 134].

LPA

Ligation-dependent probe amplification (LPA) has been proposed for food analysis, particularly for allergen detection. Basically, the method consists on the amplification of PCR products that result from the ligation of bipartite specific hybridisation probes (normally two) and uses a single pair of PCR primers to amplify several target sequences. A fragment of unique size is produced after amplification, being discriminated by capillary electrophoresis. The amplification efficiency is similar for all fragments because they are products from the same set of primers, resulting in a competitive PCR [122].

A LPA was successfully proposed by Ehlert et al. [135] for the simultaneous detection of 10 allergenic foods (peanut, cashew nut, pecan, pistachio, hazelnut, sesame, macadamia nut, almond, walnut and Brazil nut) in different matrices (cookies, pesto and chocolates). The specificity of the probes was evaluated with 56 plant and animal species, thus ensuring the absence of unspecific amplifications as result of cross-reactivity. In the case of cashew nut, the multiplex LPA system enabled its detection down to a level of 5 mg/kg in pesto matrix (Table 5).

Genosensors/Microarrays

DNA-biosensors (also known as genosensors) have become excellent tools for the detection of sequence-specific DNA fragments, mainly due to their simplicity, low-cost, automation

Table 5 DNA-based techniques (real-time PCR, LPA, microarray) for cashew nut detection in foods

Method	Target gene	Cross-reactivity	Matrices	Sensitivity level	References
Real-time PCR (TaqMan probe)	2S albumin (Ana o 3)	No cross-reactivity with walnut, peanut almond, hazelnut and Brazil nut	Model mixtures of cashew nut in chocolate cookies	5 pg cashew nut DNA 100 mg/kg cashew nut in chocolate cookies	[129]
Real-time PCR (TaqMan probe)	2S albumin (Ana o 3)	No cross-reactivity with 17-plant species (including tree nuts)	Model mixtures of cashew nut in pistachio nougat. Commercial foods (snacks, chocolate, breakfast cereals, biscuits)	1.25 pg of cashew nut DNA 100 mg/kg of cashew nut in pistachio nougat	[130]
Real-time PCR (TaqMan probe)	Ana o 3	No cross-reactivity with 56 plant and animal species (including tree nuts)	Model mixtures of cashew nut in basil pesto. Commercial foods (pesto, ice creams, sweets, cookies, spreads)	0.5 pg cashew nut DNA 2 mg/kg of cashew nut in basil pesto	[131]
Real-time PCR (TaqMan probe)	Internal transcribed spacer (ITS1)	No cross-reactivity with 49 plant and animal species	Model mixtures of raw or heat-processed cashew nut in wheat flour. Commercial foods (chocolates, bread, biscuits, cereals, meat products)	0.1 mg/kg of raw or heat-processed cashew nut in wheat flour	[132]
Real-time PCR (SYBR GreenER multiplex system)	Ana o 1	No cross-reactivity with 13 plant species	Cashew nut in different nuts. Commercial foods (biscuits, chocolates and crackers)	0.5 pg cashew nut DNA	[133]
Real-time PCR (TaqMan probe, multiplex system)	Ana o 3	Minor cross-reactivity with pistachio from a panel of 42 plant and animal species	Model mixtures of cashew nut in boiled sausages and rice cookies. Commercial foods (chocolates, sandwiches, spreads)	32 and 5 mg/kg of cashew nut in boiled sausages and rice cookies, respectively	[134]
Ligation-dependent probe amplification (LPA) (multiplex system)	Ana o 3	No cross-reactivity with 56 plant and animal species (including tree nuts)	Model mixtures of cashew nut in pesto. Commercial foods (pesto, ice creams, sweets, cookies, spreads)	5 mg/kg of cashew nut in pesto	[135]
Optical thin-film biosensor chips coupled to PCR (multiplex system)	Ana o 3	No cross-reactivity amongst the species used in biosensor (beef, shrimp, fish, chicken, soybean, wheat, peanut and cashew nut)	Model mixtures of cashew nut in maize powder. Commercial foods (biscuits, chocolate chips, cereals)	0.5 pg cashew nut DNA 10 mg/kg cashew nut in maize powder	[136]

and high selectivity/specificity. Genosensors share the same biosensing technology of immunosensors, although they are based on a different biological interaction. They rely on the hybridisation of two complementary strands: a probe and its complementary sequence (RNA or DNA), as the receptor and target analyte, respectively [122, 123]. Assembling genosensors in microarray platforms allows the simultaneous multiple target detection and thus the development of multiplex approaches.

A system of optical thin-film biosensor chips coupled to previous tetraplex PCR amplification was successfully developed for the simultaneous detection of eight allergenic ingredients (soybean, wheat, peanut, cashew nut, shrimp, fish, beef and chicken) in processed foods [136]. The proposed microarray enabled detecting down to 10 mg/kg of cashew nut in maize powder and 0.5 pg of genomic DNA (Table 5). No cross-reactivity was found amongst the species tested in microarray, being applied to a wide variety of commercial foods.

Final Remarks

Over the last years, cashew nut allergy has been regarded with special attention since its prevalence seems to be increasing in the industrialised countries. So far, three groups of seed storage allergenic proteins (Ana o 1—vicilin, Ana o 2—legumin and Ana o 3—2S albumin) have been identified and characterised in cashew nut. All proteins have been classified as major allergens of cashew nut, being included in the WHO/IUIS official list of allergens. As for other tree nuts, minute amounts of cashew nut proteins (1 mg) can induce an adverse immunological response in allergic individuals.

Allergic reactions to cashew nut are commonly severe and near fatal (anaphylaxis), often demanding epinephrine treatment and hospitalisation, especially in children. The intensity and severity of allergic reactions are most difficult to predict due to the influence of several internal and external factors. Therefore, eliminating the allergenic foods from the diet is still the most efficient means of preventing allergic reactions in sensitised/allergic individuals. Accordingly, when carefully electing the processed foods that are commercially available, the allergic individuals have to rely on proper food labelling. Correct label information represents a critical tool for allergen management. In this sense, the control of labelling compliance is much needed, which has prompted the development of proficient analytical methods for allergen analysis.

Contrarily to other pathologies, there is no cure for food allergies. Although some strategies have been used to induce immune modulation and subsequent desensitisation, dietary eviction and the provision of rescue medication in the event of accidental exposures are the most commonly applied measures. Different forms of immunotherapies have been tested aiming at inducing long-term tolerance to the allergenic food.

Oral immunotherapies to induce tolerance to cashew nut are still very scarce, even though they are foreseen as a potential therapeutics for the treatment of cashew nut allergy.

In spite of the significant number of available reports concerning cashew nut allergy, much research is still required in this field, and new advances are expected in the near future.

Acknowledgments This work was supported by FCT/MEC through national funds and co-financed by FEDER, under the Partnership Agreement PT2020 with grant no. UID/QUI/50006/2013–POCI/01/0145/FEDER/007265. Joana Costa is grateful to FCT post-doctoral grant (SFRH/BPD/102404/2014) financed by POPH-QREN (subsidised by FSE and MCTES).

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflict of interest.

References

- GRIN-USDA, Germplasm Resources Information Network—United States Department of Agriculture, Beltsville, MD, USA. <http://www.ars-grin.gov/>. Accessed 12 Jan 2016.
- Blazdell P (2000) The mighty cashew. *Interdiscip Sci Rev* 25:220–226
- Shahidi F, Tan Z (2008) Bioactive compounds from cashew nut and its coproducts. In Alasalvar C and Shahidi F (eds) *Tree nuts: composition, phytochemicals, and health effects: an overview*. Taylor & Francis Group, Boca Raton, chapter 10.
- Nair KPP (2010) Cashew nut (*Anacardium occidentale* L.). In Nair KPP (ed) *Tree crops of the developing world*. Elsevier, Amsterdam, chapter 2.
- Nussinovitch A (2010) Major plant exudates of the world. In Nussinovitch A (ed) *Plant gum exudates of the world: sources, distribution, properties, and applications*. Taylor & Francis Group, Boca Raton, chapter 3.
- FAOSTAT, Food and Agriculture Organization of the United Nations, Statistics Division, Economic and Social Development Department, Rome, Italy. <http://faostat3.fao.org/home/E>. Accessed 12 Jan 2016.
- Ros E (2010) Health benefits of nut consumption. *Nutrients* 2:652–682. doi:10.3390/nu2070652
- FDA (2003), Administration Qualified Health Claims, letter of enforcement discretion—nuts and coronary heart disease (Docket No 02P-0505), Ingredients, Packaging & Labeling, Food and Drug Administration, Washington, DC, USA. <http://www.fda.gov/Food/IngredientsPackaging/Labeling/LabelingNutrition/ucm072926.htm>. Accessed 3 February 2016.
- CODEX STAN 1 (1985) Amended in 1991, 1999, 2001, 2003, 2005, 2008 and 2010 regarding the general standard for the labelling of pre-packaged foods. FAO/WHO standards. *Off Codex Stand.* <http://www.codexalimentarius.org/standards/list-of-standards/> Accessed 3 Feb 2016.
- Directive 2003/89/EC of 10 November 2003 amending Directive 2000/13/EC regarding indication of the ingredients present in foodstuffs. *Off J Eur Union* L308:15–18.

11. Directive 2007/68/EC of 27 November 2007 amending Annex IIIa to Directive 2000/13/EC regarding certain food ingredients. *Off J Eur Union* L310:11–14.
12. Regulation (EU) No 1169/2011 of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No 1924/2006 and (EC) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/2004. *Off J Eur Union* L304:18–63.
13. Gendel SM (2012) Comparison of international food allergen labeling regulations. *Regul Toxicol Pharmacol* 63:279–285. doi:10.1016/j.yrtph.2012.04.007
14. Taylor SL, Baumert JL (2015) Worldwide food allergy labeling and detection of allergens in processed foods. *Chem Immunol Allergy* 101:227–234. doi:10.1159/000373910
15. Zuidmeer L, Goldhahn K, Rona RJ et al (2008) The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol* 121:1210–1218. doi:10.1016/j.jaci.2008.02.019
16. McWilliam V, Koplin J, Lodge C, Tang M, Dharmage S, Allen K (2015) The prevalence of tree nut allergy: a systematic review. *Curr Allergy Asthma Rep* 15:1–13. doi:10.1007/s11882-015-0555-8
17. Sicherer SH, Furlong TJ, Munoz-Furlong A, Burks AW, Sampson HA (2001) A voluntary registry for peanut and tree nut allergy: characteristics of the first 5149 registrants. *J Allergy Clin Immunol* 108:128–132. doi:10.1016/j.jaci.2014.12.1906
18. Fleischer DM, Conover-Walker MK, Matsui EC, Wood RA (2005) The natural history of tree nut allergy. *J Allergy Clin Immunol* 116:1087–1093. doi:10.1016/j.jaci.2005.09.002
19. Johnson J, Malinowski A, Alving K, Lidholm J, Borres MP, Nordvall L (2014) Ten-year review reveals changing trends and severity of allergic reactions to nuts and other foods. *Acta Paediatr* 103:862–867. doi:10.1111/apa.12687
20. Vetander M, Helander D, Flodström C et al (2012) Anaphylaxis and reactions to foods in children—a population-based case study of emergency department visits. *Clin Exp Allergy* 42:568–577. doi:10.1111/j.1365-2222.2011.03954.x
21. Le TM, Lindner TM, Pasmans SG et al (2008) Reported food allergy to peanut, tree nuts and fruit: comparison of clinical manifestations, prescription of medication and impact on daily life. *Allergy* 63:910–916. doi:10.1111/j.1398-9995.2008.01688.x
22. Moneret-Vautrin DA (2008) Épidémiologie de l'allergie alimentaire. *Rev Fr Allergol* 48:171–178. doi:10.1016/j.allerg.2008.01.018
23. De Silva IL, Mehr SS, Tey D, Tang MLK (2008) Paediatric anaphylaxis: a 5-year retrospective review. *Allergy* 63:1071–1076. doi:10.1111/j.1398-9995.2008.01719.x
24. Davoren M, Peake J (2005) Cashew nut allergy is associated with a high risk of anaphylaxis. *Arch Dis Child* 90:1084–1085. doi:10.1136/adc.2005.073817
25. Rancé F, Bidat E, Bourrier T, Sabouraud D (2003) Cashew allergy: observations of 42 children without associated peanut allergy. *Allergy* 58:1311–1314. doi:10.1046/j.1398-9995.2003.00342.x
26. Sicherer SH, Leung DYM (2015) Advances in allergic skin disease, anaphylaxis, and hypersensitivity reactions to foods, drugs, and insects in 2014. *J Allergy Clin Immunol* 135:357–367. doi:10.1016/j.jaci.2014.12.1906
27. Boyce JA, Assa'ad A, Burks AW et al (2010) Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol* 126:S1–S58. doi:10.1016/j.jaci.2010.10.007
28. Valenta R, Hochwallner H, Linhart B, Pahr S (2015) Food allergies: the basics. *Gastroenterology* 148:1120–1131.e1124. doi:10.1053/j.gastro.2015.02.006
29. Gómez E, Mayorga C, Gómez F et al (2013) Food allergy: management, diagnosis and treatment strategies. *Immunotherapy* 5:755–768. doi:10.2217/imt.13.63
30. Lupinek C, Wollmann E, Baar A et al (2014) Advances in allergen-microarray technology for diagnosis and monitoring of allergy: the MeDALL allergen-chip. *Methods* 66:106–119. doi:10.1016/j.ymeth.2013.10.008
31. Clark AT, Anagnostou K, Ewan PW (2007) Cashew nut causes more severe reactions than peanut: case-matched comparison in 141 children. *Allergy* 62:913–916. doi:10.1111/j.1398-9995.2007.01447.x
32. van der Valk JPM, Gerth van Wijk R, Dubois AEJ et al (2016) Multicentre double-blind placebo-controlled food challenge study in children sensitised to cashew nut. *PLoS One* 11:1–12. doi:10.1371/journal.pone.0151055
33. Maloney JM, Rudengren M, Ahlstedt S, Bock SA, Sampson HA (2008) The use of serum-specific IgE measurements for the diagnosis of peanut, tree nut, and seed allergy. *J Allergy Clin Immunol* 122:145–151. doi:10.1016/j.jaci.2008.04.014
34. van der Valk JPM, Gerth van Wijk R, Hoorntje E, Groenendijk L, Groenendijk IM, de Jong NW (2016) Measurement and interpretation of skin prick test results. *Clin Transl Allergy* 6:8. doi:10.1186/s13601-016-0092-0
35. Savvatanos S, Konstantinopoulos AP, Borga A et al (2015) Sensitization to cashew nut 2S albumin, Ana o 3, is highly predictive of cashew and pistachio allergy in Greek children. *J Allergy Clin Immunol* 136:192–194. doi:10.1016/j.jaci.2015.03.037
36. Muraro A, Roberts G, Worm M et al (2014) Anaphylaxis: guidelines from the European Academy of Allergy and Clinical Immunology. *Allergy* 69:1026–1045. doi:10.1111/all.12437
37. Chowdhury BA, Meyer RJ (2002) Intramuscular versus subcutaneous injection of epinephrine in the treatment of anaphylaxis. *J Allergy Clin Immunol* 109:720. doi:10.1067/mai.2002.123252
38. Bégin P, Chinthrajah RS, Nadeau KC (2014) Oral immunotherapy for the treatment of food allergy. *Hum Vaccines Immunother* 10:2295–2302. doi:10.4161/hv.29233
39. Mousallem T, Burks AW (2012) Immunology in the clinic review series; focus on allergies: immunotherapy for food allergy. *Clin Exp Immunol* 167:26–31. doi:10.1111/j.1365-2249.2011.04499.x
40. Sato S, Yanagida N, Ogura K et al (2014) Clinical studies in oral allergen-specific immunotherapy: differences among allergens. *Int Arch Allergy Immunol* 164:1–9. doi:10.1159/000361025
41. Sindher S, Fleischer DM, Spergel JM (2016) Advances in the treatment of food allergy: sublingual and epicutaneous immunotherapy. *Immunol Allergy Clin North Am* 36:39–54. doi:10.1016/j.iac.2015.08.008
42. Kulis M, Li Y, Lane H, Pons L, Burks W (2011) Single-tree nut immunotherapy attenuates allergic reactions in mice with hypersensitivity to multiple tree nuts. *J Allergy Clin Immunol* 127:81–88. doi:10.1016/j.jaci.2010.09.014
43. Kulis M, MacQueen I, Li YF, Guo RS, Zhong XP, Burks AW (2012) Pepsinized cashew proteins are hypoallergenic and immunogenic and provide effective immunotherapy in mice with cashew allergy. *J Allergy Clin Immunol* 130:716–723. doi:10.1016/j.jaci.2012.05.044
44. Okada Y, Yanagida N, Sato S et al (2015) Oral immunotherapy initiation for multi-nut allergy: a case report. *Allergol Int* 64:192–193. doi:10.1016/j.alit.2014.09.004
45. Bégin P, Dominguez T, Wilson SP et al (2014) Phase I results of safety and tolerability in a rush oral immunotherapy protocol to multiple foods using omalizumab. *Allergy Asthma Clin Immunol* 10:1–10. doi:10.1186/1710-1492-10-7
46. El-Qutob D (2015) Off-label uses of omalizumab. *Clin Rev Allergy Immunol* 50:84–96. doi:10.1007/s12016-015-8490-y

47. Entink RHK, Remington BC, Blom WM et al (2014) Food allergy population thresholds: an evaluation of the number of oral food challenges and dosing schemes on the accuracy of threshold dose distribution modeling. *Food Chem Toxicol* 70:134–143. doi:10.1016/j.fct.2014.05.001
48. Crevel RWR, Baumert JL, Baka A et al (2014) Development and evolution of risk assessment for food allergens. *Food Chem Toxicol* 67:262–276. doi:10.1016/j.fct.2014.01.032
49. Taylor SL, Baumert JL, Kruizinga AG et al (2014) Establishment of reference doses for residues of allergenic foods: report of the VITAL expert panel. *Food Chem Toxicol* 63:9–17. doi:10.1016/j.fct.2013.10.032
50. Blom WM, Vlieg-Boerstra BJ, Kruizinga AG, van der Heide S, Houben GF, Dubois AEJ (2013) Threshold dose distributions for 5 major allergenic foods in children. *J Allergy Clin Immunol* 131:172–179. doi:10.1016/j.jaci.2012.10.034
51. Allen KJ, Remington BC, Baumert JL et al (2014) Allergen reference doses for precautionary labeling (VITAL 2.0): clinical implications. *J Allergy Clin Immunol* 133:156–164. doi:10.1016/j.jaci.2013.06.042
52. Radauer C, Bublin M, Wagner S, Mari A, Breiteneder H (2008) Allergens are distributed into few protein families and possess a restricted number of biochemical functions. *J Allergy Clin Immunol* 121:847–852.e847. doi:10.1016/j.jaci.2008.01.025
53. Jenkins JA, Griffiths-Jones S, Shewry PR, Breiteneder H, Mills ENC (2005) Structural relatedness of plant food allergens with specific reference to cross-reactive allergens: an in silico analysis. *J Allergy Clin Immunol* 115:163–170. doi:10.1016/j.jaci.2004.10.026
54. Harrer A, Egger M, Gadermaier G et al (2010) Characterization of plant food allergens: an overview on physicochemical and immunological techniques. *Mol Nutr Food Res* 54:93–112. doi:10.1002/mnfr.200900096
55. Moreno FJ (2007) Gastrointestinal digestion of food allergens: effect on their allergenicity. *Biomed Pharmacother* 61:50–60. doi:10.1016/j.biopha.2006.10.005
56. Moreno FJ, Clemente A (2008) 2S albumin storage proteins: what makes them food allergens? *Open Biochem J* 2:16–28. doi:10.2174/1874091X00802010016
57. van der Valk JPM, Dubois AEJ, Gerth van Wijk R, Wichers HJ, de Jong NW (2014) Systematic review on cashew nut allergy. *Allergy* 69:692–698. doi:10.1111/all.12401
58. Grigg A, Hanson C, Davis CM (2009) Cashew allergy compared to peanut allergy in a US tertiary care center. *Pediatr Asthma Allergy Immunol* 22:101–104. doi:10.1089/pai.2009.0017
59. ALLERGEN, World Health Organization and International Union of Immunological Societies (WHO/IUIS) Allergen Nomenclature Sub-committee official site for the systematic allergen nomenclature. <http://www.allergen.org/>. Accessed 18 Feb 2016.
60. Radauer C, Breiteneder H (2007) Evolutionary biology of plant food allergens. *J Allergy Clin Immunol* 120:518–525. doi:10.1016/j.jaci.2007.07.024
61. Breiteneder H, Mills ENC (2005) Plant food allergens—structural and functional aspects of allergenicity. *Biotechnol Adv* 23:395–399. doi:10.1016/j.biotechadv.2005.05.004
62. Breiteneder H, Radauer C (2004) A classification of plant food allergens. *J Allergy Clin Immunol* 113:821–830. doi:10.1016/j.jaci.2004.01.779
63. ALLERGOME, allergome database, the platform for allergen knowledge, Latina, Italy. <http://www.allergome.org/>. Accessed 18 Feb 2016.
64. Chapman MD, Pomés A, Breiteneder H, Ferreira F (2007) Nomenclature and structural biology of allergens. *J Allergy Clin Immunol* 119:414–420. doi:10.1016/j.jaci.2006.11.001
65. Wang F, Robotham JM, Teuber SS, Tawde P, Sathe SK, Roux KH (2002) Ana o 1, a cashew (*Anacardium occidentale*) allergen of the vicilin seed storage protein family. *J Allergy Clin Immunol* 110:160–166. doi:10.1067/mai.2002.125208
66. Guan XY, Noble KA, Tao YQ et al (2015) Epitope mapping of 7S cashew antigen in complex with antibody by solution-phase H/D exchange monitored by FT-ICR mass spectrometry. *J Mass Spectrom* 50:812–819. doi:10.1002/jms.3589
67. Reitsma M, Bastiaan-Net S, Sforza S et al (2016) Purification and characterization of *Anacardium occidentale* (cashew) allergens Ana o 1, Ana o 2, and Ana o 3. *J Agric Food Chem* 64:1191–1201. doi:10.1021/acs.jafc.5b04401
68. Barre A, Sordet C, Culerrier R, Rance F, Didier A, Rouge P (2008) Vicilin allergens of peanut and tree nuts (walnut, hazelnut and cashew nut) share structurally related IgE-binding epitopes. *Mol Immunol* 45:1231–1240. doi:10.1016/j.molimm.2007.09.014
69. Kulis M, Pons L, Burks AW (2009) In vivo and T cell cross-reactivity between walnut, cashew and peanut. *Int Arch Allergy Immunol* 148:109–117. doi:10.1159/000155741
70. de Leon MP, Glaspole IN, Drew AC, Rolland JM, O'Hehir RE, Suphioglu C (2003) Immunological analysis of allergenic cross-reactivity between peanut and tree nuts. *Clin Exp Allergy* 33:1273–1280. doi:10.1046/j.1365-2222.2003.01761.x
71. Fernandez C, Fiandor A, Martinez-Garate A, Quesada JM (1995) Allergy to pistachio: cross-reactivity between pistachio nut and other Anacardiaceae. *Clin Exp Allergy* 25:1254–1259. doi:10.1111/j.1365-2222.1995.tb03050.x
72. Garcia F, Moneo I, Fernandez B et al (2000) Allergy to Anacardiaceae: description of cashew and pistachio nut allergens. *J Invest Allergol Clin Immunol* 10:173–177
73. Hasegawa M, Inomata N, Yamazaki H, Morita A, Kirino M, Ikezawa Z (2009) Clinical features of four cases with cashew nut allergy and cross-reactivity between cashew nut and pistachio. *Allergol Int* 58:209–215. doi:10.2332/allergolint.08-OA-0010
74. Rougé P, Thibaut F, Bourrier T, Saggio B, Culerrier R, Rancé F, Barre A (2011) Clinical relevance of IgE-binding cross-reactivity of allergens from cashew nut (*Anacardium occidentale*). *Rev Fr Allergol* 51:31–35. doi:10.1016/j.reval.2010.10.005
75. Willison LN, Tawde P, Robotham JM et al (2008) Pistachio vicilin, Pis v 3, is immunoglobulin E-reactive and cross-reacts with the homologous cashew allergen, Ana o 1. *Clin Exp Allergy* 38:1229–1238. doi:10.1111/j.1365-2222.2008.02998.x
76. Mills EN, Jenkins J, Marigheto N, Belton PS, Gunning AP, Morris VJ (2002) Allergens of the cupin superfamily. *Biochem Soc Trans* 30:925–929. doi:10.1042/bst0300925
77. Teuber SS, Sathe SK, Peterson WR, Roux KH (2002) Characterization of the soluble allergenic proteins of cashew nut (*Anacardium occidentale* L.). *J Agric Food Chem* 50:6543–6549. doi:10.1021/jf025757j
78. Wang F, Robotham JM, Teuber SS, Sathe SK, Roux KH (2003) Ana o 2, a major cashew (*Anacardium occidentale* L.) nut allergen of the legumin family. *Int Arch Allergy Immunol* 132:27–39. doi:10.1159/000073262
79. Robotham JM, Hoffman GG, Teuber SS et al (2009) Linear IgE-epitope mapping and comparative structural homology modeling of hazelnut and English walnut 11S globulins. *Mol Immunol* 46:2975–2984. doi:10.1016/j.molimm.2009.06.020
80. Barre A, Jacquet G, Sordet C, Culerrier R, Rougé P (2007) Homology modelling and conformational analysis of IgE-binding epitopes of Ara h 3 and other legumin allergens with a cupin fold from tree nuts. *Mol Immunol* 44:3243–3255. doi:10.1016/j.molimm.2007.01.023
81. Robotham JM, Xia LX, Willison LN, Teuber SS, Sathe SK, Roux KH (2010) Characterization of a cashew allergen, 11S globulin (Ana o 2), conformational epitope. *Mol Immunol* 47:1830–1838. doi:10.1016/j.molimm.2009.12.009
82. Xia LX, Willison LN, Porter L et al (2010) Mapping of a conformational epitope on the cashew allergen Ana o 2: a discontinuous

- large subunit epitope dependent upon homologous or heterologous small subunit association. *Mol Immunol* 47:1808–1816. doi:10.1016/j.molimm.2010.01.018
83. Kshirsagar HH, Fajer P, Sharma GM, Roux KH, Sathe SK (2011) Biochemical and spectroscopic characterization of almond and cashew nut seed 11S legumins, amandin and anacardein. *J Agric Food Chem* 59:386–393. doi:10.1021/jf1030899
 84. Chung SY, Mattison CP, Reed S, Wasserman RL, Desormeaux WA (2015) Treatment with oleic acid reduces IgE binding to peanut and cashew allergens. *Food Chem* 180:295–300. doi:10.1016/j.foodchem.2015.02.056
 85. Shewry PR, Halford NG (2002) Cereal seed storage proteins: structures, properties and role in grain utilization. *J Exp Bot* 53:947–958. doi:10.1093/jexbot/53.370.947
 86. Breiteneder H, Ebner C (2000) Molecular and biochemical classification of plant-derived food allergens. *J Allergy Clin Immunol* 106:27–36. doi:10.1067/mai.2000.106929
 87. Pantoja-Uceda D, Bruix M, Santoro J, Rico M, Monsalve R, Villaba M (2002) Solution structure of allergenic 2S albumins. *Biochem Soc Trans* 30:919–924. doi:10.1042/bst0300919
 88. Shewry PR, Napier JA, Tatham AS (1995) Seed storage proteins—structures and biosynthesis. *Plant Cell* 7:945–956. doi:10.1105/tpc.7.7.945
 89. Terras FRG, Torrekens S, Van Leuven F et al (1993) A new family of basic cysteine-rich plant antifungal proteins from Brassicaceae species. *FEBS Lett* 316:233–240. doi:10.1016/0014-5793(93)81299-F
 90. Polya GM, Chandra S, Condron R (1993) Purification and sequencing of radish seed calmodulin antagonists phosphorylated by calcium-dependent protein kinase. *Plant Physiol* 101:545–551. doi:10.1104/pp.101.2.545
 91. Sampson HA, Aceves S, Bock SA et al. (2014) Food allergy: a practice parameter update—2014. *J Allergy Clin Immunol* 134:1016–1025.e43. doi: 10.1016/j.jaci.2014.05.013
 92. Robotham JM, Wang F, Seamon V et al (2005) Ana o 3, an important cashew nut (*Anacardium occidentale* L.) allergen of the 2S albumin family. *J Allergy Clin Immunol* 115:1284–1290. doi:10.1016/j.jaci.2005.02.028
 93. Ahn K, Bardina L, Grishina G, Beyer K, Sampson HA (2009) Identification of two pistachio allergens, Pis v 1 and Pis v 2, belonging to the 2S albumin and 11S globulin family. *Clin Exp Allergy* 39:926–934. doi:10.1111/j.1365-2222.2009.03259.x
 94. Moreno FJ, Jenkins JA, Mellon FA et al (2004) Mass spectrometry and structural characterization of 2S albumin isoforms from Brazil nuts (*Bertholletia excelsa*). *Biochim Biophys Acta* 1698:175–186. doi:10.1016/j.bbapap.2003.11.007
 95. Moreno FJ, Maldonado BM, Wellner N, Mills ENC (2005) Thermostability and in vitro digestibility of a purified major allergen 2S albumin (Ses i 1) from white sesame seeds (*Sesamum indicum* L.). *Biochim Biophys Acta* 1752:142–153. doi:10.1016/j.bbapap.2005.07.022
 96. Vanga SK, Singh A, Raghavan V (in press) Review of conventional and novel food processing methods on food allergens. *Crit Rev Food Sci Nutr*. doi: 10.1080/10408398.2015.1045965.
 97. Mills ENC, Mackie AR (2008) The impact of processing on allergenicity of food. *Curr Opin Allergy Clin Immunol* 8:249–253. doi:10.1097/ACI.0b013e3282ff123
 98. Mills ENC, Sancho AI, Rigby NM, Jenkins JA, Mackie AR (2009) Impact of food processing on the structural and allergenic properties of food allergens. *Mol Nutr Food Res* 53:963–969. doi:10.1002/mnfr.200800236
 99. Su M, Venkatachalam M, Teuber SS, Roux KH, Sathe SK (2004) Impact of gamma-irradiation and thermal processing on the antigenicity of almond, cashew nut and walnut proteins. *J Sci Food Agric* 84:1119–1125. doi:10.1002/jsfa.1748
 100. Venkatachalam M, Monaghan EK, Kshirsagar HH et al (2008) Effects of processing on immunoreactivity of cashew nut (*Anacardium occidentale* L.) seed flour proteins. *J Agric Food Chem* 56:8998–9005. doi:10.1021/jf801199q
 101. Mattison CP, Desormeaux WA, Wasserman RL, Yoshioka-Tarver M, Condon B, Grimm CC (2014) Decreased immunoglobulin E (IgE) binding to cashew allergens following sodium sulfite treatment and heating. *J Agric Food Chem* 62:6746–6755. doi:10.1021/jf501117p
 102. Untersmayr E, Jensen-Jarolim E (2008) The role of protein digestibility and antacids on food allergy outcomes. *J Allergy Clin Immunol* 121:1301–1308. doi:10.1016/j.jaci.2008.04.025
 103. Mattison CP, Grimm CC, Wasserman RL (2014) In vitro digestion of soluble cashew proteins and characterization of surviving IgE-reactive peptides. *Mol Nutr Food Res* 58:884–893. doi:10.1002/mnfr.201300299
 104. Crevel RWR, Baumert JL, Luccioli S (2014) Translating reference doses into allergen management practice: challenges for stakeholders. *Food Chem Toxicol* 67:277–287. doi:10.1016/j.fct.2014.01.033
 105. Crevel RWR, Ballmer-Weber BK, Holzhauser T (2008) Thresholds for food allergens and their value to different stakeholders. *Allergy* 63:597–609. doi:10.1111/j.1398-9995.2008.01636.x
 106. Luccioli S (2012) Food allergy guidelines and assessing allergic reaction risks: a regulatory perspective. *Curr Opin Allergy Clin Immunol* 12:323–330. doi:
 107. Costa J, Carrapatoso I, Oliveira MBPP, Mafra I (2014) Walnut allergens: molecular characterization, detection and clinical relevance. *Clin Exp Allergy* 44:319–341. doi:10.1111/cea.12267
 108. Costa J, Mafra I, Carrapatoso I, Oliveira M (in press) Hazelnut allergens: molecular characterisation, detection and clinical relevance. *Crit Rev Food Sci Nutr*. doi: 10.1080/10408398.2013.826173.
 109. Johnson PE, Sancho AI, Crevel RWR, Mills ENC (2011) Detection of allergens in foods. In: Nollet LML, van Hengel AJ (eds) *Food allergens: analysis instrumentation and methods*. CRC Press, Taylor & Francis Group, Boca Raton, pp 13–27
 110. Schubert-Ullrich P, Rudolf J, Ansari et al (2009) Commercialized rapid immunoanalytical tests for determination of allergenic food proteins: an overview. *Anal Bioanal Chem* 395:69–81. doi:10.1007/s00216-009-2715-y
 111. Wei Y, Sathe SK, Teuber SS, Roux KH (2002) A sensitive sandwich ELISA for the detection of trace amount of cashew nut in foods. *J Allergy Clin Immunol* 109(1 suppl 1):S303. doi:10.1016/S0091-6749(02)82063-2
 112. Wei YH, Sathe SK, Teuber SS, Roux KH (2003) A sensitive sandwich ELISA for the detection of trace amounts of cashew (*Anacardium occidentale* L.) nut in foods. *J Agric Food Chem* 51:3215–3221. doi:10.1021/jf025977x
 113. Gaskin FE, Niemann LM, Hefle SL, Taylor SL (2009) Validated enzyme-linked immunosorbent assay (ELISA) for detection of undeclared cashew nut residues in foods. *J Allergy Clin Immunol* 123(2, Supplement):S245. doi:10.1016/j.jaci.2008.12.946
 114. Gaskin FE, Taylor SL (2011) Sandwich enzyme-linked immunosorbent assay (ELISA) for detection of cashew nut in foods. *J Food Sci* 76:T218–T226. doi:10.1111/j.1750-3841.2011.02407.x
 115. Rejeb SB, Abbott M, Davies D, Cleroux C, Delahaut P (2005) Multi-allergen screening immunoassay for the detection of protein markers of peanut and four tree nuts in chocolate. *Food Addit Contam* 22:709–715. doi:10.1080/026520030500158450
 116. Cho CY, Nowatzke W, Oliver K, Garber EAE (2015) Multiplex detection of food allergens and gluten. *Anal Bioanal Chem* 407:4195–4206. doi:10.1007/s00216-015-8645-y
 117. Rebe Raz S, Liu H, Norde W, Bremer MGE (2010) Food allergens profiling with an imaging surface plasmon resonance-based biosensor. *Anal Chem* 82:8485–8491. doi:10.1021/ac101819g

118. Bignardi C, Elviri L, Penna A, Careri M, Mangia A (2010) Particle-packed column versus silica-based monolithic column for liquid chromatography-electrospray-linear ion trap-tandem mass spectrometry multiallergen trace analysis in foods. *J Chromatogr A* 1217:7579–7585. doi:10.1016/j.chroma.2010.10.037
119. Bignardi C, Mattarozzi M, Penna A et al (2013) A rapid size-exclusion solid-phase extraction step for enhanced sensitivity in multi-allergen determination in dark chocolate and biscuits by liquid chromatography-tandem mass spectrometry. *Food Anal Meth* 6:1144–1152. doi:10.1007/s12161-012-9521-4
120. Sealey-Voyksner J, Zweigenbaum J, Voyksner R (2016) Discovery of highly conserved unique peanut and tree nut peptides by LC-MS/MS for multi-allergen detection. *Food Chem* 194:201–211. doi:10.1016/j.foodchem.2015.07.043
121. Pilolli R, Monaci L, Visconti A (2013) Advances in biosensor development based on integrating nanotechnology and applied to food-allergen management. *Trac-Trends Anal Chem* 47:12–26. doi:10.1016/j.trac.2013.02.005
122. Prado M, Ortea I, Vial S, Rivas J, Calo-Mata P, Barros-Velázquez J (in press) Advanced DNA- and protein-based methods for the detection and investigation of food allergens. *Crit Rev Food Sci Nutr*. doi:10.1080/10408398.2013.873767
123. Alves RC, Barroso MF, González-García MB, Oliveira MBPP, Delerue-Matos MC (in press) New trends in food allergens detection: towards biosensing strategies. *Crit Rev Food Sci Nutr*. doi:10.1080/10408398.2013.831026
124. Cunsolo V, Muccilli V, Saletti R, Foti S (2014) Mass spectrometry in food proteomics: a tutorial. *J Mass Spectrom* 49:768–784. doi:10.1002/jms.3374
125. Fæste CK, Rønning HT, Christians U, Granum PE (2011) Liquid chromatography and mass spectrometry in food allergen detection. *J Food Prot* 74:316–345. doi:10.4315/0362-028X.JFP-10-336
126. Monaci L, Visconti A (2009) Mass spectrometry-based proteomics methods for analysis of food allergens. *Trac-Trend Anal Chem* 28:581–591. doi:10.1016/j.trac.2009.02.013
127. Picariello G, Mamone G, Addeo F, Ferranti P (2011) The frontiers of mass spectrometry-based techniques in food allergenomics. *J Chromatogr A* 1218:7386–7398. doi:10.1016/j.chroma.2011.06.033
128. Costa J, Mafra I, Carrapatoso I, Oliveira M (2012) Almond allergens: molecular characterization, detection, and clinical relevance. *J Agric Food Chem* 60:1337–1349. doi:10.1021/jf2044923
129. Brzezinski JL (2006) Detection of cashew nut DNA in spiked baked goods using a real-time polymerase chain reaction method. *J AOAC Int* 89:1035–1038
130. Piknova L, Kuchta T (2007) Detection of cashew nuts in food by real-time polymerase chain reaction. *J Food Nutr Res* 46:101–104
131. Ehlert A, Hupfer C, Demmel A, Engel KH, Busch U (2008) Detection of cashew nut in foods by a specific real-time PCR method. *Food Anal Meth* 1:136–143. doi:10.1007/s12161-008-9023-6
132. Lopez-Calleja IM, de la Cruz S, Gonzalez I, Garcia T, Martin R (2015) Development of real-time PCR assays to detect cashew (*Anacardium occidentale*) and macadamia (*Macadamia intergrifolia*) residues in market analysis of processed food products. *LWT-Food Sci Technol* 62:233–241. doi:10.1016/j.lwt.2015.01.023
133. Pafundo S, Gulli M, Marmioli N (2010) Multiplex real-time PCR using SYBR GreenER for the detection of DNA allergens in food. *Anal Bioanal Chem* 396:1831–1839. doi:10.1007/s00216-009-3419-z
134. Koppel R, van Velsen-Zimmerli F, Bucher T (2012) Two quantitative hexaplex real-time PCR systems for the detection and quantification of DNA from twelve allergens in food. *Eur Food Res Technol* 235:843–852. doi:10.1007/s00217-012-1806-8
135. Ehlert A, Demmel A, Hupfer C, Busch U, Engel KH (2009) Simultaneous detection of DNA from 10 food allergens by ligation-dependent probe amplification. *Food Addit Contam A* 26:409–418. doi:10.1080/02652030802593529
136. Wang W, Han JX, Wu YJ, Yuan F, Chen Y, Ge YQ (2011) Simultaneous detection of eight food allergens using optical thin-film biosensor chips. *J Agric Food Chem* 59:6889–6894. doi:10.1021/jf200933b