

Boundedness and Global Exponential Stability for Delayed Differential Equations with Applications

Teresa Faria^{a,**} and José J. Oliveira^{b,†}

^aDepartamento de Matemática and CMAF, Faculdade de Ciências,
Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal
e-mail: tfaria@ptmat.fc.ul.pt

^bDepartamento de Matemática, Escola de Ciências,
Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal
e-mail: jjoliveira@math.uminho.pt

Abstract

The boundedness of solutions for a class of n -dimensional differential equations with distributed delays is established, by assuming the existence of instantaneous negative feedbacks which dominate the delay effect. As an important by-product, some criteria for global exponential stability of equilibria are obtained. The results are illustrated with applications to delayed neural networks and population dynamics models.

Keywords: global exponential stability, global asymptotic stability, delay, neural network, Cohen-Grossberg neural network, delayed population model.

2000 AMS Subject Classification: 34K20, 34K25, 34K60, 92B20, 92D25.

1. Introduction

Let $C_n := C([-τ, 0]; \mathbb{R}^n)$ be the space of continuous functions from $[-τ, 0]$ to \mathbb{R}^n , $τ > 0$, equipped with the supremum norm $\|\varphi\| = \max_{-τ \leq \theta \leq 0} |\varphi(\theta)|$, for any fixed norm $|\cdot|$ in \mathbb{R}^n . As usual, $x_t \in C_n$ is defined by $x_t(\theta) = x(t + \theta)$, $-τ \leq \theta \leq 0$.

In this paper we are particularly interested in situations modelled by systems of functional differential equations (FDEs) of the form

$$\dot{x}_i(t) = -\rho_i(t, x_t)[b_i(x_i(t)) + f_i(t, x_t)], \quad t \geq 0, \quad i = 1, \dots, n, \quad (1.1)$$

* Corresponding author. Fax: +351 21 795 4288; Tel: +351 21 790 4894.

* Work partially supported by FEDER and Fundação para a Ciência e Tecnologia (FCT)-Plurianual 2007, under CMAF.

† Work partially supported by FCT, under grant SFRH/BD/29563/2006, CMAT, and programme POCI 2010.

where $b_i : \mathbb{R} \rightarrow \mathbb{R}$, $\rho_i, f_i : [0, \infty) \times C_n \rightarrow \mathbb{R}$ are continuous, with $\rho_i(t, \varphi)$ positive, $i = 1, \dots, n$. We further assume that the negative feedback condition $b_i(x)x > 0$ for $x \neq 0$, $i = 1, \dots, n$, is satisfied.

System (1.1) is sufficiently general to include as particular classes some well-known models used in population dynamics and neural networks, such as delayed Hopfield and Cohn-Grossberg models.

If (1.1) corresponds to an n -species population model, due to its biological interpretation only positive solutions are meaningful and therefore *admissible*. In a general setting, for an FDE in C_n ,

$$\dot{x}(t) = f(t, x_t), \quad t \geq t_0, \quad (1.2)$$

with $f : D \subset \mathbb{R} \times C_n \rightarrow \mathbb{R}^n$ continuous, we often fix a set $S \subset C_n$, with $[t_0, \infty) \times S \subset D$, as the set of *admissible initial conditions*. Naturally, a solution $x(t)$ with an *admissible* initial condition $x_{t_0} = \varphi \in S$ is said to be *admissible* if $x_t \in S$ for $t > t_0$ whenever x_t is defined. Under the assumption of uniqueness of solutions, the solution of (1.2) with initial condition $x_{t_0} = \varphi$, $\varphi \in S$, is denoted by $x(t, t_0, \varphi)$.

Assuming that (1.1) has an admissible equilibrium x^* , our main purpose is to establish sufficient conditions for its global exponential or asymptotic stabilities, whose definitions we recall here.

Definition 1.1. If $x^* \in S$ is an equilibrium of (1.1), x^* is said to be *globally exponentially stable* (in the set S of admissible solutions) if there are positive constants ε, M such that

$$|x(t, 0, \varphi) - x^*| \leq M e^{-\varepsilon t} \|\varphi - x^*\|, \quad \text{for all } t \geq 0, \varphi \in S;$$

and x^* is said to be *globally asymptotically stable* if it is stable and $x(t) \rightarrow x^*$ as $t \rightarrow \infty$, for all admissible solutions $x(t)$.

The present paper is inspired by a result of the authors in [7], which establishes a criterion for boundedness of solutions to (1.2). As shown here, such criterion can also be used to derive the global exponential stability of equilibria, and has applications to numerous relevant models, either recovering or improving results in recent literature normally obtained by construction of Lyapunov functionals. Although a very effective technique for proving boundedness of solutions and global attractivity of equilibria, the construction of a Lyapunov functional is in general a very difficult task, and involves long computations. Moreover, a new Lyapunov functional for each model under consideration is often required. On the contrary, our approach does not require the use of Lyapunov functional methods, and establishes simple criteria applicable to a quite general framework.

We now set some notation. For $c \in \mathbb{R}^n$, we use both c and \hat{c} to denote the constant function $\varphi(\theta) = c$ in C_n . A vector $d = (d_1, \dots, d_n) \in \mathbb{R}^n$ is said to be positive if $d_i > 0$ for $i = 1, \dots, n$, and we write $d > 0$.

Definition 1.2. For $d = (d_1, \dots, d_n) > 0$, define the inner product $\langle x, y \rangle_d = \sum_{i=1}^n d_i x_i y_i$ for $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{R}^n$, with associated norm $|x|_{2,d} = (\sum_{i=1}^n d_i x_i^2)^{1/2}$; define also the norm $|x|_{\infty,d} = \max_{i=1}^n d_i |x_i|$.

For \mathbb{R}^n with the norms $|\cdot|_{2,d}$, respectively $|\cdot|_{\infty,d}$ (where $d \in \mathbb{R}^n, d > 0$), we denote the corresponding norms in C_n by $\|\cdot\|_{2,d}$, respectively $\|\cdot\|_{\infty,d}$: $\|\varphi\|_{2,d} = \max_{-\tau \leq \theta \leq 0} |\varphi(\theta)|_{2,d}$ and $\|\varphi\|_{\infty,d} = \max_{-\tau \leq \theta \leq 0} |\varphi(\theta)|_{\infty,d}$. Clearly, all these norms are equivalent norms in C_n . If $d_i = 1, 1 \leq i \leq n$, we write simply $|\cdot|_2$ and $|\cdot|_\infty$ for the usual Euclidean and maximum norms in \mathbb{R}^n , respectively, and $\|\cdot\|_2$ and $\|\cdot\|_\infty$ for the corresponding norms in C_n .

If $D = [d_{ij}]$ is a square matrix with non-positive off-diagonal entries, i.e., $d_{ij} \leq 0$ for all $i \neq j$, we say that D is a *non-singular M-matrix* if all the eigenvalues of D have a positive real part. The latter is also equivalent to saying that there is a positive vector c such that $Nc > 0$ [8].

We should mention that in [7] the space \mathbb{R}^n is equipped with a norm $|\cdot|_{\infty,d}$, whereas here results are obtained for norms of both types $|\cdot|_{2,d}$ and $|\cdot|_{\infty,d}$. For related results for \mathbb{R}^n with $|\cdot|_2$, see [15, 20–22]. This raises the question of whether it is possible to obtain similar results for other norms (cf. [2, 4, 5, 10, 12, 20]), a problem to be addressed in future work.

We give a summary of the remainder of this paper. In Section 2, for each one of the norms $|\cdot|_{2,d}$ and $|\cdot|_{\infty,d}$, we present a criterion for boundedness of solutions to (1.2), which is then applied to the particular case of FDEs in the form (1.1). As shown in Section 3, as a by-product of such results we deduce sufficient conditions for the global exponential stability of an (admissible) equilibrium point x^* of (1.1), if it exists. In Section 4, we illustrate the results with well-known n -dimensional neural network models; although with less success, we also apply our ideas to FDEs from population dynamics. Finally, the last section presents some examples, namely a model where the advantage of working with both norms $|\cdot|_{2,d}$ and $|\cdot|_{\infty,d}$ is exploited.

2. Boundedness of solutions

Consider an FDE in \mathbb{R}^n

$$\dot{y}(t) = f(t, y_t), \quad t \geq t_0, \quad (2.1)$$

where $f : D \subset \mathbb{R} \times C_n \rightarrow \mathbb{R}^n$ is continuous. For the sake of simplicity of notation, uniqueness of solutions to (2.1) will be assumed. Fix a set $S \subset C_n$, with $[t_0, \infty) \times S \subset D$, as the set of *admissible initial conditions* for (2.1).

Consider first a norm $|\cdot|_{\infty,d}$ in \mathbb{R}^n , $d > 0$. Next result was established in [7, Lemma 3.2].

Proposition 2.1. [7] *Let $f : [t_0, \infty) \times S \rightarrow \mathbb{R}^n$ be continuous, $f = (f_1, \dots, f_n)$. For some $d = (d_1, \dots, d_n) > 0$, assume that f satisfies*

(H) $_\infty$ *for $t \geq t_0$ and $\varphi \in S$, if $|\varphi(\theta)|_{\infty,d} < |\varphi(0)|_{\infty,d}$ for $\theta \in [-\tau, 0)$, then $\varphi_i(0)f_i(t, \varphi) < 0$ for some $i \in \{1, \dots, n\}$ such that $|\varphi(0)|_{\infty,d} = d_i |\varphi_i(0)|$.*

Then, all admissible solutions of (2.1) are defined and bounded for $t \geq t_0$. Moreover, if $y(t) = y(t, t_0, \varphi)$ ($\varphi \in S$) is an admissible solution of (2.1), then $|y(t)|_{\infty, d} \leq \|\varphi\|_{\infty, d}$ for $t \geq t_0$.

The version of Proposition 2.1 with a norm $|\cdot|_{2, d}$ in \mathbb{R}^n is as follows:

Proposition 2.2. Suppose that $f : [t_0, \infty) \times S \rightarrow \mathbb{R}^n$ is continuous. For some $d = (d_1, \dots, d_n) > 0$, assume that f satisfies

(H)₂ for $t \geq t_0$ and $\varphi \in S$, if $|\varphi(\theta)|_{2, d} < |\varphi(0)|_{2, d}$ for $\theta \in [-\tau, 0)$, then $\langle \varphi(0), f(t, \varphi) \rangle_d < 0$.

Then, all admissible solutions of (2.1) are defined and bounded for $t \geq t_0$. Moreover, if $y(t) = y(t, t_0, \varphi)$ ($\varphi \in S$) is an admissible solution of (2.1), then $|y(t)|_{2, d} \leq \|\varphi\|_{2, d}$ for $t \geq t_0$.

Proof. Let $y(t)$ be an admissible solution of (2.1) on $[t_0 - \tau, a)$ for some $a > t_0$, with $|y(t)|_{2, d} \leq K$ for $t \in [t_0 - \tau, t_0]$. Suppose that there is $t_1 > t_0$ such that $|y(t_1)|_{2, d} > K$, and define

$$T = \min\{t \in [t_0, t_1] : \max_{s \in [t_0, t_1]} |y(s)|_{2, d} = |y(t)|_{2, d}\}.$$

We have $|y(T)|_{2, d} > K$ and $|y(t)|_{2, d} < |y(T)|_{2, d}$ for $t \in [t_0, T)$. Since $|y_T(\theta)|_{2, d} = |y(T + \theta)|_{2, d} < |y(T)|_{2, d}$ for $\theta \in [-\tau, 0)$, by (H)₂ we have $\langle y(T), f(t, y_T) \rangle_d < 0$ for all $t \geq t_0$. On the other hand, the definition of T implies that the derivative of the map $t \mapsto |y(t)|_{2, d}^2$ at $t = T$ is non-negative. Hence, $\langle y(T), f(T, y_T) \rangle_d \geq 0$, a contradiction. This proves that $y(t)$ is extensible to $[t_0 - \tau, \infty)$, with $|y(t)|_{2, d} \leq K$ for all $t > t_0$. ■

Remark 2.1. In particular, we note that Propositions 2.1 and 2.2 are valid if conditions (H)_∞ and (H)₂ are replaced, respectively, by

(h)_∞ for $t \geq t_0$ and $\varphi \in S$, if $\|\varphi\|_{\infty, d} = |\varphi(0)|_{\infty, d} > 0$, then $\varphi_i(0)f_i(t, \varphi) < 0$, for all $i \in \{1, \dots, n\}$ such that $|\varphi(0)|_{\infty, d} = d_i|\varphi_i(0)|$,

and

(h)₂ for $t \geq t_0$ and $\varphi \in S$, if $\|\varphi\|_{2, d} = |\varphi(0)|_{2, d} > 0$, then $\langle \varphi(0), f(t, \varphi) \rangle_d < 0$.

We are specially motivated by situations modelled by systems of FDEs of the form

$$\dot{x}_i(t) = -\rho_i(t, x_t)[b_i(x_i(t)) + f_i(t, x_t)], \quad t \geq 0, \quad i = 1, \dots, n, \quad (2.2)$$

where $\rho_i : [0, \infty) \times S \rightarrow (0, \infty)$, $b_i : \mathbb{R} \rightarrow \mathbb{R}$, $f_i : [0, \infty) \times S \rightarrow \mathbb{R}$ are continuous, for some set $S \subset C_n$ of admissible initial conditions.

Proposition 2.3. For (2.2), assume the following assumptions:

(A1) there is an equilibrium point $x^* \in S$ of (2.2);

(A2) there are constants $\beta_i > 0$ such that $\frac{b_i(u) - b_i(v)}{u - v} \geq \beta_i$ for all $u, v \in \mathbb{R}, u \neq v$ and $i = 1, \dots, n$;

(A3) $_{\infty}$ there is a vector $d = (d_1, \dots, d_n) > 0$ such that $|f_i(t, \varphi) - f_i(t, \psi)| \leq l_i \|\varphi - \psi\|_{\infty, d}$, for all $t \geq 0, \varphi, \psi \in S$, where the Lipschitz constants l_i satisfy $d_i l_i < \beta_i, i = 1, \dots, n$.

Then, all solutions $x(t) = x(t, t_0, \varphi)$ (with $t_0 \geq 0, \varphi \in S$) of (2.2) are defined and bounded on $[t_0, \infty)$ and satisfy $|x(t) - x^*|_{\infty, d} \leq \|\varphi - x^*\|_{\infty, d}$ for $t \geq t_0$.

Proof. Let $x^* = (x_1^*, \dots, x_n^*) \in S$ be such that $b_i(x_i^*) + f_i(t, x^*) = 0, t \geq 0, i = 1, \dots, n$. By the translation $\bar{x}(t) = x(t) - x^*$, (2.2) becomes

$$\dot{\bar{x}}_i(t) = -\rho_i(t, x^* + \bar{x}_t)[b_i(x_i^* + \bar{x}_i(t)) + f_i(t, x^* + \bar{x}_t)], \quad i = 1, \dots, n. \quad (2.3)$$

Clearly, (2.3) also has the form (2.2), for which (A1), (A2) and (A3) $_{\infty}$ hold with $S - x^*$ as the set of initial conditions. Hence, without loss of generality we may consider that the equilibrium of (2.2) is $x^* = 0$.

Take $t \geq 0$ and $\varphi \in S$ with $\|\varphi\|_{\infty, d} = |\varphi(0)|_{\infty, d} > 0$. Let i be such $\|\varphi\|_{\infty, d} = d_i |\varphi_i(0)|$, and suppose that $\varphi_i(0) > 0$ (the case $\varphi_i(0) < 0$ is analogous). Then,

$$b_i(\varphi_i(0)) + f_i(t, \varphi) = [b_i(\varphi_i(0)) - b_i(0)] + [f_i(t, \varphi) - f_i(t, 0)] \geq \beta_i \varphi_i(0) - l_i \|\varphi\|_{\infty, d} > 0.$$

This proves that (h) $_{\infty}$ holds, and the result follows now from Proposition 2.1. \blacksquare

Proposition 2.4. Consider (2.2), write $f = (f_1, \dots, f_n)$ and suppose that $\rho_i(t, \varphi)$ are uniformly positive and bounded, with

$$0 < r \leq \rho_i(t, \varphi) \leq R, \quad \text{for } t \geq 0, \varphi \in S, i = 1, \dots, n, \quad (2.4)$$

for some constants r, R . Assume (A1), (A2) and

(A3) $_2$ there is a vector $d = (d_1, \dots, d_n) > 0$ such that $|f(t, \varphi) - f(t, \psi)|_{2, d} \leq l \|\varphi - \psi\|_{2, d}$, for all $t \geq 0$ and $\varphi, \psi \in S$, where the Lipschitz constant l satisfies

$$l\sqrt{R/r} < \min_{1 \leq i \leq n} \beta_i. \quad (2.5)$$

Then, all solutions $x(t) = x(t, t_0, \varphi)$ (with $t_0 \geq 0, \varphi \in S$) of (2.2) are defined and bounded on $[t_0, \infty)$ and satisfy $|x(t) - x^*|_{2, d} \leq \|\varphi - x^*\|_{2, d}$ for $t \geq t_0$.

Proof. As in the above proof, we may already assume that the equilibrium of (2.2) is $x^* = 0$. Write $g = (g_1, \dots, g_n)$ with $g_i(t, \varphi) = \rho_i(t, \varphi)[b_i(\varphi_i(0)) + f_i(t, \varphi)]$ and let $\underline{\beta} = \min_{1 \leq i \leq n} \beta_i$.

Now, consider $t \geq 0$ and $\varphi \in S$ with $\|\varphi\|_{2, d} = |\varphi(0)|_{2, d} > 0$. Define $c_i = d_i \rho_i(t, \varphi), c = (c_1, \dots, c_n)$, and note that (2.4) implies that $r^{1/2}|x|_{2, d} \leq |x|_{2, c} \leq R^{1/2}|x|_{2, d}$ for $x \in \mathbb{R}^n$. We get

$$\begin{aligned} \langle \varphi(0), g(t, \varphi) \rangle_d &\geq \underline{\beta} \sum_{i=1}^n d_i \rho_i(t, \varphi) \varphi_i^2(0) - \sum_{i=1}^n d_i \rho_i(t, \varphi) |\varphi_i(0)| |f_i(t, \varphi) - f_i(t, 0)| \\ &= \underline{\beta} |\varphi(0)|_{2, c}^2 - \langle |\varphi(0)|, |f(t, \varphi) - f(t, 0)| \rangle_c \\ &\geq \underline{\beta} |\varphi(0)|_{2, c}^2 - |\varphi(0)|_{2, c} |f(t, \varphi) - f(t, 0)|_{2, c}, \end{aligned} \quad (2.6)$$

where we used the notation $|f| = (|f_1|, \dots, |f_n|)$, $|\varphi| = (|\varphi_1|, \dots, |\varphi_n|)$. On the other hand, from (2.4) and (A3)₂

$$|f(t, \varphi) - f(t, 0)|_{2,c} \leq \sqrt{R} |f(t, \varphi) - f(t, 0)|_{2,d} \leq l\sqrt{R} |\varphi(0)|_{2,d}.$$

From (2.5) and (2.6), we thus obtain

$$\langle \varphi(0), g(t, \varphi) \rangle_d \geq (\underline{\beta}\sqrt{r} - l\sqrt{R}) |\varphi(0)|_{2,d} |\varphi(0)|_{2,c} > 0.$$

The result is now a consequence of Proposition 2.2. ■

It is interesting to notice that for autonomous functions $f_i(t, \varphi) \equiv f_i(0, \varphi)$, or more generally when $f_i(t, \hat{x}) \equiv f_i(0, \hat{x})$ for $t \geq 0, x \in \mathbb{R}^n$, then (A2) and a Lipschitz condition on f imply that there is a unique equilibrium x^* of (2.2), as shown below. This means that, in this case, (A1) simply requires that the equilibrium x^* is in S .

Lemma 2.1. *Consider (2.2), where $f_i(t, \hat{x}) \equiv f_i(0, \hat{x}) := f_i(x)$ for $t \geq 0, x \in \mathbb{R}^n$, with $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$, $i = 1, \dots, n$. Assume (A2), and either (A3)_∞ or (A3)₂ with (2.5) replaced by $l < \min_{1 \leq i \leq n} \beta_i$. Then, there is a unique $x^* = (x_1^*, \dots, x_n^*) \in \mathbb{R}^n$ such that $b_i(x_i^*) + f_i(x^*) = 0$, $i = 1, \dots, n$.*

Proof. If (A2) and (A3)_∞ hold, the result was shown in [17] (for related results, see also e.g. [5, 6, 9, 16]), so here we only consider the situation for norms $|\cdot|_{2,d}$ in \mathbb{R}^n . Define $H : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $H(x) = b(x) + f(x)$, where

$$b(x) = (b_1(x_1), \dots, b_n(x_n)), \quad f(x) = (f_1(x), \dots, f_n(x)), \quad \text{for } x = (x_1, \dots, x_n).$$

If $x, y \in \mathbb{R}^n$ and $H(x) = H(y)$, then

$$\begin{aligned} \underline{\beta}^2 |x - y|_{2,d}^2 &\leq \sum_{i=1}^n d_i \beta_i^2 |x_i - y_i|^2 \leq \sum_{i=1}^n d_i |b_i(x_i) - b_i(y_i)|^2 \\ &= \sum_{i=1}^n d_i |f_i(x) - f_i(y)|^2 \leq l^2 |x - y|_{2,d}^2, \end{aligned}$$

with $l < \underline{\beta}$. This shows that H is injective. On the other hand, one writes $H(x) = (b(x) - b(0)) + (f(x) - f(0)) + (b(0) + f(0))$, from which one deduces that

$$|H(x)|_{2,d} \geq (\underline{\beta} - l) |x|_{2,d} - |b(0) + f(0)|_{2,d} \rightarrow \infty \quad \text{as } |x|_{2,d} \rightarrow \infty.$$

By [9], this proves that H is a homeomorphism of \mathbb{R}^n , hence there is a unique root x^* of the equation $H(x) = 0$. ■

3. Exponential stability

We now address the global exponential stability of x^* by using the results in Section 2.

Theorem 3.1. Consider the FDE (2.2), with $S \subset C_n$ as the set of admissible initial conditions. Assume (A1), (A2), (A3) $_{\infty}$ and

$$\underline{\rho}_i := \inf \{ \rho_i(t, \varphi) : t \geq t_0, \varphi \in S \} > 0, \quad i = 1, \dots, n. \quad (3.1)$$

Then, the equilibrium x^* is globally exponentially stable (in the set of all admissible solutions).

Proof. As before, by translating the equilibrium to the origin we may assume that $x^* = 0$, and thus $b_i(0) + f_i(t, 0) = 0$ for $t \geq 0$ and $i = 1, \dots, n$. By Proposition 2.3, we know already that solutions satisfy $|x(t, 0, \varphi)|_{\infty, d} \leq \|\varphi\|_{\infty, d}$ for $t \geq 0$ and $\varphi \in S$.

Now, effect the change of variables $z(t) = e^{\varepsilon t} x(t)$ for $\varepsilon > 0$. Eq. (2.2) translates as

$$\dot{z}_i(t) = g_i(t, z_t), \quad t \geq 0, i = 1, \dots, n, \quad (3.2)$$

where

$$g_i(t, \varphi) = \varepsilon \varphi_i(0) - \rho_i(t, e^{-\varepsilon(t+\cdot)} \varphi) e^{\varepsilon t} \left[b_i(e^{-\varepsilon t} \varphi_i(0)) + f_i(t, e^{-\varepsilon(t+\cdot)} \varphi) \right]. \quad (3.3)$$

Take $t \geq 0$ and $\varphi \in C_n$ with $e^{-\varepsilon(t+\cdot)} \varphi \in S$. Suppose that $\|\varphi\|_{\infty, d} = |\varphi(0)|_{\infty, d} > 0$, and let i be such $\|\varphi\|_{\infty, d} = d_i |\varphi_i(0)|_{\infty, d}$, with $\varphi_i(0) > 0$ (the case $\varphi_i(0) < 0$ is analogous). Then, for ε such that $\beta_i - d_i l_i e^{\varepsilon \tau} > 0$,

$$\begin{aligned} g_i(t, \varphi) &\leq \varepsilon \varphi_i(0) - \rho_i(t, e^{-\varepsilon(t+\cdot)} \varphi) e^{\varepsilon t} \varphi_i(0) \left[\beta_i e^{-\varepsilon t} - d_i l_i e^{-\varepsilon t} e^{\varepsilon \tau} \right] \\ &\leq \varphi_i(0) \left[\varepsilon - \underline{\rho}_i(\beta_i - d_i l_i e^{\varepsilon \tau}) \right]. \end{aligned}$$

Fix $\varepsilon > 0$ such that $\varepsilon - \underline{\rho}_i(\beta_i - d_i l_i e^{\varepsilon \tau}) < 0$. By applying Proposition 2.1, we deduce that the solutions $z(t)$ of (3.2) satisfy $|z(t)|_{\infty, d} \leq \|z_0\|_{\infty, d}$ for $t \geq 0$. Thus we obtain

$$|x(t, 0, \varphi)|_{\infty, d} = |e^{-\varepsilon t} z(t, 0, e^{\varepsilon \cdot} \varphi)|_{\infty, d} \leq e^{-\varepsilon t} \|\varphi\|_{\infty, d}, \quad t \geq 0, \varphi \in S. \quad \blacksquare$$

Similarly, the application of Proposition 2.2 yields the following result:

Theorem 3.2. For (2.2) with $S \subset C_n$ as the set of admissible initial conditions, assume (2.4), (A1), (A2) and (A3) $_2$. Then, the equilibrium x^* is globally exponentially stable.

Proof. The proof follows along the lines above, so it is omitted. \(\blacksquare\)

As we shall see in Section 4, condition (3.1) in general fails for (2.2) describing a population dynamics model, hence Theorems 3.1 and 3.2 are not applicable.

4. Applications

In this section, we apply the stability criteria in Section 3 to some n -dimensional FDEs which have been widely used in both neural networks and population dynamics.

4.1. Neural Network models with delays

As a particular class included in the framework set by (2.2), we consider generalizations of the Cohen-Grossberg neural network with distributed delays:

$$\dot{x}_i(t) = -k_i(x_i(t)) \left[b_i(x_i(t)) + \sum_{j=1}^n f_{ij}(x_{j,t}) \right], \quad t \geq 0, \quad i = 1, \dots, n, \quad (4.1)$$

where $k_i : \mathbb{R} \rightarrow (0, +\infty)$, $b_i : \mathbb{R} \rightarrow \mathbb{R}$ e $f_{ij} : C_1 \rightarrow \mathbb{R}$ are continuous, $i, j = 1, \dots, n$, and $x_t = (x_{1,t}, \dots, x_{n,t})$. Here, we take the entire phase space C_n as the set of admissible initial conditions. We note that (4.1) generalizes several systems studied in [1, 2, 4–6, 17, 20–25]; see also references therein.

For (4.1), denote $f = (f_1, \dots, f_n) : C_n \rightarrow \mathbb{R}^n$, where

$$f_i(\varphi) = \sum_{j=1}^n f_{ij}(\varphi_j), \quad \varphi = (\varphi_1, \dots, \varphi_n) \in C_n. \quad (4.2)$$

Proposition 4.1. *For (4.1), assume that b_i satisfy (A2), f_{ij} are Lipschitzian with*

$$|f_{ij}(\varphi) - f_{ij}(\psi)| \leq l_{ij} \|\varphi - \psi\|_\infty, \quad \varphi, \psi \in C_1, \quad i, j = 1, \dots, n, \quad (4.3)$$

and

$$\underline{k}_i := \inf_{x \in \mathbb{R}} k_i(x) > 0, \quad i = 1, \dots, n, \quad (4.4)$$

For β_i as in (A2), suppose in addition that the matrix

$$N = \text{diag}(\beta_1, \dots, \beta_n) - [l_{ij}]$$

is a non-singular M-matrix. Then, there is a unique equilibrium x^* of (4.1), which is globally exponentially stable.

Proof. Eq. (4.1) has the form (2.2) with $f_i(t, \varphi) = f_i(\varphi)$ as in (4.2) and $\rho_i(t, \varphi) = k_i(\varphi_i(0))$ for $t \geq 0, \varphi \in C_n$ and $i = 1, \dots, n$. If N is a non-singular M-matrix, then there is $c = (c_1, \dots, c_n) > 0$ such that $Nc > 0$ [8, Theorem 5.1], i.e.,

$$\beta_i > c_i^{-1} \sum_{j=1}^n l_{ij} c_j, \quad i = 1, \dots, n. \quad (4.5)$$

Clearly (A3) $_\infty$ holds with $d = (c_1^{-1}, \dots, c_n^{-1})$. The result is a consequence of Lemma 2.1 and Theorem 3.1. \blacksquare

We remark that system (4.1) is sufficiently general to include most of the autonomous delayed neural network models considered in the literature, namely Hopfield, Cohen-Grossberg, static, bidirectional associative memory models. Proposition 4.1 strongly improves several criteria established in e.g. [6, Theorem 2], [16, Theorems 2.1 and 2.2], [22, Theorem 3.1].

In an analogous way, we now apply Theorem 3.2 to (4.1) (for comparison of results, see e.g. [2, 3, 15, 21, 22]).

Proposition 4.2. For (4.1), assume (A2) and that there are constants $\underline{k}, \bar{k} > 0$ such that

$$\underline{k} \leq k_i(x) \leq \bar{k}, \quad \text{for } x \in \mathbb{R}, i = 1, \dots, n. \quad (4.6)$$

Suppose in addition that there is $d = (d_1, \dots, d_n) > 0$ such that

$$|f(\varphi) - f(\psi)|_{2,d} \leq l \|\varphi - \psi\|_{2,d}, \quad \varphi, \psi \in C_n, \quad (4.7)$$

with

$$l\sqrt{\bar{k}/\underline{k}} < \min_{1 \leq i \leq n} \beta_i, \quad (4.8)$$

for β_i as in (A2). Then, there is an equilibrium x^* of (4.1), which is globally exponentially stable.

Recently, the following Cohen-Grossberg model with time-dependent discrete delays has been studied [3, 12, 25]:

$$\dot{x}_i(t) = -k_i(x_i(t)) \left[b_i(x_i(t)) + \sum_{j=1}^n \sum_{p=1}^P h_{ijp}(x_j(t - \tau_{ijp}(t))) + J_i \right], \quad i = 1, \dots, n, t \geq 0, \quad (4.9)$$

where: $k_i : \mathbb{R} \rightarrow (0, +\infty)$, $b_i, h_{ijp} : \mathbb{R} \rightarrow \mathbb{R}$, $\tau_{ijp} : [0, +\infty) \rightarrow [0, +\infty)$ are continuous functions, with k_i uniformly positive on $[0, +\infty)$, $J_i \in \mathbb{R}$ and τ_{ijp} bounded, $i, j = 1, \dots, n$, $p = 1, \dots, P$.

For $\tau \geq 0$ such that $\tau_{ijp}(t) \in [0, \tau]$ for $t \geq 0$ and $i, j = 1, \dots, n$, $p = 1, \dots, P$, we take $C_n = C([- \tau, 0]; \mathbb{R}^n)$ as the phase space for (4.9) as well as the set of admissible initial conditions.

Proposition 4.3. For system (4.9), assume (A2), (4.4) and that h_{ijp} are Lipschitz continuous with Lipschitz constants l_{ijp} for $i, j = 1, \dots, n$, $p = 1, \dots, P$. For β_i as in (A2) and $l_{ij} = \sum_{p=1}^P l_{ijp}$, suppose also that the matrix $N = \text{diag}(\beta_1, \dots, \beta_n) - [l_{ij}]$ is a non-singular M-matrix. Then, there is a unique equilibrium x^* of (4.9), which is globally exponentially stable.

Proof. Write (4.9) as (2.2), where

$$f_i(t, \varphi) = \sum_{j=1}^n \sum_{p=1}^P h_{ijp}(\varphi_j(-\tau_{ijp}(t))) + J_i, \quad \rho_i(t, \varphi) = k_i(\varphi_i(0)), \quad i = 1, \dots, n,$$

for $t \geq 0, \varphi \in C_n$. Let $c = (c_1, \dots, c_n) > 0$ be such that (4.5) holds. For $\varphi, \psi \in C_n$ and $t \geq 0$ we have

$$|f_i(t, \varphi) - f_i(t, \psi)| \leq \sum_{j=1}^n \sum_{p=1}^P c_j l_{ijp} \|\varphi - \psi\|_{\infty, c^{-1}} = \left(\sum_{j=1}^n c_j l_{ij} \right) \|\varphi - \psi\|_{\infty, c^{-1}},$$

and therefore assumption (A3) $_{\infty}$ holds with $d = c^{-1} := (c_1^{-1}, \dots, c_n^{-1}) > 0$ and $l_i = \sum_{j=1}^n c_j l_{ij}$. Again, the result follows from Lemma 2.1 and Theorem 3.1. \blacksquare

The exponential stability for Eq. (4.9) was investigated in [12]. We note however that our Proposition 4.3 significantly improves the following criterion given in [12, Theorem 4]:

Proposition 4.4. [12] Consider Eq. (4.9) with $h_{ijp}(x) = c_{ijp}g_{ijp}(x)$, for $c_{ijp} \in \mathbb{R}$ and $g_{ijp}(x)$ continuous functions. Assume that, for $i, j = 1, \dots, n$ and $p = 1, \dots, P$:

(i) $k_i(x)$ are locally Lipschitz continuous and there are constants $\overline{k}_i, \underline{k}_i > 0$ such that

$$\underline{k}_i \leq k_i(x) \leq \overline{k}_i, \quad x \in \mathbb{R};$$

(ii) $b_i \in C^1(\mathbb{R}, \mathbb{R})$ with $b'_i(x) \geq \beta_i > 0$ for $x \in \mathbb{R}$;

(iii) g_{ijp} are bounded Lipschitzian functions, with Lipschitz constants μ_{ijp} ;

(iv) for $\overline{k}_i, \underline{k}_i$ as in (i) and β_i as in (ii), there are constants $\alpha_{ijp}, \gamma_{ijp} \in \mathbb{R}$, $\omega_i > 0$, $r > 1$ and $\sigma > 0$ such that

$$r\omega_i \underline{k}_i \beta_i - (r-1) \sum_{j=1}^n \sum_{p=1}^P \omega_j \overline{k}_i \mu_{ijp}^{\frac{r-\gamma_{ijp}}{r-1}} |c_{ijp}|^{\frac{r-\alpha_{ijp}}{r-1}} - \sum_{j=1}^n \sum_{p=1}^P \omega_j \overline{k}_i \mu_{ijp}^{\gamma_{ijp}} |c_{ijp}|^{\alpha_{ijp}} > \sigma. \quad (4.10)$$

Then, there is a unique equilibrium x^* of (4.9), which is globally exponentially stable.

Comparing the hypotheses in Proposition 4.3 and Proposition 4.4, we remark that, in the latter case, not only the constraints imposed on the functions k_i, b_i and g_{ijp} are clearly more restrictive, but also condition (4.10) is stronger than the requirement of N being a non-singular M-matrix, as shown in the next Lemma. See also Example 5.3.

Lemma 4.1. With the above notation, suppose that there are constants $r > 1$, $\sigma > 0$, $\alpha_{ijp}, \gamma_{ijp} \in \mathbb{R}$, $\omega_i > 0$, for $i, j = 1, \dots, n$ and $p = 1, \dots, P$, for which condition (4.10) is satisfied. Then, for $l_{ij} = \sum_{p=1}^P |c_{ijp}| \mu_{ijp}$, the matrix $N = \text{diag}(\beta_1, \dots, \beta_n) - [l_{ij}]$ is a non-singular M-matrix.

Proof. Condition (4.10) can be written as

$$r\omega_i \underline{k}_i \beta_i - \overline{k}_i \sum_{j=1}^n \omega_j \sum_{p=1}^P \left((r-1) \mu_{ijp}^{\frac{r-\gamma_{ijp}}{r-1}} |c_{ijp}|^{\frac{r-\alpha_{ijp}}{r-1}} + \mu_{ijp}^{\gamma_{ijp}} |c_{ijp}|^{\alpha_{ijp}} \right) > \sigma,$$

or, equivalently,

$$\omega_i \underline{k}_i \beta_i > \frac{\sigma}{r} + \overline{k}_i \sum_{j=1}^n \omega_j \sum_{p=1}^P \left[\frac{r-1}{r} \left(\mu_{ijp}^{\frac{1-\gamma_{ijp}}{r}} |c_{ijp}|^{1-\frac{\alpha_{ijp}}{r}} \right)^{\frac{r}{r-1}} + \frac{1}{r} \left(\mu_{ijp}^{\frac{\gamma_{ijp}}{r}} |c_{ijp}|^{\frac{\alpha_{ijp}}{r}} \right)^r \right].$$

Applying Young's inequality (see e.g. [12]), we therefore obtain

$$\omega_i \underline{k}_i \beta_i > \frac{\sigma}{r} + \overline{k}_i \sum_{j=1}^n \omega_j \sum_{p=1}^P \mu_{ijp} |c_{ijp}|. \quad (4.11)$$

Since $\overline{k}_i \geq \underline{k}_i > 0$, from (4.11) we have $\omega_i \beta_i > \sum_{j=1}^n \omega_j l_{ij}$, $i = 1, \dots, n$, from which we conclude that N is a non-singular M-matrix [8]. \blacksquare

4.2. Delayed Kolmogorov-type population models

Consider multiple species Kolmogorov systems with delays and instantaneous negative feedbacks given by

$$\dot{x}_i(t) = r_i(t)x_i(t)[-b_ix_i(t) + f_i(x_t)] \quad t \geq 0, \quad i = 1, \dots, n, \quad (4.12)$$

where $b_i > 0$, $r_i : [0, \infty) \rightarrow (0, \infty)$ are continuous and $f_i : C_n \rightarrow \mathbb{R}$ are Lipschitz continuous functions, $1 \leq i \leq n$. In a more general way, we can also study models with non-autonomous competition interactions $f_i(t, x_t)$, instead of $f_i(x_t)$, and suppose that $f_i(t, \varphi)$ are continuous and uniform Lipschitzian with respect to φ . For biological discussions and related results, see [7, 11, 13, 14], also for cited references.

In biological terms, only *positive* solutions of (4.12) are meaningful. We take

$$C_{\hat{0}} = \{\varphi = (\varphi_1, \dots, \varphi_n) \in C_n : \varphi_i(\theta) \geq 0 \text{ for } \theta \in [-\tau, 0), \varphi_i(0) > 0, i = 1, \dots, n\},$$

as the set of *admissible initial conditions*, and only consider solutions of (4.12) with initial conditions $x_{t_0} = \varphi \in C_{\hat{0}}$ for some $t_0 \geq 0$. A solution $x(t, t_0, \varphi)$ with initial condition in $C_{\hat{0}}$ is *admissible*, since $x_i(t, t_0, \varphi) = x_i(t_0) \exp(\int_{t_0}^t r_i(s)[-b_ix_i(s) + f_i(x_s)] ds) > 0$, for $t > t_0$ whenever it is defined.

If $x^* = (x_1^*, \dots, x_n^*)$ is a positive equilibrium of (4.1), for $y_i(t) = x_i(t) - x_i^*$ system (4.12) becomes

$$\dot{y}_i(t) = -r_i(t)(y_i(t) + x_i^*)[b_i y_i(t) - f_i(y_t + x^*) + f_i(x^*)], \quad i = 1, \dots, n, \quad (4.13)$$

with $C_{\hat{0}} - x^* := C_{-x^*}$ as the set of admissible initial conditions.

Proposition 4.5. *For (4.12), suppose that there is a positive equilibrium $x^* = (x_1^*, \dots, x_n^*)$, and that there is $d = (d_1, \dots, d_n) > 0$ such that for $\varphi, \psi \in C_{\hat{0}}$ and $i = 1, \dots, n$,*

$$|f_i(\varphi) - f_i(\psi)| \leq l_i \|\varphi - \psi\|_{\infty, d}, \quad \text{with } d_i l_i < b_i. \quad (4.14)$$

(i) If

$$\underline{r}_i := \inf_{t \geq 0} r_i(t) > 0, \quad i = 1, \dots, n, \quad (4.15)$$

then, for any $\delta > 0$ small, x^* is globally exponentially stable in the set of solutions satisfying $x_i(t) \geq \delta$ for $t \geq 0, i = 1, \dots, n$.

(ii) If (4.14) also holds with $\varphi, \psi \in \overline{C_{\hat{0}}}$ and

$$\sup_{t \geq 0} r_i(t) := \overline{r}_i < \infty \quad \text{and} \quad \int_0^\infty r_i(t) dt = \infty, \quad i = 1, \dots, n, \quad (4.16)$$

then x^* is globally asymptotically stable.

Proof. After translating x^* to the origin, consider (4.13). This equation has the form (2.2), for which condition $(A3)_\infty$ is satisfied by virtue of (4.14).

For the proof of (i), we now assume (4.15), and fix $\delta > 0$ with $\delta < x_i^*$, $i = 1, \dots, n$. Consider (4.13) with $S_\delta = \{(\varphi_1, \dots, \varphi_n) \in C_n : \varphi_i(\theta) + x_i^* \geq \delta \text{ for } \tau \leq \theta \leq 0, i = 1, \dots, n\}$ as the set of admissible initial conditions. We have $\rho_i(t, \varphi) := r_i(t)(\varphi_i(0) + x_i^*) \geq \delta \underline{r}_i$ for $t \geq 0, \varphi \in S_\delta, i = 1, \dots, n$. Thus, Theorem 3.1 implies the global exponential stability of zero on S_δ .

Now, suppose that (4.16) holds. From Proposition 2.3, all solutions of (4.13) are defined and bounded on $[0, \infty)$ and the origin is uniformly stable. It remains to prove that $y(t) \rightarrow 0$ as $t \rightarrow \infty$, for all (admissible) solutions of (4.13). We now argue as in the proof of [7, Theorem 3.4], where details can be found.

After the scaling $y_i(t) \mapsto d_i y_i(t), i = 1, \dots, n$, we may assume that (4.14) holds with $d = (1, \dots, 1)$. Let $y(t) = (y_i(t))_{i=1}^n$ be a solution to (4.13), and set

$$\begin{aligned} -v_i &= \liminf_{t \rightarrow \infty} y_i(t), & u_i &= \limsup_{t \rightarrow \infty} y_i(t), & i &= 1, \dots, n, \\ v &= \max_{1 \leq i \leq n} v_i, & u &= \max_{1 \leq i \leq n} u_i. \end{aligned}$$

Note that $-x_i^* \leq -v_i \leq u_i < \infty$, $i = 1, \dots, n$. It is sufficient to prove that $\max(u, v) = 0$. Assume e.g. that $v \leq u$, so that $\max(u, v) = u$ (the case $u \leq v$ is analogous). Let i be such that $u_i = u$, and fix $\varepsilon > 0$ small. Proceeding as in [7], we can show that there is a sequence $t_k \rightarrow \infty$ such that

$$y_i(t_k) \rightarrow u, \quad b_i y_i(t_k) - f_i(y_{t_k} + x^*) + f_i(x^*) \rightarrow 0 \quad \text{and} \quad \|y_{t_k}\|_\infty \leq u + \varepsilon. \quad (4.17)$$

The boundedness of $y(t)$ and $r_i(t)$ implies that $\dot{y}(t)$ is also bounded on $[0, \infty)$. Hence, the sequence (y_{t_k}) in C_n is uniformly bounded and equicontinuous. Up to a subsequence, we may suppose that $y_{t_k} \rightarrow \varphi$ in C_n , for some $\varphi \in \overline{C_{-x^*}}$. From (4.17), by letting $k \rightarrow \infty$ and $\varepsilon \rightarrow 0$ we obtain

$$\varphi_i(0) = u, \quad b_i u - f_i(\varphi + x^*) + f_i(x^*) = 0 \quad \text{and} \quad \|\varphi\|_\infty = u. \quad (4.18)$$

From (4.14) and (4.18), we deduce that $0 \geq b_i u - l_i \|\varphi\|_\infty = (b_i - l_i)u$. Since $l_i < b_i$ this implies that $u = 0$, showing that $y(t) \rightarrow 0$ as $t \rightarrow \infty$. The proof is complete. \blacksquare

If $f_i(\varphi) = \sum_{j=1}^n f_{ij}(\varphi_j)$ for $\varphi = (\varphi_1, \dots, \varphi_n)$ where $f_{ij} : C_1 \rightarrow \mathbb{R}$ satisfy the Lipschitz conditions in (4.3), then, as proven in Section 4.1, there is a positive vector d for which (4.14) is satisfied if and only if the matrix $N = \text{diag}(b_1, \dots, b_n) - [l_{ij}]$ is a non-singular M-matrix.

Remark 4.1. As a particular case of (4.12), consider Lotka-Volterra type systems

$$\dot{x}_i(t) = r_i(t)x_i(t) \left[\alpha_i - b_i x_i(t) - \sum_{j=1}^n l_{ij} \int_{-\tau}^0 x_j(t + \theta) d\eta_{ij}(\theta) \right], \quad i = 1, \dots, n, \quad (4.19)$$

where $\alpha_i \in \mathbb{R}, b_i > 0, l_{ij} \geq 0, \tau > 0, r_i(t)$ are positive continuous functions and $\eta_{ij} : [-\tau, 0] \rightarrow \mathbb{R}$ are normalized bounded variation functions. If $N = \text{diag}(b_1, \dots, b_n) - [l_{ij}]$ is a non-singular M-matrix and (4.16) holds, from Proposition 4.5 then the positive equilibrium x^* (if it exists) is globally

asymptotically stable. In [7], this stability was proven by imposing (4.16), that N is an M-matrix, and replacing $\det N \neq 0$ by the weaker condition $\det \tilde{M} \neq 0$, for $\tilde{M} = \text{diag}(b_1, \dots, b_n) + [\tilde{a}_{ij}]$, where $\tilde{a}_{ii} = a_{ii}$, $\tilde{a}_{ij} = -|a_{ij}|$ for $i \neq j$, and $a_{ij} = l_{ij} \int_{-\tau}^0 d\eta_{ij}(\theta)$. For more results, see also [11].

Remark 4.2. From Lemma 2.1, condition (4.14) implies that there is an equilibrium x^* of (4.12), but possibly non-positive. Note also that, for population models (4.12), the stability result in Proposition 4.5 is weaker than the criterion in Proposition 4.1 for neural networks, due to the fact that $\inf_{\varphi \in C_{-x^*}} (\varphi_i(0) + x_i^*) = 0$, so that Theorem 3.1 cannot be directly applied. On the other hand, our boundedness and stability results relative to norms $|\cdot|_{2,d}$ are not of interest in the present situation, since their application requires the estimates (2.4) and (2.5). We should however mention that Saito et al. [18] and Saito and Takeuchi [19] considered a predator-prey model (4.19) with $n = 2$ and the symmetry given by $N = \text{diag}(b, b) - L$, for $L := \begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix}$ and N defined as above, and showed its global asymptotic stability if and only if $\sqrt{\alpha^2 + \beta^2} \leq b$. Note that in this case we can choose $l = \|L\|_2 = \sqrt{\alpha^2 + \beta^2}$ for the Lipschitz constant l in (4.7).

5. Examples

We complement the above criteria with some illustrative examples.

Example 5.1. Consider the two neuron network with distributed delays

$$\begin{aligned} \dot{x}_1(t) &= -b_1 x_1(t) + a_{11} f_{11}(x_{1,t}) + a_{12} f_{12}(x_{2,t}) \\ \dot{x}_2(t) &= -b_2 x_2(t) + a_{21} f_{21}(x_{1,t}) + a_{22} f_{22}(x_{2,t}) \end{aligned} \quad (5.1)$$

where $b_i > 0$, $a_{ij} \in \mathbb{R}$ and $f_{ij} : C_1 := C([- \tau, 0]; \mathbb{R}) \rightarrow \mathbb{R}$ are Lipschitzian functions normalized so that $|f_{ij}(\varphi) - f_{ij}(\psi)| \leq \|\varphi - \psi\|_\infty$ for $\varphi, \psi \in C_1$, $i, j = 1, 2$.

Define $N = \text{diag}(b_1, b_2) - [|a_{ij}|]$. By [8, Theorem 5.1], N is a non-singular M-matrix if and only if all its principal minors are positive, which translates here as

$$b_1 > |a_{11}| \quad \text{and} \quad (b_1 - |a_{11}|)(b_2 - |a_{22}|) > |a_{12} a_{21}|. \quad (5.2)$$

By Proposition 4.1 and Lemma 2.1, (5.2) implies that there is a globally exponentially stable equilibrium x^* . That is the case if, for instance, $b_1 = b_2 = 2.1$, $a_{11} = a_{22} = 1$, $a_{12} = 1/2$, $a_{21} = 2$.

With b_i, f_{ij} as in (5.1), consider now the 2-species delayed population system

$$\begin{aligned} \dot{x}_1(t) &= x_1(t) \left[\alpha_1 - b_1 x_1(t) - a_{11} f_{11}(x_{1,t}) - a_{12} f_{12}(x_{2,t}) \right] \\ \dot{x}_2(t) &= x_2(t) \left[\alpha_2 - b_2 x_2(t) - a_{21} f_{21}(x_{1,t}) - a_{22} f_{22}(x_{2,t}) \right]. \end{aligned}$$

Assuming (5.2) and the existence of a positive equilibrium x^* , from Proposition 4.5 we conclude that x^* is globally asymptotically stable and that, for each $\delta > 0$ with $\delta < x_1^*, x_2^*$, x^* is globally exponentially stable in the set of solutions $x(t)$ such that $x_i(t) \geq \delta$, $i = 1, 2$, $t \geq 0$.

Example 5.2. Consider the 2-neuron network with one discrete delay

$$\begin{aligned}\dot{x}_1(t) &= -b_1x_1(t) + a_{11}f_1(x_1(t-\tau)) + a_{12}f_2(x_2(t-\tau)) \\ \dot{x}_2(t) &= -b_2x_2(t) + a_{21}f_1(x_1(t-\tau)) + a_{22}f_2(x_2(t-\tau)).\end{aligned}\tag{5.3}$$

Here, τ, b_i, a_{ij} are constants, $\tau, b_1, b_2 > 0$, $f_i : \mathbb{R} \rightarrow \mathbb{R}$ are Lipschitz continuous functions, normalized so that $|f_i(u) - f_i(v)| \leq |u - v|, u, v \in \mathbb{R}$, for $i, j = 1, 2$. The coefficients of the connection matrix $A := [a_{ij}]$ are arbitrary real numbers, representing either excitatory or inhibitory connections, according to their signs. We assume that there are self-connections, i.e., $a_{ii} \neq 0$. Without loss of generality we take $a_{11}, a_{22} > 0$.

We first suppose that $a_{12}a_{21} < 0$. Define

$$f(\varphi) = A \begin{pmatrix} f_1(\varphi_1(-\tau)) \\ f_2(\varphi_2(-\tau)) \end{pmatrix}, \quad \varphi = (\varphi_1, \varphi_2) \in C_2.\tag{5.4}$$

Choose $d = (d_1, d_2) > 0$ such that $d_1a_{11}a_{12} + d_2a_{21}a_{22} = 0$. Then for $\varphi, \psi \in C_2$ we have

$$\begin{aligned}\|f(\varphi) - f(\psi)\|_{2,d} &= \left[(d_1a_{11}^2 + d_2a_{21}^2) \left(f_1(\varphi_1(-\tau)) - f_1(\psi_1(-\tau)) \right)^2 \right. \\ &\quad \left. + (d_1a_{12}^2 + d_2a_{22}^2) \left(f_2(\varphi_2(-\tau)) - f_2(\psi_2(-\tau)) \right)^2 \right]^{1/2} \\ &= (\det A)^{1/2} \left[d_1 \frac{a_{11}}{a_{22}} \left(f_1(\varphi_1(-\tau)) - f_1(\psi_1(-\tau)) \right)^2 \right. \\ &\quad \left. + d_2 \frac{a_{22}}{a_{11}} \left(f_2(\varphi_2(-\tau)) - f_2(\psi_2(-\tau)) \right)^2 \right]^{1/2} \\ &\leq \left(\max \left\{ \frac{a_{11}}{a_{22}}, \frac{a_{22}}{a_{11}} \right\} \det A \right)^{1/2} \|\varphi - \psi\|_{2,d}.\end{aligned}$$

Hence, for $d > 0$ chosen as above, f given by (5.4) satisfies (4.8) if

$$\left(\max \left\{ \frac{a_{11}}{a_{22}}, \frac{a_{22}}{a_{11}} \right\} \det A \right)^{1/2} < \min\{b_1, b_2\}.\tag{5.5}$$

In this situation, Lemma 2.1 and Proposition 4.2 imply that there is an equilibrium point x^* of (5.3) which is globally exponentially stable.

Now let $a_{11}, a_{22} > 0$ as before, but consider both situations $a_{12}a_{21} < 0$ and $a_{12}a_{21} > 0$. Define $N = \text{diag}(b_1, b_2) - [a_{ij}]$. As in example 5.1, N is a non-singular M-matrix if

$$b_1 > a_{11} \quad \text{and} \quad (b_1 - a_{11})(b_2 - a_{22}) > |a_{12}a_{21}|,\tag{5.6}$$

in which case Proposition 4.1 allows to conclude the global exponential stability of x^* .

For instance, take system (5.3), with $b_1 = b_2 = 2.1, a_{11} = a_{22} = 1$ and $a_{12}a_{21} = -3$. In this case, (5.6) fails but (5.5) holds. Reversely, if $b_1 = 1.5, b_2 = 7.1, a_{11} = a_{22} = 1$ and $a_{12}a_{21} = -3$,

then (5.5) fails but (5.6) is satisfied. In both situations, we conclude that the equilibrium x^* , which exists by Lemma 2.1, is a global attractor of all solutions of (5.3), with an exponential rate of attraction. This shows the advantage of the possible use of both Euclidean and maximum norms (with weights d_i) in \mathbb{R}^n accounted for by our method.

Example 5.3. Consider the neural network model

$$\begin{aligned} \dot{x}_1(t) &= -(8 + 2 \sin x_1(t)) \left[7x_1(t) - \tanh x_1(t) - 2 \tanh x_2(t) - \tanh \left(x_1(t - \frac{1}{3} \sin t - 1) \right) \right. \\ &\quad \left. - \tanh \left(x_2(t - \frac{1}{4} e^{-\sin t} - 1) \right) + 2 \right] \\ \dot{x}_2(t) &= -(5 + 2 \cos x_2(t)) \left[10x_2(t) - 2 \tanh x_1(t) - \tanh x_2(t) - \tanh \left(x_1(t - \frac{1}{4} e^{-\sin t} - 1) \right) \right. \\ &\quad \left. - 2 \tanh \left(x_2(t - \frac{1}{3} \sin t - 1) \right) + 3 \right]. \end{aligned} \tag{5.7}$$

System (5.7) is very similar to the one considered in [12, Example 2]. It has the form (4.9), and satisfies all the assumptions in Proposition 4.3, with $\underline{k}_1 = 6, \underline{k}_2 = 3, \beta_1 = 7, \beta_2 = 10$ and $N = \text{diag}(7, 10) - \begin{pmatrix} 2 & 3 \\ 3 & 3 \end{pmatrix}$, clearly a non-singular M-matrix. From Proposition 4.3. the unique equilibrium of (5.7) is globally exponentially stable.

We now show that (5.7) does not satisfy condition (4.10), so that the criterion of Jiang et al. [12] given in Proposition 4.4 cannot be invoked. Besides the above coefficients, we have $\overline{k}_1 = 10, \overline{k}_2 = 7, \mu_{ijp} = 1, i, j, p = 1, 2, c_{111} = c_{112} = c_{121} = c_{221} = c_{212} = -1, c_{121} = c_{211} = c_{222} = -2$, thus conditions (i)-(iii) in the statement of Proposition 4.4 are satisfied; for the sake of contraction, suppose that condition (iv) there also holds. From (4.10), we derive (4.11), translated here as

$$\begin{cases} 22\omega_1 - 30\omega_2 > \sigma/r \\ -21\omega_1 + 9\omega_2 > \sigma/r \end{cases} \tag{5.8}$$

for some constants $\omega_1, \omega_2, \sigma > 0$ and $r > 1$. Since $A = \begin{pmatrix} 22 & -30 \\ -21 & 9 \end{pmatrix}$ is not a non-singular M-matrix, then it is not possible to find a positive vector $\omega = (\omega_1, \omega_2)$ such that $A\omega > 0$, which contradicts (5.8). This shows the advantage of the stability criterion given by Proposition 4.3 over the result established in [12].

References

- [1] C. Bai, Global exponential stability and existence of periodic solution of Cohen-Grossberg type neural networks with delays and impulses, *Nonlinear Anal. Real World Appl.* 9 (2008) 747–761.
- [2] J. Cao, New results concerning exponential stability and periodic solutions of delayed cellular neural networks, *Phys. Lett. A*, 307 (2003) 136–147.

- [3] J. Cao and J. Liang, Boundedness and stability for Cohen-Grossberg neural network with time-varying delays, *J. Math. Anal. Appl.* 296 (2004) 665–685.
- [4] J. Cao and Q. Song, Stability in Cohen-Grossberg-type bidirectional associative memory neural networks with time-varying delays, *Nonlinearity* 19 (2006) 1601–1617.
- [5] J. Cao and J. Wang, Global exponential stability and periodicity of recurrent neural networks with time delays, *IEEE Trans. Circuits Syst.* 52 (2005) 920–931.
- [6] Y. Chen, Global asymptotic stability of a general class of recurrent neural networks with time-varying delays, *IEEE Trans. Circuits Syst.* 52 (2005) 417–426.
- [7] T. Faria and J. Oliveira, Local and global stability for Lotka-Volterra systems with distributed delays and instantaneous feedbacks, *J. Differential Equations* 244 (2008) 1049–1079.
- [8] M. Fiedler, *Special Matrices and Their Applications in Numerical Mathematics*, Martinus Nijhoff Publ. (Kluwer), Dordrecht, 1986.
- [9] M. Forti and A. Tesi, New conditions for global stability of neural networks with applications to linear and quadratic programming problems, *IEEE Trans. Circuits Syst. I* 42 (1995) 354–366.
- [10] K. Gopalsamy, Stability of artificial neural networks with impulses, *Appl. Math. Comput.* 4 (2004) 783–813.
- [11] J. Hofbauer and J. So, Diagonal dominance and harmless off-diagonal delays, *Proc. Amer. Math. Soc.* 128 (2000) 2675–2682.
- [12] H. Jiang, J. Cao and Z. Teng, Dynamics of Cohen-Grossberg neural networks with time-varying delays, *Phys. Lett. A* 354 (2006) 414–422.
- [13] Y. Kuang, Global stability in delayed nonautonomous Lotka-Volterra type systems without saturated equilibrium, *Differential and Integral Equations* 9 (1996) 557–567.
- [14] Y. Kuang and H. L. Smith, Global stability for infinite delay Lotka-Volterra type systems, *J. Differential Equations* 103 (1993) 221–246.
- [15] X. Liao, C. Li and K. Wong, Criteria for exponential stability of Cohen-Grossberg neural networks, *Neural Networks* 17 (2004) 1401–1414.
- [16] S. Mohamad and K. Gopalsamy, Exponential stability of continuous-time and discrete-time cellular neural networks with delays, *Appl. Math. Comput.* 135 (2003) 17–38.
- [17] J. Oliveira, Global asymptotic stability for neural networks models, *preprint*.
- [18] Y. Saito, T. Hara and W. Ma, Necessary and sufficient conditions for permanence and global stability of a Lotka-Volterra system with two delays, *J. Math. Anal. Appl.* 236 (1999) 534–556.
- [19] Y. Saito and Y. Takeuchi, Sharp conditions for global stability of Lotka-Volterra systems with delayed intraspecific competitions, *Fields Inst. Commun.* 36 (2003) 195–211.
- [20] Q. Song and J. Cao, Global exponential stability of bidirectional associative memory neural networks with distributed delays, *J. Comput. Appl. Math.* 202 (2007) 266–279.
- [21] P. van den Driessche and X. Zou, Global stability in delayed Hopfield neural network models,

- SIAM J. Appl. Math.* 58 (1998) 1878–1890.
- [22] L. Wang and X. Zou, Harmless delays in Cohen-Grossberg neural networks, *Physica D* 170 (2002) 162–173.
- [23] L. Wang and X. Zou, Stability and bifurcation of bidirectional associative memory neural networks with delayed self-feedback, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **15** (2005), 2145–2159.
- [24] X. Yang, C. Li, X. Liao, D. Evans and G. Megson, Global exponential periodicity of a class of bidirectional associative memory networks with finite distributed delays, *Appl. Math. Comput.* 171 (2005) 108–121.
- [25] Z. Yang and D. Xu, Implulsive effects on stability of Cohen-Grossberg neural networks with variable delays, *Appl. Math. Comput.* 177 (2006) 63–78.