

Variational problems with non-constant gradient constraints

Lisa Santos
Centro de Matemática, University of Minho
Campus of Gualtar, 4700-030 Braga
Portugal

Abstract

This paper studies existence, uniqueness, continuous dependence on the given data and the asymptotic behavior of the solution of an evolutive variational inequality with non-constant gradient constraint and homogeneous Dirichlet boundary condition.

With assumptions on the given data, we prove existence of solution for a variational inequality with two obstacles, a Lagrange multiplier problem and an equation with gradient constraint. Equivalence of these problems with the variational inequality with gradient constraint is proved. An example of non-equivalence among these problems is given in order to show the necessity of the assumptions.

Mathematics subject classification: 35K85, 35K55, 35R35.

Key words: Parabolic variational inequality, parabolic PDE, free boundary problem.

1 Introduction

Variational problems with gradient constraint have been studied by several authors, in many different situations. The well known elastic-plastic torsion problem, a linear elliptic variational inequality, with constant coefficients and gradient constraint γ (the threshold of plasticity, which, for simplicity, is assumed here to be 1), in a simply connected domain, was solved by Brézis in [2]. Brézis also proved the equivalence of this problem with two other problems, a double obstacle variational inequality and a Lagrange multiplier problem. The first equivalence was generalized by Caffarelli and Friedman ([3]) to problems with non-homogeneous boundary conditions and the second one by Gerhardt ([6]) to multiply connected domains and also non-homogeneous boundary conditions. A general elliptic variational inequality with a convex set defined by a convex nonlinear function of the gradient, bounded from above by 1, was studied by Jensen in [9]. Evans studied general linear elliptic equations with a non-constant gradient constraint in [5] and his regularity result was extended by Ishii and Koike ([8]). Choe and Shim ([4]) obtained a regularity result for a variational inequality for the p -Laplacian, with non-constant gradient constraint and non-homogeneous boundary condition.

Parabolic variational inequalities with gradient constraint have also been considered (see, for instance [20], [21] and [23]).

Recently, the interest in problems with gradient constraint increased, since the critical state model of type-II superconductors in a longitudinal geometry turns out to be a nonlinear

evolution equation involving the p-Laplacian, for the relevant component of the magnetic field, together with a gradient constraint the threshold of which depends on the solution (see [1] and [17]). More explicitly, the model turns out to be a quasivariational inequality with a gradient constraint depending on the solution itself. Another model, the description of the growth of a sandpile, is also a quasi-variational inequality with gradient constraint depending on the solution ([16]). Some recent works about quasi-variational inequalities with gradient constraint are [10], [11] and [19].

In this paper we study an abstract evolutive variational inequality, with non-constant gradient constraint and homogeneous Dirichlet boundary condition. Part of the results presented here were announced in a symposium and are shortly described in [21].

This paper has two sections:

- the first section is divided in three subsections. The first one establishes briefly the existence of solution of the variational inequality, being this proof related with the one presented in [19], for the quasi-variational case with the p-Laplacian. Obviously, the result obtained here for the variational case (with the Laplacian) is stronger. The second subsection studies the continuous dependence of the solution on the data and the third one obtains the asymptotic limit of the solution, when $t \rightarrow +\infty$;

- in section two we suppose the given function depends only on the t variable. With assumptions on the gradient constraint we prove, in the first subsection, existence of solution of a double obstacle problem, deducing easily the $W_p^{2,1}(Q_T)$ regularity of the solution in this case. Afterwards, equivalence of this problem with the variational inequality with gradient constraint is proved. In the second subsection we establish existence of solution of a Lagrange multiplier problem and we prove that its solution is solution of the variational inequality. In the third subsection, existence for an equation with gradient constraint is proved, as well as the equivalence between this problem and the variational inequality. The fourth subsection is dedicated to the presentation of an example that shows the non-equivalence, in general, among these problems.

2 The variational problem

The main purpose of this section is to define the variational inequality problem and to present a brief proof of existence of solution. We also present a result about the continuous dependence of the solution on the given data and we study the asymptotic behavior of the solution, when $t \rightarrow +\infty$.

We assume that Ω is an open, bounded subset of \mathbb{R}^N , with a smooth boundary $\partial\Omega$. We denote by $I = [0, T]$ ($T \in \mathbb{R}^+$) a closed interval of \mathbb{R} and by Q_T the cylinder $\Omega \times]0, T[$. The set $\Sigma = \partial\Omega \times I$ is the lateral boundary and $\Omega_0 = \bar{\Omega} \times \{0\}$.

Let f and g be functions defined in Q_T , $g \geq 0$, and let h be defined in Ω .

Define, for a.e. $t \in I$, the following closed convex subset of $H_0^1(\Omega)$,

$$\mathcal{K}_{g(t)} = \{v \in H_0^1(\Omega) : |\nabla v(x)| \leq g(x, t) \text{ for a.e. in } x \in \Omega\}. \quad (1)$$

We consider the following variational inequality problem:

To find $u \in L^\infty(0, T; H_0^1(\Omega))$ such that

$$\left\{ \begin{array}{l} u(t) \in \mathbb{K}_{g(t)} \text{ for a.e. } t \in I, \quad u(0) = h, \\ \int_{\Omega} u_t(t)(v(t) - u(t)) + \int_{\Omega} \nabla u(t) \cdot \nabla (v(t) - u(t)) \geq \int_{\Omega} f(t)(v(t) - u(t)), \\ \forall v \in L^\infty(0, T; H_0^1(\Omega)) : v(t) \in \mathbb{K}_{g(t)}, \text{ for a.e. } t \in I. \end{array} \right. \quad (2)$$

2.1 Existence of solution

This subsection is dedicated to the proof of existence of solution, since the uniqueness is obvious. In [19], it can be found the proof of existence of solution for a related quasi-variational inequality problem with the p-Laplacian. There are many similarities between both proofs. Obviously, the solution for the variational inequality presented here is more regular and we prove here with more detail the facts which are specific of this problem.

Let us impose some assumptions on the given data:

$$\left\{ \begin{array}{l} g \in C^0(\overline{Q_T}) \cap W^{1,\infty}(0, T; L^\infty(\Omega)), \\ \exists m > 0 \forall (x, t) \in Q_T \quad g(x, t) \geq m, \\ f \in L^\infty(Q_T), \\ h \in H_0^1(\Omega), \quad |\nabla h| \leq g(0) \text{ a.e. in } \Omega. \end{array} \right. \quad (3)$$

Let $f^\varepsilon \in C_{\alpha, \alpha/2}^0(Q_T)$, $g_\varepsilon \in C_{\alpha, \alpha/2}^{1,0}(Q_T)$ and $h_\varepsilon \in C_\alpha^2(\overline{\Omega})$ ($0 < \alpha < 1$) be smooth approximations of f , g and h in the spaces $L^\infty(Q_T)$, $C^0(\overline{Q_T}) \cap W^{1,\infty}(0, T; L^\infty(\Omega))$, $H_0^1(\Omega)$ respectively, verifying h_ε and g_ε the additional conditions $|\nabla h_\varepsilon| \leq g_\varepsilon(0)$ a.e. in Ω , $g_\varepsilon \geq m$. Let k_ε be a C^2 , nondecreasing function, such that $k_\varepsilon(s) = 1$ if $s \leq 0$, $k_\varepsilon(s) = e^{s/\varepsilon}$ if $\varepsilon \leq s$. Consider now a family of approximate quasilinear parabolic problems, defined as follows,

$$\left\{ \begin{array}{l} u_t^\varepsilon - \nabla \cdot (k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) \nabla u^\varepsilon) = f^\varepsilon \text{ in } Q_T, \\ u^\varepsilon(0) = h_\varepsilon \text{ in } \Omega_0, \quad u^\varepsilon = 0 \text{ on } \Sigma. \end{array} \right. \quad (4)$$

The following theorem is the main result of this section.

Theorem 2.1 *With the assumption (3), problem (2) has a unique solution u belonging to $L^\infty(0, T; W_0^{1,\infty}(\Omega)) \cap C^0(\overline{Q_T}) \cap H^1(0, T; L^2(\Omega))$. Besides that, u is the weak limit in $L^p(0, T; W_0^{1,p}(\Omega))$ (for any $p \in]N, +\infty[$) of a subsequence $(u^{\varepsilon_n})_n$ of solutions of the family of approximate problems (4), $u^{\varepsilon_n} \rightarrow u$ in $C^0(\overline{Q_T})$, $u_t^{\varepsilon_n} \rightharpoonup u_t$ in $L^2(Q_T)$ -weak.*

We begin presenting first some auxiliary results.

Proposition 2.2 *Problem (4) has a unique solution $u^\varepsilon \in C_{\alpha, \alpha/2}^{2,1}(Q_T) \cap C^0(\overline{Q_T})$, $0 < \alpha < 1$.*

Proof: This result is an immediate consequence of the well known theory of quasilinear parabolic equations (see theorem 6.2, page 457 of [12]). ■

Lemma 2.3 *Let u^ε be the solution of problem (4) and suppose that the assumption (3) is verified. Then*

$$\exists C_0 > 0 \quad \forall \varepsilon \in]0, 1[\quad \forall (x, t) \in Q_T \quad |u^\varepsilon(x, t)| \leq C_0, \quad (5)$$

the constant C_0 being dependent on $\|f\|_{L^\infty(Q_T)}$ and $\|h\|_{L^\infty(\Omega)}$.

Proof: This result is an immediate consequence of the well known maximum principle for quasilinear parabolic equations (see [12], theorem 7.1, page 181). Notice that, since $h \in H_0^1(\Omega)$ and $|\nabla h| \leq g(0)$, then $h \in L^\infty(\Omega)$. ■

Lemma 2.4 *Let u^ε be the solution of problem (4) and suppose that the assumption (3) is verified. Then*

$$\exists C_1 > 0 \quad \forall \varepsilon \in]0, 1[\quad \|k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)\|_{L^1(Q_T)} \leq C_1, \quad (6)$$

the constant C_1 being dependent on $\frac{1}{m^2}$, $\|f\|_{L^2(Q_T)}^2$, $\|g\|_{L^2(Q_T)}^2$ and $\|h\|_{L^2(\Omega)}^2$.

Proof: Multiply the equation of the problem (4) by u^ε and integrate over $Q_t = \Omega \times]0, t[$. Then,

$$\frac{1}{2} \int_\Omega [u^\varepsilon(t)]^2 + \int_{Q_t} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)|\nabla u^\varepsilon|^2 = \int_{Q_t} f^\varepsilon u^\varepsilon + \frac{1}{2} \int_\Omega h_\varepsilon^2.$$

Using Hölder and Poincaré inequalities, denoting by C the Poincaré constant, we have

$$\int_{Q_t} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)|\nabla u^\varepsilon|^2 \leq C \left(\int_{Q_t} (f^\varepsilon)^2 \right)^{\frac{1}{2}} \left(\int_{Q_t} |\nabla u^\varepsilon|^2 \right)^{\frac{1}{2}} + \frac{1}{2} \int_\Omega h_\varepsilon^2$$

and using Young's inequality and the fact that $k_\varepsilon \geq 1$, we have

$$\int_{Q_T} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)|\nabla u^\varepsilon|^2 \leq C^2 \|f^\varepsilon\|_{L^2(Q_T)}^2 + \|h_\varepsilon\|_{L^2(\Omega)}^2.$$

Now,

$$\int_{Q_T} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)|\nabla u^\varepsilon|^2 = \int_{Q_T} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) [|\nabla u^\varepsilon|^2 - g_\varepsilon^2] + \int_{Q_T} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) g_\varepsilon^2$$

and, since $k_\varepsilon(s) = 1$ for $s \leq 0$ and $k_\varepsilon(s)s \geq 0$, $\forall s \in \mathbb{R}_0^+$, then

$$\begin{aligned} \int_{Q_t} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) [|\nabla u^\varepsilon|^2 - g_\varepsilon^2] &= \int_{\{|\nabla u^\varepsilon|^2 \leq g_\varepsilon^2\}} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) [|\nabla u^\varepsilon|^2 - g_\varepsilon^2] \\ &\quad + \int_{\{|\nabla u^\varepsilon|^2 > g_\varepsilon^2\}} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) [|\nabla u^\varepsilon|^2 - g_\varepsilon^2] \\ &\geq - \int_{Q_T} g_\varepsilon^2. \end{aligned}$$

Then we conclude that

$$\begin{aligned} \int_{Q_t} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) &\leq \frac{1}{m^2} \left[\int_{Q_T} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) |\nabla u^\varepsilon|^2 + \int_{Q_T} g_\varepsilon^2 \right] \\ &\leq \frac{1}{m^2} \left[C^2 \|f^\varepsilon\|_{L^2(Q_T)}^2 + \|h_\varepsilon\|_{L^2(\Omega)}^2 + \|g_\varepsilon\|_{L^2(Q_T)}^2 \right] \\ &\leq \frac{1}{m^2} \left[C^2 \|f\|_{L^2(Q_T)}^2 + \|h\|_{L^2(\Omega)}^2 + \|g\|_{L^2(Q_T)}^2 + 1 \right], \end{aligned}$$

since f^ε , g_ε and h_ε are approximations of f , g and h . ■

Lemma 2.5 *Let u^ε be the solution of problem (4) and suppose that the assumption (3) is verified. Then*

$$\exists C_2 > 0 \quad \forall \varepsilon \in]0, 1[\quad \|u_t^\varepsilon\|_{L^2(Q_T)}^2 \leq C_2, \quad (7)$$

the constant C_2 being dependent on C_1 and on $\|g\|_{W^{1,\infty}(0,T;L^\infty(\Omega))}^2$.

Proof: Multiply the equation of problem (4) by u_t^ε , noticing that $u_t^\varepsilon|_\Sigma \equiv 0$, and integrate over Q_t . Calling $\phi_\varepsilon(s) = \int_0^s k_\varepsilon(\tau) d\tau$, we have

$$\int_{Q_t} [u_t^\varepsilon]^2 + \frac{1}{2} \int_{Q_t} \frac{d}{dt} [\phi_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)] + \int_{Q_t} k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) g_\varepsilon g_{\varepsilon t} = \int_{Q_t} f^\varepsilon u_t^\varepsilon,$$

and, consequently,

$$\int_{Q_T} [u_t^\varepsilon]^2 \leq \int_{Q_T} [f^\varepsilon]^2 + 2C_1 \|g_\varepsilon\|_{L^\infty(Q_T)} \|g_{\varepsilon t}\|_{L^\infty(Q_T)} - \int_\Omega [\phi_\varepsilon(|\nabla u^\varepsilon(t)|^2 - g_\varepsilon^2(t))],$$

since

$$\int_\Omega [\phi_\varepsilon(|\nabla u^\varepsilon(0)|^2 - g_\varepsilon^2(0))] \leq 0, \quad \text{because } |\nabla u^\varepsilon(0)| = |\nabla h_\varepsilon| \leq g_\varepsilon(0).$$

Let $\Lambda = \{(x, t) \in Q_T : |\nabla u^\varepsilon(x, t)| < g_\varepsilon(x, t)\}$. Then we have:

$$\left\{ \begin{array}{l} \text{for a.e. } (x, t) \in \Lambda \quad \phi_\varepsilon(|\nabla u^\varepsilon(x, t)|^2 - g_\varepsilon^2(x, t)) = |\nabla u^\varepsilon(x, t)|^2 - g_\varepsilon^2(x, t) \geq -g_\varepsilon^2(x, t), \\ \text{for a.e. } (x, t) \in Q_T \setminus \Lambda \quad \phi_\varepsilon(|\nabla u^\varepsilon(x, t)|^2 - g_\varepsilon^2(x, t)) \geq 0 \geq -g_\varepsilon^2(x, t), \end{array} \right.$$

Consequently, for a.e. $t_0 \in I$,

$$-\int_{\Omega} \phi_{\varepsilon}(|\nabla u^{\varepsilon}(t_0)|^2 - g_{\varepsilon}^2(t_0)) \leq \|g_{\varepsilon}\|_{L^{\infty}(0,T;L^2(\Omega))}^2$$

and the proof is concluded. ■

Lemma 2.6 *Let u^{ε} be the solution of problem (4) and suppose that the assumption (3) is verified. Then*

$$\forall p \in [1, +\infty[\quad \exists D_p \in \mathbb{R}^+ \quad \forall \varepsilon \in]0, 1[: \quad \|\nabla u^{\varepsilon}\|_{L^p(Q_T)} \leq D_p, \quad (8)$$

the constant D_p being dependent on p , C_1 and on $\|g\|_{L^2(Q_T)}^2$.

Proof: We know, from (6), that there exists a constant C_1 , independent of ε , such that, for any $\varepsilon \in]0, 1[$,

$$\int_{Q_T} k_{\varepsilon}(|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2) \leq C_1.$$

So,

$$C_1 \geq \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} k_{\varepsilon}(|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2) = \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} e^{\frac{|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2}{\varepsilon}}.$$

Recalling that,

$$\forall s \in \mathbb{R}^+ \quad \forall j \in \mathbb{N} \quad e^s \geq \frac{s^j}{j!},$$

we obtain

$$\begin{aligned} \forall j \in \mathbb{N} \quad \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2\}} [|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2]^j \\ \leq j! \varepsilon^j \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} e^{\frac{|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2}{\varepsilon}} \leq j! \varepsilon^j C_1. \end{aligned}$$

Given $p \in [1, +\infty[$, we have

$$\int_{Q_T} |\nabla u^{\varepsilon}|^p = \int_{\{|\nabla u^{\varepsilon}|^2 \leq g_{\varepsilon}^2 + \varepsilon\}} |\nabla u^{\varepsilon}|^p + \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} |\nabla u^{\varepsilon}|^p. \quad (9)$$

Since there exists a constant $M > 0$, not depending on ε , such that $\|g_{\varepsilon}\|_{L^{\infty}(Q_T)} \leq M$, we can estimate, for $p \in \mathbb{N}$, the second integral in the second term of (9) as follows,

$$\begin{aligned} \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} |\nabla u^{\varepsilon}|^{2p} &\leq \\ &= \int_{\{|\nabla u^{\varepsilon}|^2 > g_{\varepsilon}^2 + \varepsilon\}} \sum_{j=0}^p \binom{p}{j} \|g_{\varepsilon}\|_{L^{\infty}(Q_T)}^{2p-2j} [|\nabla u^{\varepsilon}|^2 - g_{\varepsilon}^2]^j \\ &\leq \sum_{j=0}^p \binom{p}{j} \|g_{\varepsilon}\|_{L^{\infty}(Q_T)}^{2p-2j} j! \varepsilon^j C_1. \end{aligned}$$

The first integral in the second term of (9) is obviously bounded. In fact,

$$\int_{\{|\nabla u^\varepsilon|^2 \leq g_\varepsilon^2 + \varepsilon\}} |\nabla u^\varepsilon|^{2p} \leq \int_{Q_T} (g_\varepsilon^2 + 1)^p$$

and the conclusion follows easily, not only for $p \in \mathbb{N}$, but also for any $p \in [1, +\infty[$. ■

Lemma 2.7 *Define*

$$\mathbb{K}_{g_\varepsilon(t)} = \{v \in H_0^1(\Omega) : |\nabla v(x)| \leq g_\varepsilon(x, t) \text{ for a.e. } x \text{ in } \Omega\}. \quad (10)$$

Then, for any $v \in L^\infty(0, T; H_0^1(\Omega))$ such that $v(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in I$, there exists $v^\varepsilon \in L^\infty(0, T; H_0^1(\Omega))$ such that $v^\varepsilon(t) \in \mathbb{K}_{g_\varepsilon(t)}$ and

$$v^\varepsilon \longrightarrow v \quad \text{when } \varepsilon \rightarrow 0 \quad \text{in } L^\infty(0, T; H_0^1(\Omega)).$$

Proof: Let $\alpha_\varepsilon(t) = \|g_\varepsilon(t) - g(t)\|_{L^\infty(\Omega)}$. Obviously,

$$\alpha_\varepsilon \longrightarrow 0 \quad \text{when } \varepsilon \rightarrow 0 \quad \text{in } C^0([0, T]).$$

Define $\psi_\varepsilon(t) = 1 + \frac{\alpha_\varepsilon(t)}{m}$ and, given $v \in L^\infty(0, T; H_0^1(\Omega))$ such that $v(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in [0, T]$, define $v^\varepsilon = \frac{1}{\psi_\varepsilon} v \in L^\infty(0, T; H_0^1(\Omega))$.

Then,

$$|\nabla v^\varepsilon(x, t)| = \frac{1}{\psi_\varepsilon(x, t)} |\nabla v(x, t)| \leq \frac{1}{\psi_\varepsilon(x, t)} g(x, t) \leq g_\varepsilon(x, t),$$

because

$$\psi_\varepsilon(x, t) = 1 + \frac{\alpha_\varepsilon(t)}{m} \geq 1 + \frac{\alpha_\varepsilon(t)}{g_\varepsilon(x, t)} \geq \frac{g_\varepsilon(x, t) + g(x, t) - g_\varepsilon(x, t)}{g_\varepsilon(x, t)} = \frac{g(x, t)}{g_\varepsilon(x, t)}.$$

So, $v^\varepsilon(t) \in \mathbb{K}_{g_\varepsilon(t)}$ for a.e. $t \in [0, T]$ and $v^\varepsilon \longrightarrow v$ in $L^\infty(0, T; H_0^1(\Omega))$, when $\varepsilon \rightarrow 0$, since

$$\|v^\varepsilon(t) - v(t)\|_{H_0^1(\Omega)} = \left| \frac{1}{\psi_\varepsilon(t)} - 1 \right| \|v\|_{L^\infty(0, T; H_0^1(\Omega))} \leq \frac{|\alpha_\varepsilon(t)|}{m} \|v\|_{L^\infty(0, T; H_0^1(\Omega))},$$

and so,

$$v^\varepsilon \longrightarrow v \quad \text{when } \varepsilon \rightarrow 0 \text{ in } L^\infty(0, T; H_0^1(\Omega)). \quad \blacksquare$$

Proof of Theorem 2.1: We have proved that there are constants C_2 and C_p (independent of ε), $\forall p \in [1, +\infty[$, such that

$$\|u_t^\varepsilon\|_{L^2(Q_T)} \leq C_2, \quad \|u^\varepsilon\|_{L^p(0, T; W^{1, p}(\Omega))} \leq C_p.$$

So, for $p > N$, by a well known compactness theorem ([22], page 84), $\{u^\varepsilon\}_{\varepsilon \in]0, 1[}$ is relatively compact in $C(0, T; C(\overline{\Omega}))$ and so, at least for a subsequence, we have

$$u^\varepsilon(t) \longrightarrow u \quad \text{when } \varepsilon \rightarrow 0 \quad \text{uniformly in } t \text{ in } C^0(\overline{\Omega}),$$

and we also know that,

$$\begin{aligned} u^\varepsilon &\rightharpoonup u & \text{when } \varepsilon \rightarrow 0 & \text{ weakly in } L^p(0, T; W^{1,p}\Omega), \quad p \in [1, +\infty[\\ u_t^\varepsilon &\rightharpoonup u_t & \text{when } \varepsilon \rightarrow 0 & \text{ weakly in } L^2(Q_T). \end{aligned}$$

Let us prove now that u is, in fact, solution of the variational inequality (2). Given $v \in L^\infty(0, T; H_0^1(\Omega))$ such that $v(t) \in \mathcal{K}_{g(t)}$ for a.e. $t \in [0, T]$, let $v^\varepsilon \in L^\infty(0, T; H_0^1(\Omega))$ be defined as in Lemma 2.7. Multiply the equation of problem (4) by $v^\varepsilon(t) - u^\varepsilon(t)$ and use the monotonicity of k_ε and integration over $]s, t[\times \Omega$, $0 \leq s < t \leq T$ to conclude that

$$\int_s^t \int_\Omega u_t^\varepsilon (v^\varepsilon - u^\varepsilon) + \int_s^t \int_\Omega \nabla v^\varepsilon \cdot \nabla (v^\varepsilon - u^\varepsilon) \geq \int_s^t \int_\Omega f^\varepsilon (v^\varepsilon - u^\varepsilon).$$

Letting $\varepsilon \rightarrow 0$ and because s and t are arbitrary, we conclude that

$$\int_\Omega u_t(t)(v(t) - u(t)) + \int_\Omega \nabla v(t) \cdot \nabla (v(t) - u(t)) \geq \int_\Omega f(t)(v(t) - u(t)), \quad (11)$$

$$\forall v \in L^\infty(0, T; H_0^1(\Omega)) : \quad v(t) \in \mathcal{K}_{g(t)} \text{ for a.e. } t \in]0, T[.$$

Calling, for $M \geq \varepsilon$, $A_{M,\varepsilon} = \{(x, t) \in Q_T : |\nabla u^\varepsilon|^2 - g_\varepsilon^2 \geq M\}$, we see that, since in $A_{M,\varepsilon}$ we have $k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) \geq e^{\frac{M}{\varepsilon}}$,

$$|A_{M,\varepsilon}| = \int_{A_{M,\varepsilon}} 1 \leq \int_{A_{M,\varepsilon}} \frac{k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)}{e^{\frac{M}{\varepsilon}}} \leq C_1 e^{-\frac{M}{\varepsilon}},$$

since $\|k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)\|_{L^1(Q_T)} \leq C_1$, C_1 independent of ε . So, choosing $M = \sqrt{\varepsilon}$,

$$\begin{aligned} \int_{Q_T} (|\nabla u|^2 - g^2)^+ &\leq \liminf_{\varepsilon \rightarrow 0} \int_{Q_T} (|\nabla u^\varepsilon|^2 - g_\varepsilon^2 - \sqrt{\varepsilon})^+ \\ &= \liminf_{\varepsilon \rightarrow 0} \int_{A_{\sqrt{\varepsilon}, \varepsilon}} (|\nabla u^\varepsilon|^2 - g_\varepsilon^2 - \sqrt{\varepsilon}) \leq \lim_{\varepsilon \rightarrow 0} D |A_{\sqrt{\varepsilon}, \varepsilon}|^{\frac{1}{2}} = 0, \end{aligned}$$

where D is an upper bound of $\left[\int_{Q_T} (|\nabla u^\varepsilon|^2 - g_\varepsilon^2 - \sqrt{\varepsilon})^2 \right]^{\frac{1}{2}}$, D independent of ε . Consequently,

$$|\nabla u| \leq g \quad \text{a.e. in } Q_T.$$

So $u \in \mathcal{K}_{g(t)}$ and, to complete the proof, it is necessary to show that, by a variant of Minty's Lemma (see [18], lemma 4.2, page 99), it is possible to substitute the term $\int_\Omega \nabla v(t) \cdot \nabla (v(t) - u(t))$ in (11) by $\int_\Omega \nabla u(t) \cdot \nabla (v(t) - u(t))$, in order to obtain the variational inequality (2). So, let $w \in L^\infty(0, T; H_0^1(\Omega))$ be such that $w(t) \in \mathcal{K}_{g(t)}$ for a.e. $t \in [0, T]$. Define $v = u + \theta(w - u)$, $\theta \in]0, 1]$. Notice that $v(t) \in \mathcal{K}_{g(t)}$ for a.e. $t \in I$. Then, substituting v in (11) and dividing both sides by θ , we obtain

$$\int_\Omega u_t(t)(w(t) - u(t)) + \int_\Omega \nabla u(t) \cdot \nabla (w(t) - u(t)) + \theta \int_\Omega |\nabla (w(t) - u(t))|^2 \geq \int_\Omega f(t)(w(t) - u(t))$$

and, letting $\theta \rightarrow 0$, we prove that

$$\int_{\Omega} u_t(t)(w(t) - u(t)) + \int_{\Omega} \nabla u(t) \cdot \nabla (w(t) - u(t)) \geq \int_{\Omega} f(t)(w(t) - u(t)),$$

$$\forall w \in L^{\infty}(0, T; H_0^1(\Omega)) : w(t) \in \mathcal{K}_{g(t)} \text{ for a.e. } t \in]0, T[.$$

It remains to prove uniqueness, which is an immediate consequence of Theorem 2.8 below. ■

2.2 Continuous dependence on the data

This subsection is dedicated to the study of the continuous dependence of the solution of problem (2) on the given data.

Let u^1 and u^2 denote, respectively, the solution of problem (2) with data (f_1, g_1, h_1) and (f_2, g_2, h_2) . Denote

$$\mathcal{K}_{g_i(t)} = \{v \in H_0^1(\Omega) : |\nabla v| \leq g_i(t) \text{ a.e. in } \Omega\}, \quad i = 1, 2. \quad (12)$$

Theorem 2.8 *Suppose that (f_1, g_1, h_1) and (f_2, g_2, h_2) satisfy assumption (3), with the same m for g_1 and g_2 . Then*

$$\begin{aligned} \exists C_0, C_1, C_2 > 0 : \quad & \|u^1 - u^2\|_{L^{\infty}(0, T; L^2(\Omega))}^2 + \|\nabla(u^1 - u^2)\|_{L^2(Q_T)}^2 \leq \\ & C_0 \|f_1 - f_2\|_{L^2(Q_T)}^2 + C_1 \|h_1 - h_2\|_{L^2(\Omega)}^2 + \frac{C_2}{m} \|g_1 - g_2\|_{L^2(0, T; L^{\infty}(\Omega))}. \end{aligned} \quad (13)$$

Proof: Let $\theta(t) = \|g_1(t) - g_2(t)\|_{L^{\infty}(\Omega)}$ and $\psi(t) = 1 + \frac{\theta(t)}{m}$. Define $v^1(x, t) = \frac{1}{\psi(t)} u^1(x, t)$ and $v^2(x, t) = \frac{1}{\psi(t)} u^2(x, t)$.

Notice that

$$\frac{g_1(x, t)}{g_2(x, t)} = 1 + \frac{g_1(x, t) - g_2(x, t)}{g_2(x, t)} \leq \psi(t) \quad \text{and also} \quad \frac{g_2(x, t)}{g_1(x, t)} \leq \psi(t).$$

Since

$$|\nabla v^1(x, t)| = \left| \frac{1}{\psi(t)} \nabla u^1(x, t) \right| \leq \frac{1}{\psi(t)} g_1(x, t) \leq g_2(x, t)$$

and

$$|\nabla v^2(x, t)| = \left| \frac{1}{\psi(t)} \nabla u^2(x, t) \right| \leq \frac{1}{\psi(t)} g_2(x, t) \leq g_1(x, t),$$

we have that $v^1(t) \in \mathcal{K}_{g_2(t)}$ and $v^2(t) \in \mathcal{K}_{g_1(t)}$.

Putting in (2), for data (f_1, g_1, h_1) , $v = v^2$, we obtain

$$\begin{aligned} \int_{\Omega} u_t^1(t)(u^2(t) - u^1(t)) + \int_{\Omega} \nabla u^1(t) \cdot \nabla (u^2(t) - u^1(t)) &\geq \int_{\Omega} f_1(t)(u^2(t) - u^1(t)) \\ &+ \int_{\Omega} \left(1 - \frac{1}{\psi(t)}\right) \left[u_t^1(t)u^2(t) + \nabla u^1(t) \cdot \nabla u^2(t) - f_1(t)u^2(t) \right], \end{aligned} \quad (14)$$

and an analogous expression substituting the superscripts 1 by 2 and 2 by 1. Summing both inequalities, integrating between 0 and t and using Poincaré inequality (being C the Poincaré constant), we obtain

$$\begin{aligned} &\frac{1}{2} \int_{\Omega} (u^1(t) - u^2(t))^2 + \frac{1}{2} \int_{Q_T} |\nabla(u^1 - u^2)|^2 \\ &\leq \frac{C^2}{2} \int_{Q_T} (f_1 - f_2)^2 + \frac{1}{2} \int_{\Omega} (u^1(0) - u^2(0))^2 \\ &\quad + \int_{Q_T} \left| 1 - \frac{1}{\psi} \right| \left| u_t^1 u^2 + u_t^2 u^1 + 2\nabla u^1 \cdot \nabla u^2 - f_1 u^2 - f_2 u^1 \right| \\ &\leq \frac{C^2}{2} \int_{Q_T} (f_1 - f_2)^2 + \frac{1}{2} \int_{\Omega} (u^1(0) - u^2(0))^2 \\ &\quad + \left[\int_{Q_T} \left(1 - \frac{1}{\psi}\right)^2 \right]^{\frac{1}{2}} \left[\int_{Q_T} \left(u_t^1 u^2 + u_t^2 u^1 + 2\nabla u^1 \cdot \nabla u^2 - f_1 u^2 - f_2 u^1 \right)^2 \right]^{\frac{1}{2}}. \end{aligned}$$

Noticing that

$$\exists D > 0 \text{ (depending only on the data) :}$$

$$\|u_t^1 u^2 + u_t^2 u^1 + 2\nabla u^1 \cdot \nabla u^2 - f_1 u^2 - f_2 u^1\|_{L^2(Q_T)} \leq D,$$

since $u_t^1, u_t^2 \in L^2(Q_T)$, $u^1, u^2 \in L^\infty(0, T; W^{1,\infty}(\Omega))$, $f_1, f_2 \in L^\infty(Q_T)$ and that

$$\left[\int_{Q_T} \left(1 - \frac{1}{\psi}\right)^2 \right]^{\frac{1}{2}} \leq C \left(\int_{Q_T} \theta^2(t) \right)^{\frac{1}{2}},$$

(13) is proved. ■

2.3 Asymptotic behavior in time

In this subsection we are going to study the asymptotic limit, when $t \rightarrow +\infty$, of the solution of the variational inequality (2).

Considering $T = +\infty$, we begin proving that there exists a global solution of the variational inequality, defined in $\Omega \times \mathbb{R}_0^+$.

Lemma 2.9 *Suppose that the assumption (3) is verified, with $T = +\infty$. Then problem (2) has a solution u such that*

$$u \in L^\infty(0, +\infty; W_0^{1,\infty}(\Omega)) \cap C^0(\bar{\Omega} \times \mathbb{R}_0^+), \quad u_t \in L_{loc}^2(\mathbb{R}_0^+; L^2(\Omega)).$$

Proof: For each $T \in]0, +\infty[$, let $u_T : \Omega \times [0, T] \rightarrow \mathbb{R}$ denote the unique solution of the variational inequality (2). Let $u : \Omega \times \mathbb{R}_0^+ \rightarrow \mathbb{R}$ be defined as follows: given $(x, t) \in \Omega \times \mathbb{R}_0^+$, fix $T > t$ and define $u(x, t) = u_T(x, t)$. Clearly, u is well defined, due to the uniqueness (for each T) of solution of the variational inequality (2), and u solves (2) with $T = +\infty$. By the estimates obtained in the previous section, it is obvious that $u \in C^0(\bar{\Omega} \times \mathbb{R}_0^+)$ and that $u_t \in L_{loc}^2(\mathbb{R}_0^+; L^2(\Omega))$.

From the fact that $u(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in \mathbb{R}_0^+$, we obtain that

$$|\nabla u(x, t)| \leq g(x, t) \quad \text{for a.e. } (x, t) \in \Omega \times \mathbb{R}_0^+$$

and, since $g \in L^\infty(\Omega \times \mathbb{R}_0^+)$, it follows that $u \in L^\infty(0, +\infty; W^{1,\infty}(\Omega))$. ■

Let us now define, for given functions (f_∞, g_∞) , satisfying the assumption

$$\begin{cases} g_\infty \in C^0(\bar{\Omega}), \\ \exists m_\infty > 0 \forall x \in \Omega \quad g_\infty(x) \geq m_\infty, \\ f_\infty \in L^\infty(\Omega), \end{cases} \quad (15)$$

the limiting elliptic problem

$$\begin{cases} \text{To find } u^\infty \in \mathbb{K}_{g_\infty} : \\ \int_{\Omega} \nabla u^\infty \cdot \nabla (w - u^\infty) \geq \int_{\Omega} f_\infty (w - u^\infty), \quad \forall w \in \mathbb{K}_{g_\infty}, \end{cases} \quad (16)$$

where

$$\mathbb{K}_{g_\infty} = \{w \in H_0^1(\Omega) : |\nabla w| \leq g_\infty \text{ a.e. in } \Omega\}. \quad (17)$$

Existence of solution for the variational inequality (16) follows immediately from Stampacchia Theorem (see [18], Corollary 3.3 i), page 95).

Lemma 2.10 ([7], pg. 286) *Let $\zeta : \mathbb{R}_0^+ \rightarrow \mathbb{R}$ be a nonnegative function, absolutely continuous in any compact subinterval of \mathbb{R}_0^+ , $\Phi \in L_{loc}^1(0, +\infty)$ a nonnegative function and λ a positive constant such that*

$$\zeta'(t) + \lambda \zeta(t) \leq \Phi(t), \quad \forall t \in \mathbb{R}_0^+. \quad (18)$$

Then

$$\forall s, t \in \mathbb{R}_0^+ \quad \zeta(t + s) \leq e^{-\lambda t} + \frac{1}{1 - e^{-\lambda}} \left[\sup_{\tau \geq s} \int_{\tau}^{\tau+1} \Phi(\xi) d\xi \right]. \quad (19)$$

■

Theorem 2.11 *Suppose that (f, g, h) satisfy the assumption (3) with $T = +\infty$ and (f_∞, g_∞) satisfy the assumption (15). Suppose, in addition, that*

$$\int_t^{t+1} \int_\Omega [f(\tau) - f_\infty]^2 d\tau dx \longrightarrow 0, \text{ when } t \rightarrow +\infty, \quad (20)$$

$$\exists D > 0 \exists \gamma > \frac{1}{2} : \quad \|g(t) - g_\infty\|_{L^\infty(\Omega)} \leq \frac{D}{t^\gamma}.$$

Then we have

$$u(t) \longrightarrow u^\infty \text{ in } C^{0,\alpha}(\overline{\Omega}), \quad (0 < \alpha < 1) \quad \text{when } t \rightarrow +\infty. \quad (21)$$

Proof: Let, once more, $\theta(t) = \|g(t) - g_\infty\|_{L^\infty(\Omega)}$, $m_0 = \min\{m, m_\infty\}$ and $\psi(t) = 1 + \frac{\theta(t)}{m_0}$. Then we have that $\bar{v}(t) = \frac{1}{\psi(t)}u^\infty \in \mathbb{K}_{g(t)}$ and $v^\infty = \frac{1}{\psi(t)}u(t) \in \mathbb{K}_{g_\infty}$, for a.e. $t \in \mathbb{R}^+$. Substitute v by $\bar{v}(t)$ in (2) and w by v^∞ in (16). Then, we obtain,

$$\begin{aligned} & \int_\Omega u_t(t) \left(\frac{1}{\psi(t)}u^\infty - u(t) \right) + \int_\Omega \nabla u(t) \cdot \nabla \left(\frac{1}{\psi(t)}u^\infty - u(t) \right) \\ & \geq \int_\Omega f(t) \left(\frac{1}{\psi(t)}u^\infty - u(t) \right) \end{aligned}$$

and

$$\int_\Omega \nabla u^\infty \cdot \nabla \left(\frac{1}{\psi(t)}u(t) - u^\infty \right) \geq \int_\Omega f_\infty \left(\frac{1}{\psi(t)}u(t) - u^\infty \right)$$

and so,

$$\begin{aligned} & \int_\Omega (u(t) - u^\infty)_t (u(t) - u^\infty) + \int_\Omega |\nabla(u(t) - u^\infty)|^2 \leq \int_\Omega (f(t) - f_\infty)(u(t) - u^\infty) \\ & + \int_\Omega \left(1 - \frac{1}{\psi(t)} \right) \left[u_t(t)u^\infty + 2\nabla u(t) \cdot \nabla u^\infty - f(t)u^\infty - f_\infty u(t) \right]. \end{aligned}$$

Using Poincaré's inequality (denoting by C the Poincaré constant) and denoting $w(t) = u(t) - u^\infty$, we obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_\Omega w^2(t) + \frac{1}{2} \int_\Omega |\nabla w(t)|^2 \leq \frac{C^2}{2} \|f(t) - f_\infty\|_{L^2(\Omega)}^2 \\ & + C_1 \left| 1 - \frac{1}{\psi(t)} \right| \left[\|u_t(t)\|_{L^2(\Omega)} + 1 \right]. \end{aligned} \quad (22)$$

since $u^\infty, f_\infty \in L^\infty(\Omega)$, $u \in L^\infty(0, +\infty; W^{1,\infty}(Q_\infty))$, $u_t \in L^2_{loc}(\mathbb{R}^+; L^2(\Omega))$ and $f \in L^\infty(Q_\infty)$. Looking at the estimate of $\|u_t^\varepsilon\|_{L^2(\Omega \times [0, T])}^2$ presented in the proof of Lemma 2.5, we see that there exist constants D_0, D_1 , such that

$$\|u_t\|_{L^2(\Omega \times [0, t])}^2 \leq D_0 t + D_1,$$

where D_0 and D_1 are independent of t .

Calling $\zeta(t) = \|w(t)\|_{L^2(\Omega)}^2$ and

$$\Phi(t) = C^2 \|f(t) - f_\infty\|_{L^2(\Omega)}^2 + 2 \frac{C_1}{m} \|g(t) - g_\infty\|_{L^\infty(\Omega)} \left[\|u_t(t)\|_{L^2(\Omega)} + 1 \right]$$

we see that, in fact, $\Phi \in L^1_{loc}(0, +\infty)$ and ζ and Φ satisfy (18). So, for this choice of ζ and Φ , (19) is verified.

Applying Lemma 2.10, the result is proved, since

$$\begin{aligned} & \int_t^{t+1} \|g(\tau) - g_\infty\|_{L^\infty(\Omega)} \|u_t(\tau)\|_{L^2(\Omega)} d\tau \\ & \leq \|g(\tau) - g_\infty\|_{L^\infty(\Omega \times]t, t+1])} \left(\int_{\Omega \times]t, t+1[} |u_t|^2 \right)^{\frac{1}{2}} \\ & \leq \frac{D\tilde{D}_0}{t^{\gamma-\frac{1}{2}}} + \frac{D\tilde{D}_1}{t^\gamma} \longrightarrow 0 \quad \text{when } t \rightarrow +\infty, \end{aligned}$$

being \tilde{D}_0 and \tilde{D}_1 constants (independent of t). ■

3 Other problems with gradient constraint

There are problems with gradient constraint, well known in the literature, and which are related with this one.

We are going to define now three other problems related with the variational problem (2). It is our aim in this section to study whether this problem is equivalent to each one of the three problems defined here.

In this section we impose the additional assumptions on f and g :

$$\begin{aligned} f &= f(t) \quad (\text{i.e. } f \text{ is independent of } x), & f &\in L^\infty(0, T), \\ g &\in L^\infty(0, T; C^2(\bar{\Omega})) \cap W^{1,\infty}(0, T; L^\infty(\Omega)), & \partial\Omega &\text{ is of class } C^2. \end{aligned} \tag{23}$$

There are two main reasons for the choice of the function f depending only on t . The first one is a historical reason. In fact, the first problem with gradient constraint known in the literature is the elastic-plastic torsion problem, the elliptic variational inequality considered in (16), with gradient constraint $g_\infty \equiv 1$ and f a positive constant. For this first problem, with a clear physical meaning, equivalence with a double obstacle problem, with obstacles $\bar{\varphi}(x) = d(x, \partial\Omega)$ and $\underline{\varphi}(x) = -d(x, \partial\Omega)$, was proved by Caffarelli and Friedman in [3]. Equivalence of this specific problem with a Lagrange multiplier problem was also proved, for simply connected domains, by Brèzis in [2], and for multiply connected domains by Gerhardt, in [6]. Even in the case of the equation with gradient constraint, Evans ([5]) refers, without

proving, the truth of the equivalence, when the gradient constraint is one, of the equation with gradient constrain and the variational elastic-plastic torsion problem, alerting to the fact that this equivalence is not true, in the general case. The second and more important reason to consider f depending only on the time parameter is that any natural way to establish equivalence among all these problems depends on the application of the maximum principle and on the obtainance of very precise estimates on the gradient of the solutions of each of these problems. Considering f not constant in x would allow us to obtain gradient bounds but not the necessary ones to prove equivalence among the problems.

We consider first the *double obstacle problem*. To define it, let, for $x, z \in \overline{\Omega}$,

$$L_t(x, z) = \inf \left\{ \int_0^\delta g(\xi(s), t) ds : \delta > 0, \xi : [0, \delta] \rightarrow \Omega, \xi \text{ smooth}, \xi(0) = x, \xi(\delta) = z, \right. \\ \left. |\xi'| \leq 1 \right\},$$

$$\overline{\varphi}(x, t) = \bigvee \{w(x) : w \in \mathbb{K}_{g(t)}\}, \quad (24)$$

$$\underline{\varphi}(x, t) = \bigwedge \{w(x) : w \in \mathbb{K}_{g(t)}\}. \quad (25)$$

The function L_t is a metric and it can be shown (see [15], theorem 5.1, page 117) that

$$\overline{\varphi}(x, t) = \inf_{z \in \partial\Omega} \{L_t(x, z)\} = L_t(x, \partial\Omega)$$

and that

$$\underline{\varphi}(x, t) = \sup_{z \in \partial\Omega} \{-L_t(x, z)\} = -L_t(x, \partial\Omega).$$

In the special case where $g \equiv 1$, then L_t is the *geodesic distance* to $\partial\Omega$ and, if Ω is convex, L_t is the usual distance to $\partial\Omega$.

Consider the following closed convex set with two obstacles

$$\mathbb{K}(t) = \{w \in H_0^1(\Omega) : \underline{\varphi}(x, t) \leq w(x) \leq \overline{\varphi}(x, t) \text{ for a.e. } x \text{ in } \Omega\}. \quad (26)$$

The *double obstacle problem* is defined as follows:

To find $u \in L^\infty(0, T; H_0^1(\Omega))$ such that

$$\left\{ \begin{array}{l} u(t) \in \mathbb{K}(t) \text{ for a.e. } t \in I, \quad u(0) = h, \\ \int_\Omega u_t(t)(v(t) - u(t)) + \int_\Omega \nabla u(t) \cdot \nabla(v(t) - u(t)) \geq \int_\Omega f(t)(v(t) - u(t)), \\ \forall v \in L^\infty(0, T; H_0^1(\Omega)) : v(t) \in \mathbb{K}(t), \text{ for a.e. } t \in I. \end{array} \right. \quad (27)$$

It was shown by Caffarelli and Friedman in [3], that the elliptic formulation of the variational inequality with constant gradient constraint is equivalent to the double obstacle problem. The equivalence is still true for the parabolic case and constant gradient constraint (see

[20]). We intend to prove here the equivalence between these two problems, with non-constant gradient constraint g , as long as g satisfy suitable assumptions.

The following *Lagrange multiplier problem* is also related with the problem (2):

To find $(u, \lambda) \in [L^\infty(0, T; W_0^{1,\infty}(\Omega)) \cap W^{1,\infty}(0, T; L^2(\Omega))] \times L^\infty(Q_T)$ such that

$$\begin{cases} u_t - \nabla \cdot (\lambda \nabla u) = f \text{ in } Q_T, \\ \lambda \in k(|\nabla u|^2 - g^2), \\ u(0) = h \text{ in } \Omega_0, \quad u|_\Sigma = 0, \end{cases} \quad (28)$$

where k is the maximal monotone graph defined by $k(s) = 1$ if $s < 0$, $k(0) = [1, +\infty[$.

The existence of solution for the elliptic case was proved by Gerhardt ([6]) in the case where f is constant, the boundary condition is zero and $g \equiv 1$, as well as its equivalence with the elastic-plastic torsion problem. The parabolic case with non-homogeneous boundary condition was considered in [20].

Let us consider also the following *parabolic equation with gradient constraint*: to find $u \in W^{1,\infty}(0, T; L^\infty(\Omega)) \cap L^\infty(0, T; W^{1,\infty}(\Omega)) \cap W_{p,loc}^{2,1}(Q_T)$ such that

$$\begin{cases} \max\{u_t - \Delta u - f, |\nabla u| - g\} = 0 \text{ in } Q_T, \\ u(0) = h \text{ in } \Omega_0, \quad u|_\Sigma = 0. \end{cases} \quad (29)$$

It is easily seen that, when g is constant and $f = f(t)$, problems (2) and (29) are equivalent.

Zhu ([23]) has proved existence and (additional) regularity of solution for a similar problem, if $f \geq 0$ (depending on (x, t)). More precisely, he studied the problem

$$\begin{cases} \min\{u_t + Lu + f, -|\nabla u| + g\} = 0 \quad \text{in } \mathbb{R}^N \times \mathbb{R}_0^+, \\ u(x, T) = 0 \quad \forall x \in \mathbb{R}^N, \end{cases}$$

where L is an elliptic operator and T is a fixed instant.

3.1 Equivalence with the double obstacle problem

In this subsection, we present firstly a brief proof of existence of solution of problem (27). The equivalence between problem (2) and problem (27) is obtained when

$$\left(g^2\right)_t - \Delta \left(g^2\right) \geq 0. \quad (30)$$

Theorem 3.1 *With the assumption (3), problem (27) has a unique solution.*

In order to prove this theorem, we are going to present some auxiliary propositions first.

Proposition 3.2 *Let $\bar{\varphi}$ and $\underline{\varphi}$ be the obstacles defined, respectively, in (24) and in (25). We have*

$$\bar{\varphi}, \underline{\varphi} \in L^\infty(0, T; W^{1,\infty}(\Omega)) \cap W^{1,\infty}(0, T; L^\infty(\Omega)),$$

$$\text{for a.e. } (x, t) \in Q_T \quad |\nabla \bar{\varphi}(x, t)| = |\nabla \underline{\varphi}(x, t)| = g(x, t), \quad (31)$$

$$\text{for a.e. } t \in I \ \forall x_0 \in \partial\Omega \quad |\nabla \bar{\varphi}(x_0, t)| = |\nabla \underline{\varphi}(x_0, t)| = g(x_0, t), \quad (32)$$

$$\exists C > 0: \quad \Delta \bar{\varphi} \leq C, \quad \Delta \underline{\varphi} \geq -C, \quad \text{in } L^\infty(0, T; \mathcal{D}'(\Omega)). \quad (33)$$

Proof: Since $\bar{\varphi}(x, t) = L_t(x, \partial\Omega)$, L_t is continuous and $\partial\Omega$ is compact, there exists $z \in \partial\Omega$ such that $\bar{\varphi}(x, t) = L_t(x, z)$. So,

$$\forall \varepsilon > 0 \ \exists \delta_0^\varepsilon > 0 \ \exists \xi_\varepsilon : [0, \delta_0^\varepsilon] \rightarrow \Omega : \quad \xi_\varepsilon(0) = x, \ \xi_\varepsilon(\delta_0^\varepsilon) = z, \ |\xi'_\varepsilon| \leq 1,$$

$$\int_0^{\delta_0^\varepsilon} g(\xi_\varepsilon(s), t) ds - \varepsilon \leq \bar{\varphi}(x, t) \leq \int_0^{\delta_0^\varepsilon} g(\xi_\varepsilon(s), t) ds.$$

Since, obviously, given $h > 0$, we have $L_{t+h}(x, z) \leq \int_0^{\delta_0^\varepsilon} g(\xi_\varepsilon(s), t+h) ds$, then

$$\begin{aligned} \frac{L_{t+h}(x, z) - L_t(x, z)}{h} &\leq \int_0^{\delta_0^\varepsilon} \frac{g(\xi_\varepsilon(s), t+h) - g(\xi_\varepsilon(s), t)}{h} ds + \frac{\varepsilon}{h} \\ &= \int_0^{\delta_0^\varepsilon} [g_t(\xi_\varepsilon(s), \eta(h))] ds + \varepsilon/h \leq C, \end{aligned}$$

where $t < \eta(h) < t+h$ and C is a constant, if we choose, for instance, $\varepsilon = h^2$, noticing that $g_t \in L^\infty(Q_T)$ and that δ_0^ε is bounded from above independently of h (depending on $\|g\|_{L^\infty(Q_T)}$ and on Ω).

Analogously,

$$\forall h > 0 \ \forall \varepsilon > 0 \ \exists \delta_0^{\varepsilon, h} > 0 \ \exists \zeta_\varepsilon^h : [0, \delta_0^{\varepsilon, h}] \rightarrow \Omega : \quad \zeta_\varepsilon^h(0) = x, \ \zeta_\varepsilon^h(\delta_0^{\varepsilon, h}) = z, \ |(\zeta_\varepsilon^h)'| \leq 1,$$

$$\int_0^{\delta_0^{\varepsilon, h}} g(\zeta_\varepsilon^h(s), t+h) ds - \varepsilon \leq \bar{\varphi}(x, t+h) \leq \int_0^{\delta_0^{\varepsilon, h}} g(\zeta_\varepsilon^h(s), t+h) ds$$

and, of course, $L_t(x, z) \leq \int_0^{\delta_0^{\varepsilon, h}} g(\zeta_\varepsilon^h(s), t) ds$, so,

$$\begin{aligned} \frac{L_{t+h}(x, z) - L_t(x, z)}{h} &\geq \int_0^{\delta_0^{\varepsilon, h}} \frac{g(\zeta_\varepsilon^h(s), t+h) - g(\zeta_\varepsilon^h(s), t)}{h} ds - \frac{\varepsilon}{h} \\ &= \int_0^{\delta_0^{\varepsilon, h}} [g_t(\zeta_\varepsilon^h(s), \alpha(h))] ds - \varepsilon/h \geq C, \end{aligned}$$

being $t < \alpha(h) < t + h$ and, for a choice of $\varepsilon = h^2$, C is, as above, independent of h . Then $\bar{\varphi}_t \in L^\infty(Q_T)$ and so, $\bar{\varphi} \in W^{1,\infty}(0, T; L^\infty(\Omega))$.

By theorem 5.1, page 117 of [15], we know that, for a.e. $t \in I$, $\bar{\varphi}(t) \in W^{1,\infty}(\Omega)$ and that (31) is verified.

By theorem 8.2, page 179 of [15], we know that, if $\Omega_\delta = \{x \in \Omega : d(x, \partial\Omega) \geq \delta\}$ (where $\delta > 0$), then

$$\exists \delta_0 > 0 \quad \text{for a.e. } t \in I \quad \bar{\varphi}(t) \in C^2(\bar{\Omega} \setminus \Omega_{\delta_0}). \quad (34)$$

So, we have (32) and, using again theorem 5.1, page 117 of [15], we also have that

$$\forall \delta > 0 \quad \exists C_\delta > 0 \quad \text{for a.e. } t \in I : \quad \Delta \bar{\varphi}(t) \leq C_\delta \quad \text{in } \mathcal{D}'(\Omega_\delta). \quad (35)$$

So, (33) follows immediately for $\bar{\varphi}$ from (34) and (35). The proof for the function $\underline{\varphi}$ is analogous. ■

Consider the following family of penalized problems

$$\begin{cases} z_t^\varepsilon - \Delta z^\varepsilon + \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi}) = f & \text{in } Q_T, \\ z^\varepsilon(0) = h, \quad z|_\Sigma^\varepsilon = 0. \end{cases} \quad (36)$$

Proposition 3.3 *Problem (36) has a unique solution $z^\varepsilon \in W_p^{2,1}(Q_T)$, for any $p \in [1, +\infty[$ and*

- i) *the set $\left\{ \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi}) : \varepsilon \in]0, 1[\right\}$ is bounded in $L^\infty(Q_T)$;*
- ii) *the set $\{z^\varepsilon : \varepsilon \in]0, 1[\}$ is bounded in $W_p^{2,1}(Q_T)$, for any $p \in [1, +\infty[$.*

Proof: Let $w^\varepsilon = z^\varepsilon - h$. Then

$$\begin{cases} w_t^\varepsilon - \Delta w^\varepsilon + \frac{1}{\varepsilon} \left((w^\varepsilon + h) - ((w^\varepsilon + h) \wedge \bar{\varphi}) \vee \underline{\varphi} \right) = f - \Delta h & \text{in } Q_T, \\ w^\varepsilon = 0 & \text{on } \Sigma \cup \Omega_0. \end{cases}$$

This problem has a unique solution $w \in L^2(0, T; H_0^1(\Omega))$ (see [14], pg. 162) and so, problem (36) has a unique solution $z^\varepsilon \in L^2(0, T; H^1(\Omega))$.

Let $\bar{\psi} = \bar{\varphi} + M\varepsilon$, where M is a positive constant to be chosen later. By (33), we know that $\Delta \bar{\varphi}(x, t) \leq C$ in $L^\infty(0, T; \mathcal{D}'(\Omega))$ and we also know that $\bar{\varphi}_t \in L^\infty(Q_T)$. Then, $\bar{\psi}$ is a supersolution of problem (36), i.e. , if $L\xi = \xi_t - \Delta\xi + \frac{1}{\varepsilon} (\xi - (\xi \wedge \bar{\varphi}) \vee \underline{\varphi})$, then

$$L\bar{\psi} - f \geq 0 \quad \text{in } L^\infty(0, T; \mathcal{D}'(\Omega)), \quad (37)$$

as long as we impose $M \geq \|\bar{\varphi}_t\|_{L^\infty(Q_T)} + C + \|f\|_{L^\infty(0, T)}$. And, for $M \geq \|\underline{\varphi}_t\|_{L^\infty(Q_T)} + C + \|f\|_{L^\infty(0, T)}$, $\underline{\psi} = \underline{\varphi} - M\varepsilon$ is a subsolution of problem (36). So

$$(Lz^\varepsilon - L\bar{\psi}) \Phi^+ \leq 0 \quad \forall \Phi \in L^\infty(0, T; \mathcal{D}(\Omega))$$

and, consequently, as we can approximate $z^\varepsilon - \bar{\psi}$ by functions in $L^\infty(0, T; \mathcal{D}(\Omega))$, we have

$$(Lz^\varepsilon - L\bar{\psi})(z^\varepsilon - \bar{\psi})^+ \leq 0.$$

Easy calculations show that

$$\frac{1}{2} \int_{\Omega} [(\bar{\psi}(t) - z^\varepsilon(t))^+]^2 + \int_{Q_t} |\nabla(\bar{\psi} - z^\varepsilon)^+|^2 \leq 0$$

and, as a consequence, $z^\varepsilon \leq \bar{\psi}$ a.e. in Q_T .

Analogously we prove that $z^\varepsilon \geq \underline{\psi}$ a.e. in Q_T .

In particular, we conclude that

$$-M \leq \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi}) \leq M,$$

which proves i).

From the classical theory for parabolic equations (see [12], theorem 9.1, page 341), since $f - \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi})$ is bounded in $L^\infty(Q_T)$ independently of ε ,

$$\forall p \in [1, +\infty[\quad \exists C > 0, \quad C \text{ independent of } \varepsilon : \quad \|z^\varepsilon\|_{W_p^{2,1}(Q_T)} \leq C.$$

■

Proof of Theorem 3.1: By the preceding proposition, we know that $\{z^\varepsilon : \varepsilon \in]0, 1[\}$ is a bounded subset in $W_p^{2,1}(Q_T)$, for any $p \geq 1$. So, there exists a subsequence converging weakly to some function u^* , in this space. This convergence is strong in $L^2(0, T; H^1(\Omega))$ (see, for instance, [13], pg. 58). On the other hand, $z_t^\varepsilon \rightharpoonup u_t^*$ weakly in $L^2(Q_T)$.

Multiplying the first equation of problem (36) by $v - z^\varepsilon$, being $v(t)$ a function belonging to $\mathcal{K}(t)$, for a.e. $t \in I$, integrating over Q_T , and using the fact that

$$\int_{Q_T} \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi}) (v - z^\varepsilon) \leq 0,$$

we obtain

$$\int_{Q_T} z_t^\varepsilon (v - z^\varepsilon) + \int_{Q_T} \nabla z^\varepsilon \cdot \nabla (v - z^\varepsilon) \geq \int_{Q_T} f(v - z^\varepsilon),$$

$$\forall v : v(t) \in \mathcal{K}(t) \text{ for a.e. } t \in [0, T].$$

Letting $\varepsilon \rightarrow 0$, we see that

$$\int_{Q_T} u_t^* (v - u^*) + \int_{Q_T} \nabla u^* \cdot \nabla (v - u^*) \geq \int_{Q_T} f(v - u^*),$$

$$\forall v : v(t) \in \mathcal{K}(t) \text{ for a.e. } t \in [0, T].$$

Since

$$\underline{\varphi}(x, t) - M\varepsilon \leq z^\varepsilon(x, t) \leq \bar{\varphi}(x, t) + M\varepsilon, \quad \text{for a.e. } (x, t) \in Q_T,$$

letting $\varepsilon \rightarrow 0$, we obtain

$$\underline{\varphi}(x, t) \leq u^*(x, t) \leq \overline{\varphi}(x, t), \quad \text{for a.e. } (x, t) \in Q_T,$$

which means that $u^*(t) \in \mathbb{K}(t)$ for a.e. $t \in I$.

Given v such that $v(t) \in \mathbb{K}(t)$ for a.e. $t \in I$ and given $t_0 \in I$, $\delta > 0$ such that $I_\delta =]t_0 - \delta, t_0 + \delta[\subset I$, define

$$w(t) = \begin{cases} u^*(t) & \text{if } t \in I \setminus I_\delta, \\ v(t) & \text{if } t \in I_\delta. \end{cases}$$

Obviously, $w(t) \in \mathbb{K}(t)$ for a.e. $t \in I$ and so,

$$\int_{Q_T} u_t^*(w - u^*) + \int_{Q_T} \nabla u^* \cdot \nabla (w - u^*) \geq \int_{Q_T} f(w - u^*),$$

and, dividing the inequality by δ , we obtain

$$\frac{1}{\delta} \int_{t_0 - \delta}^{t_0 + \delta} \int_{\Omega} u_t^*(v - u^*) + \frac{1}{\delta} \int_{t_0 - \delta}^{t_0 + \delta} \int_{\Omega} \nabla u^* \cdot \nabla (v - u^*) \geq \frac{1}{\delta} \int_{t_0 - \delta}^{t_0 + \delta} \int_{\Omega} f(v - u^*),$$

and, letting $\delta \rightarrow 0$, where t_0 is a Lebesgue point, we have

$$\int_{\Omega} u_t^*(t_0)(v(t_0) - u^*(t_0)) + \int_{\Omega} \nabla u^*(t_0) \cdot \nabla (v(t_0) - u^*(t_0)) \geq \int_{\Omega} f(t_0)(v(t_0) - u^*(t_0)).$$

So, u^* is solution of problem (27).

The uniqueness of solution follows immediately from the fact that, if u_1 and u_2 are two solutions of problem (27) then, substituting $v = u_2(t)$ in the variational inequality when u_1 is considered as a solution and reciprocally and subtracting the inequalities obtained, one from the other, we get

$$\int_{\Omega} (u_1(t) - u_2(t))^2 + \int_{Q_t} |\nabla(u_1 - u_2)|^2 \leq 0$$

and so $u_1 = u_2$ a.e. in Q_T . ■

Proposition 3.4 *Suppose that the assumptions (3) and (23) are verified. Then*

$$\exists M > 0 \quad \forall (x, t) \in \Sigma \cup \Omega_0 \quad |\nabla z^\varepsilon(x, t)| \leq g(x, t) + M\sqrt{\varepsilon}. \quad (38)$$

Proof: Since $\partial\Omega$ is of class C^2 , it satisfies the *exterior sphere condition*, i.e.

$$\exists R > 0 \quad \forall x_0 \in \partial\Omega \quad \exists y_0 \in \mathbb{R}^N \quad D_R(y_0) \cap \overline{\Omega} = \{x_0\},$$

where $D_R(y_0) = \{x \in \mathbb{R}^N : d(x, y_0) \leq R\}$. Fixed $x_0 \in \partial\Omega$ we can, with a linear change of variables, suppose that $y_0 = 0$.

Let $\xi_\varepsilon(s) = e^{-\frac{s}{\sqrt{\varepsilon}}}$ and define

$$\overline{\psi}(x, t) = \overline{\varphi}(x, t) + M\varepsilon(1 - \xi_\varepsilon(\|x\| - R)), \quad \underline{\psi}(x, t) = \underline{\varphi}(x, t) - M\varepsilon(1 - \xi_\varepsilon(\|x\| - R)).$$

We are going to prove that $\bar{\psi}$ and $\underline{\psi}$ are, respectively, a supersolution and a subsolution of problem (36), in the same sense as in (37).

Notice that

$$\begin{aligned}\bar{\psi}|_{\Sigma} \geq 0 = z_{|\Sigma}^{\varepsilon} \geq \underline{\psi}|_{\Sigma}, \quad \bar{\psi}(x_0, t) = 0 = z^{\varepsilon}(x_0, t) = \underline{\psi}(x_0, t), \\ \bar{\psi}|_{\Omega_0} \geq \bar{\varphi}|_{\Omega_0} \geq h \geq \underline{\varphi}|_{\Omega_0} \geq \underline{\psi}|_{\Omega_0}.\end{aligned}$$

Let us denote $\xi_{\varepsilon}(\|x\| - R)$ simply by $\xi_{\varepsilon}(-)$. Easy calculation show that

$$\begin{aligned}\bar{\psi}_t = \bar{\varphi}_t, \quad \frac{1}{\varepsilon}(\bar{\psi} - (\bar{\psi} \wedge \bar{\varphi}) \vee \underline{\varphi}) = M(1 - \xi_{\varepsilon}(-)), \\ \Delta \bar{\psi} = \Delta \bar{\varphi} - M\xi_{\varepsilon}(-) + M\sqrt{\varepsilon}\xi_{\varepsilon}(-)\frac{n-1}{\|x\|}.\end{aligned}$$

Then,

$$\begin{aligned}\bar{\psi}_t - \Delta \bar{\psi} + \frac{1}{\varepsilon}(\bar{\psi} - (\bar{\psi} \wedge \bar{\varphi}) \vee \underline{\varphi}) \\ = \bar{\varphi}_t - \Delta \bar{\varphi} + M\xi_{\varepsilon}(-) - M\sqrt{\varepsilon}\xi_{\varepsilon}(-)\frac{n-1}{\|x\|} + M(1 - \xi_{\varepsilon}(-)) \\ \geq \bar{\varphi}_t - \Delta \bar{\varphi} + M\left(1 - \sqrt{\varepsilon}\frac{n-1}{R}\right).\end{aligned}$$

Choosing ε such that $\sqrt{\varepsilon}\frac{n-1}{R} \leq \frac{1}{2}$ and $M \geq 2\left(\|\bar{\varphi}_t\|_{L^{\infty}(Q_T)} + C + \|f\|_{L^{\infty}(0,T)}\right)$, we verify that $\bar{\psi}$ is a supersolution of problem (36). Analogously, for $M \geq 2\left(\|\underline{\varphi}_t\|_{L^{\infty}(Q_T)} + C + \|f\|_{L^{\infty}(0,T)}\right)$, $\underline{\psi}$ is a subsolution, so $\underline{\psi} \leq z^{\varepsilon} \leq \bar{\psi}$.

In particular, recalling that $z^{\varepsilon} \in W_p^{2,1}(Q_T)$, for any $p \in [1, +\infty[$ and the inclusion $W_p^{2,1}(Q_T) \hookrightarrow C_{\alpha}^{1,0}(\bar{\Omega} \times [0, T])$, if $p > n$ (and $\alpha = 1 - n/p$), then, $\nabla z^{\varepsilon}(x_0, t)$ exists for every $(x_0, t) \in \Sigma \cup \Omega_0$ and

$$|\nabla z^{\varepsilon}(x_0, t)| \leq \max\{|\nabla \bar{\psi}(x_0, t)|, |\nabla \underline{\psi}(x_0, t)|\}.$$

But,

$$|\nabla \bar{\psi}(x_0, t)| = \left| \nabla \bar{\varphi}(x_0, t) + M\sqrt{\varepsilon}\xi_{\varepsilon}(-)\frac{x_0}{\|x_0\|} \right| \leq g(x_0, t) + M\sqrt{\varepsilon},$$

and also $|\nabla \underline{\psi}(x_0, t)| \leq g(x_0, t) + M\sqrt{\varepsilon}$.

Besides that, $|\nabla z^{\varepsilon}(x, 0)| = |\nabla h(x)| \leq g(x, 0)$, which completes the proof. ■

Theorem 3.5 *Suppose that the assumptions (3), (23) and (30) are verified. Then problem (2) is equivalent to problem (27).*

Proof: Differentiate the first equation of problem (36) with respect to x_k , multiply by $z_{x_k}^{\varepsilon}$ and sum over k , denoting $v = |\nabla z^{\varepsilon}|^2$. Since $\frac{1}{2}v_t = z_{x_k}^{\varepsilon} z_{x_k t}^{\varepsilon}$ and $\frac{1}{2}\Delta v = (z_{x_i x_k}^{\varepsilon})^2 + z_{x_k}^{\varepsilon} \Delta z_{x_k}^{\varepsilon}$, we get then

$$\frac{1}{2}v_t - \frac{1}{2}\Delta v + \frac{1}{\varepsilon}(v - \nabla \tilde{z}^{\varepsilon} \cdot \nabla z^{\varepsilon}) \leq 0,$$

where $\tilde{z}^\varepsilon = (z^\varepsilon \wedge \bar{\varphi}) \vee \underline{\varphi}$ and using the Cauchy-Schwartz inequality,

$$\frac{1}{2}v_t - \frac{1}{2}\Delta v + \frac{1}{\varepsilon}(v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}}) \leq 0. \quad (39)$$

Since we have proved in Proposition 3.4 that $v(x, t) \leq (g(x, t) + M\sqrt{\varepsilon})^2$, for $(x, t) \in \Sigma \cup \Omega_0$, there exists N independent of ε such that $v(x, t) \leq g^2(x, t) + N\sqrt{\varepsilon}$ on $\Sigma \cup \Omega_0$. Then $(v - (g^2 + N\sqrt{\varepsilon}))^+$ is zero on $\Sigma \cup \Omega_0$.

Notice that the expression $\frac{2}{\varepsilon}(v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}})(v - (g^2 + N\sqrt{\varepsilon}))^+$ is always nonnegative. In fact,

$$|\nabla \tilde{z}^\varepsilon(x, t)| = \begin{cases} v(x, t) & \text{if } \underline{\varphi}(x, t) < z^\varepsilon(x, t) < \bar{\varphi}(x, t), \\ g(x, t) & \text{if } z^\varepsilon(x, t) \geq \bar{\varphi}(x, t) \text{ or } z^\varepsilon(x, t) \leq \underline{\varphi}(x, t). \end{cases}$$

Then, at a given point $(x, t) \in Q_T$,

$$v \leq g^2 + N\sqrt{\varepsilon} \implies \frac{2}{\varepsilon}(v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}})(v - (g^2 + N\sqrt{\varepsilon}))^+ = 0,$$

$$v > g^2 + N\sqrt{\varepsilon} \text{ and } \underline{\varphi} < z^\varepsilon < \bar{\varphi} \implies v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}} = 0 \implies \frac{2}{\varepsilon}(v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}})(v - (g^2 + N\sqrt{\varepsilon}))^+ = 0,$$

$$v > g^2 + N\sqrt{\varepsilon} \text{ and } z^\varepsilon \geq \bar{\varphi} \text{ or } z^\varepsilon \leq \underline{\varphi} \implies v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}} = v^{\frac{1}{2}}(v^{\frac{1}{2}} - g) > 0$$

$$\implies \frac{2}{\varepsilon}(v - |\nabla \tilde{z}^\varepsilon|v^{\frac{1}{2}})(v - (g^2 + N\sqrt{\varepsilon}))^+ > 0.$$

Multiplying the inequality (39) by $(v - (g^2 + N\sqrt{\varepsilon}))^+$ and integrating over $Q_t = \Omega \times [0, t]$, we have

$$\int_{Q_t} v_t(v - (g^2 + N\sqrt{\varepsilon}))^+ + \int_{Q_t} \nabla v \cdot \nabla(v - (g^2 + N\sqrt{\varepsilon}))^+ \leq 0$$

and so

$$\begin{aligned} & \frac{1}{2} \int_{\Omega} \left[(v(t) - (g^2(t) + N\sqrt{\varepsilon}))^+ \right]^2 - \frac{1}{2} \int_{\Omega} \left[(v(0) - (g^2(0) + N\sqrt{\varepsilon}))^+ \right]^2 \\ & + \int_{Q_t} \left| \nabla(v - (g^2 + N\sqrt{\varepsilon}))^+ \right|^2 \\ & \leq \int_{Q_t} - \left[(g^2 + N\sqrt{\varepsilon})_t - \Delta(g^2 + N\sqrt{\varepsilon}) \right] (v - (g^2 + N\sqrt{\varepsilon}))^+ \leq 0, \end{aligned}$$

using the assumption (30).

Since $(v(0) - (g^2(0) + N\sqrt{\varepsilon}))^+ \equiv 0$, we conclude that $(v - (g^2 + N\sqrt{\varepsilon}))^+ \equiv 0$, which means that

$$|\nabla z^\varepsilon|^2 \leq g^2 + N\sqrt{\varepsilon} \quad \text{a.e. in } Q_T,$$

and so, if u^* is the solution of problem (27), since u^* is the limit in $L^\infty(0, T; H_0^1(\Omega))$ of z^ε , when $\varepsilon \rightarrow 0$, we have $|\nabla u^*| \leq g$ a.e. in Q_T . In particular, $u^*(t) \in \mathcal{IK}_{g(t)}$. Since $\mathcal{IK}_{g(t)} \subseteq \mathcal{IK}(t)$, then u^* (by uniqueness) is also the solution of problem (2). \blacksquare

3.2 Equivalence with the Lagrange multiplier problem

We begin this subsection by proving existence of a solution for the Lagrange multiplier problem (28), when the following assumption is verified:

$$\left(g^2\right)_t \geq 0 \quad \Delta\left(g^2\right) \leq 0. \quad (40)$$

We would like to refer that the proof of existence of solution of this problem is very technical, even in the case where $g \equiv 1$. As we have done in the first section, the problem is approximated by a family of quasilinear parabolic problems (depending on a parameter ε) and the necessary estimates to pass to the limit are obtained. The more difficult estimates are the uniform boundedness of the gradient and the uniform (local) estimate in $L^2(0, T; H^2(\Omega))$. Although the procedure for both cases (g constant and non-constant) is the same, these two estimates are more difficult in the second case, since the partial derivatives of g are not zero.

Afterwards, with the same assumptions, we prove that if (u, λ) is a solution of (28) then u solves the variational inequality (2).

Consider the approximated problem (4), with $k_\varepsilon(s) = e^{\frac{Ns}{\varepsilon}}$ if $s \geq \varepsilon$, where N is a constant to be chosen later. In addition to the conditions imposed in the definition of the problem, we impose that k_ε is a $C^{2,1}$ function.

Recall that problem (4) has a solution $u^\varepsilon \in C_{\alpha, \alpha/2}^{2,1}(Q_T)$, $0 < \alpha < 1$.

Proposition 3.6 *Suppose that the assumptions (3) and (23) are verified. Then*

$$\exists C > 0 \quad |\nabla u^\varepsilon(x, t)|^2 \leq g_\varepsilon^2(x, t) + C\varepsilon \quad \text{for a.e. } (x, t) \in \Sigma \cup \Omega_0. \quad (41)$$

Proof: Let $\bar{\varphi}_\varepsilon$ and $\underline{\varphi}_\varepsilon$ be defined as in (24) and (25), respectively, with g replaced by $\sqrt{g_\varepsilon^2 + \varepsilon}$.

Let, for $s \in \mathbb{R}$, $\eta_\varepsilon(s) = s + \varepsilon(1 - e^{-Bs})$, where B is a positive constant, to be chosen later, depending on the given data and independent of ε .

We are going to prove that $\bar{\psi} = \eta_\varepsilon(\bar{\varphi}_\varepsilon)$ and $\underline{\psi} = \eta_\varepsilon(\underline{\varphi}_\varepsilon)$ are, respectively, a supersolution and a subsolution of problem (4).

Define $L\psi = \psi_t - \nabla \cdot (k_\varepsilon(|\nabla\psi|^2 - g_\varepsilon^2)\nabla\psi)$. Due to the monotonicity of k_ε , it is enough to prove that

$$L\bar{\psi} \geq f = Lu^\varepsilon \text{ in } Q_T, \quad \bar{\psi}|_{\Sigma \cup \Omega_0} \geq u^\varepsilon|_{\Sigma \cup \Omega_0}, \quad (42)$$

and

$$L\underline{\psi} \leq f = Lu^\varepsilon \text{ in } Q_T, \quad \underline{\psi}|_{\Sigma \cup \Omega_0} \leq u^\varepsilon|_{\Sigma \cup \Omega_0}. \quad (43)$$

We present here only the calculations for the supersolution, since the calculations for the subsolution are similar.

Obviously, since η_ε is an increasing function and $\eta_\varepsilon(0) = 0$, we have:

- for $(x, t) \in \Sigma$, $\bar{\psi}(x, t) = \eta_\varepsilon(\bar{\varphi}_\varepsilon(x, t)) = \eta_\varepsilon(0) = 0$;

- for $(x, t) \in \Omega_0$, $\bar{\psi}(x, 0) = \eta_\varepsilon(\bar{\varphi}_\varepsilon(x, 0)) \geq \bar{\varphi}_\varepsilon(x, 0) \geq h_\varepsilon(x) = u^\varepsilon(x, 0)$.

Easy calculations show that

$$\bar{\psi}_{x_i} = \eta'_\varepsilon(\bar{\varphi}^\varepsilon) \bar{\varphi}_{x_i}^\varepsilon, \quad |\nabla \bar{\psi}| = \eta'_\varepsilon(\bar{\varphi}^\varepsilon) \sqrt{g_\varepsilon^2 + \varepsilon},$$

$$\bar{\psi}_{x_i x_j} = \eta''_\varepsilon(\bar{\varphi}^\varepsilon) \bar{\varphi}_{x_i}^\varepsilon \bar{\varphi}_{x_j}^\varepsilon + \eta'_\varepsilon(\bar{\varphi}^\varepsilon) \bar{\varphi}_{x_i x_j}^\varepsilon,$$

$$\Delta \bar{\psi} = \eta''_\varepsilon(\bar{\varphi}^\varepsilon) (g_\varepsilon^2 + \varepsilon) + \eta'_\varepsilon(\bar{\varphi}^\varepsilon) \Delta \bar{\varphi}^\varepsilon,$$

$$\bar{\psi}_{x_i} \bar{\psi}_{x_j} \bar{\psi}_{x_i x_j} = (\eta'_\varepsilon(\bar{\varphi}^\varepsilon))^3 \bar{\varphi}_{x_i}^\varepsilon \bar{\varphi}_{x_j}^\varepsilon \bar{\varphi}_{x_i x_j}^\varepsilon + (\eta'_\varepsilon(\bar{\varphi}^\varepsilon))^2 \eta''_\varepsilon(\bar{\varphi}^\varepsilon) |\nabla \bar{\varphi}^\varepsilon|^4$$

and, noticing that $\bar{\varphi}_{x_i}^\varepsilon \bar{\varphi}_{x_i x_j}^\varepsilon = g_\varepsilon g_{\varepsilon x_j}$, then $\bar{\varphi}_{x_i}^\varepsilon \bar{\varphi}_{x_j}^\varepsilon \bar{\varphi}_{x_i x_j}^\varepsilon = g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon$.

Denoting $\xi(s) = e^{-Bs}$, we have

$$\eta'_\varepsilon(s) = 1 + \varepsilon B \xi(s), \quad \eta''_\varepsilon(s) = -\varepsilon B^2 \xi(s).$$

Calculate now $L\bar{\psi}$ (to simplify, we will omit the argument $\bar{\varphi}^\varepsilon$ in η'_ε and in ξ):

$$\begin{aligned} L\bar{\psi} &= \bar{\psi}_t - k'_\varepsilon (|\nabla \bar{\psi}|^2 - g_\varepsilon^2) (2\bar{\psi}_{x_i} \bar{\psi}_{x_j} \bar{\psi}_{x_i x_j} - 2g_\varepsilon g_{\varepsilon x_i} \bar{\psi}_{x_i}) - k_\varepsilon (|\nabla \bar{\psi}|^2 - g_\varepsilon^2) \Delta \bar{\psi} \\ &= \eta'_\varepsilon \bar{\varphi}_t^\varepsilon + 2k'_\varepsilon \left((\eta'_\varepsilon)^2 (g_\varepsilon^2 + \varepsilon) - g_\varepsilon^2 \right) \left\{ \left[-(\eta'_\varepsilon)^2 \eta''_\varepsilon (g_\varepsilon^2 + \varepsilon)^2 - (\eta'_\varepsilon)^3 g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \right] + \right. \\ &\quad \left. \eta'_\varepsilon g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \right\} + k_\varepsilon \left((\eta'_\varepsilon)^2 (g_\varepsilon^2 + \varepsilon) - g_\varepsilon^2 \right) \left[-\eta'_\varepsilon \Delta \bar{\varphi}^\varepsilon - \eta''_\varepsilon (g_\varepsilon^2 + \varepsilon) \right]. \end{aligned}$$

Notice that:

- for $s \geq \varepsilon$ we have $k'_\varepsilon(s) = \frac{N}{\varepsilon} k_\varepsilon(s)$ and $(\eta'_\varepsilon)^2 (g_\varepsilon^2 + \varepsilon) - g_\varepsilon^2 \geq \varepsilon$;
- $\exists C_0 > 0$ (depending only on $\|\bar{\varphi}\|_\infty$) such that $1 \geq \xi = \xi(\bar{\varphi}_\varepsilon) \geq e^{-BC_0}$;
- $k_\varepsilon \geq 1$, $g_\varepsilon^2 \geq m^2$;
- $\bar{\varphi}_t^\varepsilon$ is bounded independently of ε , $\Delta \bar{\varphi}^\varepsilon \leq C$, C positive constant independent of ε ;
- $1 \leq \eta'_\varepsilon(\bar{\varphi}^\varepsilon) = 1 + \varepsilon B \xi(\bar{\varphi}^\varepsilon) \leq 1 + \varepsilon B$;
- $\eta'_\varepsilon \frac{1 - (\eta'_\varepsilon)^2}{\varepsilon} g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon = -(2B\eta'_\varepsilon \xi + \varepsilon B^2 \eta'_\varepsilon \xi^2) g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon$.

To prove (42) we only need to find B and N sufficiently large, independent of ε , such that

$$\begin{aligned} k_\varepsilon \left\{ 2N \left[B^2 (\eta'_\varepsilon)^2 \xi (g_\varepsilon^2 + \varepsilon)^2 + \eta'_\varepsilon \frac{1 - (\eta'_\varepsilon)^2}{\varepsilon} g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \right] - \eta'_\varepsilon \Delta \bar{\varphi}^\varepsilon - \eta''_\varepsilon (g_\varepsilon^2 + \varepsilon) \right\} &\geq \\ &\geq \eta'_\varepsilon(\bar{\varphi}^\varepsilon) \|\bar{\varphi}_t^\varepsilon\|_{L^\infty(Q_T)} + \|f\|_{L^\infty(0,T)}, \end{aligned}$$

The second term is bounded from above by a positive constant C_1 independent of ε . Working with ε such that $\varepsilon \leq \frac{1}{B}$ and noting that $\|g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon\|_\infty \leq X_0$, X_0 not depending on ε , we see that

$$\begin{aligned} -\eta'_\varepsilon \Delta \bar{\varphi}^\varepsilon &\geq -(1 + \varepsilon B)C \geq -2C, \\ -\eta''_\varepsilon (g_\varepsilon^2 + \varepsilon) &\geq 0, \\ B^2 (\eta'_\varepsilon)^2 \xi (g_\varepsilon^2 + \varepsilon)^2 + \eta'_\varepsilon \frac{1 - (\eta'_\varepsilon)^2}{\varepsilon} g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \\ &\geq B^2 \xi m^4 - (2B\eta'_\varepsilon \xi + \varepsilon B^2 \eta'_\varepsilon \xi^2) X_0 \\ &= B\xi (Bm^4 - \eta'_\varepsilon (2 + \varepsilon B\xi) X_0) \\ &\geq B\xi (Bm^4 - 6X_0). \end{aligned}$$

Choose $B = (1 + 6X_0)/m^4$ (this choice comes from imposing that $Bm^4 - 6X_0 = 1$). So,

$$\begin{aligned} 2N \left[B^2 (\eta'_\varepsilon)^2 \xi (g_\varepsilon^2 + \varepsilon)^2 + \eta'_\varepsilon \frac{1 - (\eta'_\varepsilon)^2}{\varepsilon} g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \right] - \eta'_\varepsilon \Delta \bar{\varphi}^\varepsilon - \eta''_\varepsilon (g_\varepsilon^2 + \varepsilon) \\ \geq 2NB\xi (Bm^4 - 6X_0) - 2C \\ \geq 2NB\varepsilon^{-BC_0} - 2C \geq C_1 \geq 0, \end{aligned}$$

as long as $N \geq \frac{C_1 + 2C}{2B\varepsilon^{-BC_0}}$.

Then, since $k_\varepsilon \geq 1$, $C_1 \geq 0$

$$\begin{aligned} k_\varepsilon \left\{ 2N \left[B^2 (\eta'_\varepsilon)^2 \xi (g_\varepsilon^2 + \varepsilon)^2 + \eta'_\varepsilon \frac{1 - (\eta'_\varepsilon)^2}{\varepsilon} g_\varepsilon \nabla g_\varepsilon \cdot \nabla \bar{\varphi}^\varepsilon \right] - \eta'_\varepsilon \Delta \bar{\varphi}^\varepsilon - \eta''_\varepsilon (g_\varepsilon^2 + \varepsilon) \right\} \\ \geq k_\varepsilon C_1 \geq C_1, \end{aligned}$$

as we wanted to prove.

Since we have

$$\underline{\psi} \leq u^\varepsilon \leq \bar{\psi} \text{ in } Q_T, \quad \underline{\psi}(x, t) = u^\varepsilon(x, t) = \bar{\psi}(x, t) = 0 \text{ if } (x, t) \in \Sigma,$$

then

$$\forall (x, t) \in \Sigma \quad |\nabla u^\varepsilon(x, t)| \leq \max\{|\nabla \bar{\psi}(x, t)|, |\nabla \underline{\psi}(x, t)|\}.$$

But, for $(x, t) \in \Sigma$,

$$|\nabla \bar{\psi}(x, t)|^2 = (\eta'_\varepsilon (\bar{\varphi}_\varepsilon(x, t)))^2 |\nabla \bar{\varphi}_\varepsilon(x, t)| \leq (1 + \varepsilon B)^2 (g_\varepsilon^2(x, t) + \varepsilon) \leq g_\varepsilon^2(x, t) + C\varepsilon,$$

where C is a constant independent of ε . Analogously, $|\nabla \underline{\psi}(x, t)|^2 \leq g_\varepsilon^2(x, t) + C\varepsilon$.

Since $|\nabla u^\varepsilon(x, 0)| = |\nabla h_\varepsilon(x)| \leq g_\varepsilon(x, 0)$, the proof is concluded. ■

Proposition 3.7 *Suppose that the assumptions (3), (23) and (40) are verified. Then*

$$\exists C > 0 \forall (x, t) \in Q_T \quad |\nabla u^\varepsilon(x, t)|^2 \leq g_\varepsilon^2(x, t) + C\varepsilon. \quad (44)$$

Proof: Let $v = |\nabla u^\varepsilon|^2$ and $w = v - g_\varepsilon^2$. Since g satisfies (40), we may assume that the approximations g_ε of g also verify $(g_\varepsilon^2)_t \geq 0$ and $\Delta g_\varepsilon^2 \leq 0$. Differentiate the first equation of problem (4) with respect to x_k . Then,

$$\begin{aligned} u_{x_k t}^\varepsilon - k_\varepsilon''(w)w_{x_k}w_{x_i}u_{x_i}^\varepsilon - k_\varepsilon'(w)w_{x_i x_k}u_{x_i}^\varepsilon - k_\varepsilon'(w)w_{x_i}u_{x_i x_k}^\varepsilon \\ - k_\varepsilon'(w)w_{x_k}u_{x_i x_i}^\varepsilon - k_\varepsilon(w)u_{x_i x_i x_k}^\varepsilon = 0. \end{aligned} \quad (45)$$

Multiplying (45) by $u_{x_k}^\varepsilon$, summing over k , we obtain

$$\begin{aligned} u_{x_k}^\varepsilon u_{x_k t}^\varepsilon - k_\varepsilon''(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon w_{x_i}w_{x_k} - k_\varepsilon'(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon w_{x_i x_k} - k_\varepsilon'(w)u_{x_k}^\varepsilon u_{x_i x_k}^\varepsilon w_{x_i} - k_\varepsilon'(w)u_{x_k}^\varepsilon u_{x_i x_i}^\varepsilon w_{x_k} \\ - k_\varepsilon(w)u_{x_k}^\varepsilon u_{x_i x_i x_k}^\varepsilon = 0. \end{aligned}$$

Notice that

$$u_{x_k}^\varepsilon u_{x_i x_i x_k}^\varepsilon = \frac{1}{2}v_{x_i x_i} - (u_{x_k x_i}^\varepsilon)^2.$$

Then

$$\begin{aligned} \frac{1}{2}v_t + \left(-k_\varepsilon''(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon w_{x_i} - k_\varepsilon'(w)\left(u_{x_i}^\varepsilon u_{x_k x_i}^\varepsilon + u_{x_i x_i}^\varepsilon u_{x_k}^\varepsilon\right)\right)w_{x_k} - \left(k_\varepsilon'(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon\right)w_{x_i x_k} \\ - \frac{1}{2}k_\varepsilon(w)\Delta v + k_\varepsilon(w)(u_{x_i x_k}^\varepsilon)^2 = 0 \end{aligned}$$

and, denoting

$$a_{ik} = k_\varepsilon'(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon + \frac{1}{2}k_\varepsilon(w)\delta_{ij}, \quad b_k = -k_\varepsilon'(w)\left(u_{x_i}^\varepsilon u_{x_i x_k}^\varepsilon + u_{x_i x_i}^\varepsilon u_{x_k}^\varepsilon\right) - k_\varepsilon''(w)u_{x_i}^\varepsilon u_{x_k}^\varepsilon w_{x_i}$$

and recalling that $v = w + g_\varepsilon^2$, we obtain

$$\frac{1}{2}w_t - a_{ik}w_{x_i x_k} - b_k w_{x_k} \leq -\frac{1}{2}\left[\left(g_\varepsilon^2\right)_t - k_\varepsilon(w)\Delta\left(g_\varepsilon^2\right)\right],$$

and, by the assumption (40) and the previous proposition, we have, in fact,

$$\begin{cases} \frac{1}{2}w_t - a_{ik}w_{x_i x_k} - b_k w_{x_k} \leq 0, \\ w|_{\Sigma \cup \Omega_0} \leq C\varepsilon, \end{cases} \quad (46)$$

where C is a constant independent of ε .

Since u^ε is a function belonging to the class $C_{\alpha, \alpha/2}^{2,1}(Q_T)$ and k_ε is also a $C^{2,\alpha}$ function, the coefficients a_{ik} and b_k are Hölder continuous functions. On the other hand, we have $\sum_{i,k} a_{ik}\xi_i\xi_k \geq 0$ for all $\xi \in \mathbb{R}^n$. So, by the weak maximum principle for parabolic equations, if $z = w - C\varepsilon$,

$$z(x, t) \leq \max_{\Sigma \cup \Omega_0} \{z\} \leq 0,$$

by the previous proposition. Since $|\nabla u^\varepsilon|^2 = v = z + g_\varepsilon^2 + C\varepsilon$, we conclude (44). ■

Proposition 3.8 *Suppose that the assumption (23) is verified. Then*

$$\{u^\varepsilon : \varepsilon \in]0, 1[\} \text{ is bounded in } L^2(0, T; H_{loc}^2(\Omega)). \quad (47)$$

Proof: In this proof, since there is no risk of confusion, we are going to omit the subscripts and the superscripts ε .

Given $\Omega' \subset\subset \Omega$, let η belonging to $\mathcal{D}(\Omega)$ be such that $\eta|_{\Omega'} \equiv 1$. Multiply the equation of problem (4) by $-u_{x_k x_k} \eta^2$ and integrate over $Q_t = \Omega \times]0, t[$. Then

$$\int_{Q_t} -u_t u_{x_k x_k} \eta^2 + \int_{Q_t} \left(k(|\nabla u|^2 - g^2) u_{x_i} \right)_{x_i} u_{x_k x_k} \eta^2 = \int_{Q_t} f u_{x_k x_k} \eta^2.$$

We are going to consider each term of the equality above separately.

Notice that

$$\begin{aligned} \int_{Q_t} -u_t u_{x_k x_k} \eta^2 &= \int_{Q_t} \left(u_t \eta^2 \right)_{x_k} u_{x_k} \\ &= \int_0^t \int_{\Omega} \frac{1}{2} \left[(u_{x_k})^2 \right]_t \eta^2 + \int_0^t \int_{\Omega} 2u_t u_{x_k} \eta \eta_{x_k} \\ &= \frac{1}{2} \int_{\Omega} \left[(u_{x_k})^2(t) - (u_{x_k})^2(0) \right] \eta^2 + \int_{Q_t} 2u_t u_{x_k} \eta \eta_{x_k}. \end{aligned} \quad (48)$$

The second term of the equality above is treated as follows:

$$\begin{aligned} \int_{Q_t} \left(k(|\nabla u|^2 - g^2) u_{x_i} \right)_{x_i} u_{x_k x_k} \eta^2 &= - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} \left(u_{x_k x_k} \eta^2 \right)_{x_i} \\ &= - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} \left(u_{x_k x_k x_i} \eta^2 + u_{x_k x_k} 2\eta \eta_{x_i} \right) \\ &= \int_{Q_t} \left(k(|\nabla u|^2 - g^2) u_{x_i} \eta^2 \right)_{x_k} u_{x_k x_i} - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k} \\ &= \int_{Q_t} k'(|\nabla u|^2 - g^2) \left(2u_{x_j} u_{x_j x_k} - 2g g_{x_k} \right) u_{x_i} \eta^2 u_{x_i x_k} \\ &\quad + \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2 \\ &\quad + \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} u_{x_i x_k} 2\eta \eta_{x_k} - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k} \\ &= \int_{Q_t} k'(|\nabla u|^2 - g^2) \left(2u_{x_j} u_{x_j x_k} - 2g g_{x_k} \right) \left((u_{x_i} u_{x_i x_k} - g g_{x_k}) + g g_{x_k} \right) \eta^2 \\ &\quad + \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2 + \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} u_{x_i x_k} 2\eta \eta_{x_k} \\ &\quad - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k} \end{aligned} \quad (49)$$

$$\begin{aligned}
&= 2 \int_{Q_t} k'(|\nabla u|^2 - g^2) (u_{x_j} u_{x_j x_k} - g g_{x_k})^2 \eta^2 + \frac{1}{2} \int_{Q_t} [k(|\nabla u|^2 - g^2)]_{x_k} [g^2]_{x_k} \eta^2 \\
&\quad + \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2 + \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} u_{x_i x_k} 2\eta \eta_{x_k} \\
&\quad - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k}.
\end{aligned}$$

So, we have

$$\begin{aligned}
&\frac{1}{2} \int_{\Omega} [(u_{x_k})^2(t) - (u_{x_k})^2(0)] \eta^2 + \int_{Q_t} 2u_t u_{x_k} \eta \eta_{x_k} \\
&\quad + \int_{Q_t} k'(|\nabla u|^2 - g^2) (2u_{x_j} u_{x_j x_k} - 2g g_{x_k})^2 \eta^2 + \frac{1}{2} \int_{Q_t} [k(|\nabla u|^2 - g^2)]_{x_k} [g^2]_{x_k} \eta^2 \\
&\quad + \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2 + \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} u_{x_i x_k} 2\eta \eta_{x_k} \\
&\quad - \int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k} \eta^2 = - \int_{Q_t} f u_{x_k x_k} \eta^2.
\end{aligned} \tag{50}$$

Notice that

$$\begin{aligned}
&\int_{\Omega} (u_{x_k})^2(t) \eta^2 \geq 0, \quad \int_{\Omega} (u_{x_k})^2(0) = \int_{\Omega} (h_{x_k})^2 \eta^2, \\
&\int_{Q_t} 2u_t u_{x_k} \eta \eta_{x_k} \leq C_0, \quad C_0 \text{ constant independent of } \varepsilon, \\
&\int_{Q_t} k'(|\nabla u|^2 - g^2) (2u_{x_j} u_{x_j x_k} - 2g g_{x_k})^2 \eta^2 \geq 0, \\
&\int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2 \geq \int_{Q_t} (u_{x_i x_k})^2 \eta^2,
\end{aligned}$$

using Hölder and Young inequalities, we see that

$$\begin{aligned}
&\int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} u_{x_i x_k} 2\eta \eta_{x_k} \leq \\
&\quad \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i})^2 \eta_{x_k}^2 + \frac{1}{4} \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i x_k})^2 \eta^2,
\end{aligned}$$

and, obviously,

$$\int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i})^2 \eta_{x_k}^2 \leq C_1, \quad C_1 \text{ constant independent of } \varepsilon.$$

Analogously,

$$\int_{Q_t} k(|\nabla u|^2 - g^2) u_{x_i} 2\eta \eta_{x_i} u_{x_k x_k} \leq \int_{Q_t} k(|\nabla u|^2 - g^2) (u_{x_i})^2 \eta_{x_i}^2$$

$$+\frac{1}{4} \int_{Q_t} k(|\nabla u|^2 - g^2)(u_{x_k x_k})^2 \eta^2$$

and

$$\int_{Q_t} k(|\nabla u|^2 - g^2)(u_{x_i})^2 \eta_{x_i}^2 \leq C_1, \quad C_1 \text{ defined above.}$$

On the other hand,

$$\int_{Q_t} f u_{x_k x_k} \eta^2 \leq \frac{1}{4} \int_{Q_t} (u_{x_k x_k})^2 \eta^2 + \int_{Q_t} f^2 \eta^2$$

and

$$\int_{Q_t} \left[k(|\nabla u|^2 - g^2) \right]_{x_k} \left[g^2 \right]_{x_k} \eta^2 = - \int_{Q_t} k(|\nabla u|^2 - g^2) \left(\left[g^2 \right]_{x_k} \eta^2 \right)_{x_k} \leq C_2,$$

C_3 constant independent of ε .

Then,

$$\frac{1}{4} \int_{Q_T} (u_{x_i x_k})^2 \eta^2 \leq \int_{\Omega} (h_{x_k})^2 \eta^2 + C_0 + 2C_1 + 2C_2 + \int_{Q_T} f^2 \eta^2$$

and the proof is concluded. ■

Theorem 3.9 *With the assumptions (3), (23) and (40), problem (28) has a solution.*

Proof: We have proved that

$$\{u^\varepsilon : \varepsilon \in]0, 1[\} \text{ is uniformly bounded in } W = \{v \in L^2(0, T; H_{loc}^2(\Omega)) : v_t \in L^2(Q_T)\}.$$

If we consider W with the weak topology, we know that $\{u^\varepsilon : \varepsilon \in]0, 1[\}$ belongs to a compact subset of W (see [13], pg. 58). So, there exists $u \in W$ such that, for the weak topology, $u^\varepsilon \rightharpoonup u$ in this space, when $\varepsilon \rightarrow 0$.

So, $u^\varepsilon \rightarrow u$ strongly in $L^2(0, T; H_0^1(\Omega'))$, for all Ω' with smooth boundary and compactly included in Ω and $\nabla u^\varepsilon(x, t) \rightarrow \nabla u(x, t)$ for a.e. $(x, t) \in Q_T$.

Recalling that

$$\exists C > 0 \quad \forall (x, t) \in Q_T \quad |\nabla u^\varepsilon(x, t)|^2 \leq g_\varepsilon^2(x, t) + C\varepsilon,$$

we have

$$1 \leq k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) \leq e^{NC}.$$

So, with the additional assumptions introduced in this subsection, $k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2)$ is uniformly bounded not only in $L^1(Q_T)$, as we have proved in section 1, but also in $L^\infty(Q_T)$. So, there exists $\lambda \in L^\infty(Q_T)$ such that

$$k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) \rightharpoonup \lambda \quad \text{in } L^\infty(Q_T) \text{ weak-}^*, \quad \text{when } \varepsilon \rightarrow 0$$

and so

$$k_\varepsilon(|\nabla u^\varepsilon|^2 - g_\varepsilon^2) \nabla u^\varepsilon \rightharpoonup \lambda \nabla u \quad \text{weakly in } L^2(Q_T), \quad \text{when } \varepsilon \rightarrow 0.$$

Since $\|u_t^\varepsilon\|_{L^2(Q_T)} \leq C$, C independent of ε , we also have $u_t^\varepsilon \rightharpoonup u_t$ weakly in $L^2(Q_T)$ and so, passing to the limit in problem (4), we see that, in fact,

$$\begin{cases} u_t - \nabla \cdot (\lambda \nabla u) = f \text{ in } Q_T, \\ u|_\Sigma = 0, \quad u(0) = h, \end{cases}$$

and it only remains to prove that $\lambda \in k(|\nabla u|^2 - g_\varepsilon^2)$ to conclude that the pair (u, λ) is solution of problem (28).

Since $\Lambda = \{(x, t) \in Q_T : |\nabla u(x, t)| - g(x, t) < 0\}$ is a measurable set and

$$k_\varepsilon(|\nabla u^\varepsilon(x, t)|^2 - g_\varepsilon^2(x, t)) \longrightarrow 1 \quad \text{when } \varepsilon \rightarrow 0, \quad \text{for a.e. } (x, t) \in \Lambda,$$

$$k_\varepsilon(|\nabla u^\varepsilon(x, t)|^2 - g_\varepsilon^2(x, t)) \geq 1 \text{ for a.e. } (x, t) \in Q_T$$

we conclude that $\lambda = 1$ in Λ and $\lambda \geq 1$ in Q_T . So, $\lambda \in k(|\nabla u|^2 - g_\varepsilon^2)$, as we wanted to prove. ■

Theorem 3.10 *Suppose that the assumptions (3), (23) and (40) are verified. Then, if (u, λ) is a solution of problem (28), then u is solution of problem (2).*

Remark 3.11 *From this theorem we may conclude that, under the assumptions (3), (23) and (40), if (u, λ) is a solution of problem (28), then u is unique, but nothing is established about the uniqueness of λ . Uniqueness for λ was proved by Brézis in [1], for the elliptic case, with $g \equiv 1$ and homogeneous boundary condition. In [20], examples of non-uniqueness of λ can be found, when the boundary condition is not homogeneous.* ■

Proof of Theorem 3.10: Multiply the equation $u_t - \nabla \cdot (\lambda \nabla u) = f$ by $v(t) - u(t)$, with $v \in L^\infty(0, T; H_0^1(\Omega))$, $v(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in I$, and integrate over Ω , to obtain

$$\int_\Omega u_t(t)(v(t) - u(t)) + \int_\Omega ((\lambda(t) - 1) + 1) \nabla u(t) \cdot \nabla(v(t) - u(t)) = \int_\Omega f(t)(v(t) - u(t)).$$

Notice that

$$\begin{aligned} (\lambda(t) - 1) \nabla u(t) \cdot \nabla(v(t) - u(t)) &\leq (\lambda(t) - 1) |\nabla u(t)| [|\nabla v(t)| - |\nabla u(t)|] \\ &\leq (\lambda(t) - 1) |\nabla u(t)| [g(t) - |\nabla u(t)|] = 0, \end{aligned}$$

since $\lambda(x, t) = 1$ whenever $|\nabla u(x, t)| < g(x, t)$.

Then,

$$\int_\Omega u_t(t)(v(t) - u(t)) + \int_\Omega \nabla u(t) \nabla(v(t) - u(t)) \geq \int_\Omega f(t)(v(t) - u(t)), \quad \forall v \in \mathbb{K}_{g(t)}, \text{ for a.e. } t \in I,$$

as we wanted to prove. Besides that, $u(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in I$. ■

3.3 Equivalence with the equation with gradient constraint

In this subsection, the equation with gradient constraint (29) is considered. We begin proving existence of solution of problem (29) if assumptions (3) and

$$\begin{aligned} \Delta h &\in L^\infty(\Omega), & f &\in W^{1,\infty}(0, T; L^\infty(\Omega)), \\ -\Delta h(x) &\leq f(x, t) \text{ for a.e. } (x, t) \in Q_T, & (g^2)_t &\leq 0, \end{aligned} \tag{51}$$

are verified. We would like to remark that, in order to prove existence of solution, we don't need to assume that f is independent of the spatial variable x . The proof has many similarities with the proof presented by Zhu in [23], where a general linear parabolic equation with gradient constraint is considered in an unbounded domain and for arbitrarily large times, with a zero condition given at a fixed instant T , as well as with the proof of Evans ([5]), for the elliptic case.

The proof of equivalence between the variational inequality (2) and problem (29) is presented if assumptions (3), (40) and (51) are verified (which implies, in particular, that g is independent of t).

Theorem 3.12 *Suppose that the assumptions (3) and (51) are verified. Then problem (29) has a solution.*

Proof: Consider the following family of problems

$$\begin{cases} w_t^\varepsilon - \Delta w^\varepsilon + \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) = f^\varepsilon \text{ in } Q_T, \\ w^\varepsilon|_\Sigma = 0, & w^\varepsilon(0) = h_\varepsilon, \end{cases} \tag{52}$$

where $\gamma_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ is a C^2 , nondecreasing, convex function such that $\gamma_\varepsilon(s) = 0$ if $s \leq 0$ and $\gamma_\varepsilon(s) = \frac{s - \varepsilon}{\varepsilon}$ for $s \geq 2\varepsilon$, $f^\varepsilon \in C_{\alpha, \alpha/2}^{2,1}(Q_T)$, and $h_\varepsilon \in C_\alpha^2(\Omega)$ are approximations of f and h , respectively, satisfying $-\Delta h_\varepsilon \leq f^\varepsilon$ and $|\nabla h_\varepsilon|^2 \leq g^2 + \varepsilon$.

Problem (52) has a unique solution $w^\varepsilon \in C_{\alpha, \alpha/2}^{2,1}(\overline{Q}_T)$, by the classical theory of quasilinear parabolic equations (see [12], theorem 4.1, page 558).

Since $\gamma_\varepsilon \geq 0$, we have $w_t^\varepsilon - \Delta w^\varepsilon \leq f^\varepsilon$ and so, by the maximum principle for parabolic equations,

$$\exists C > 0 \text{ independent of } \varepsilon : \quad \|w^\varepsilon\|_{L^\infty(Q_T)} \leq C.$$

Let us prove now that w_t^ε is bounded in $L^\infty(Q_T)$ independently of ε : differentiate the first equation of problem (52) in order to t and call $z = w_t^\varepsilon$. Then

$$\begin{cases} z_t - \Delta z + \gamma'_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)(2w_{x_i}^\varepsilon z_{x_i} - (g^2)_t) = f_t^\varepsilon \text{ in } Q_T, \\ z|_\Sigma = 0, & z(0) = \Delta h_\varepsilon + f^\varepsilon(0), \end{cases}$$

Since $\gamma'_\varepsilon \geq 0$ and $(g^2)_t \leq 0$, we have, in fact that

$$z_t - \Delta z + b_i z_{x_i} \leq f_t^\varepsilon, \quad \text{where } b_i = 2\gamma'_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)w_{x_i}^\varepsilon.$$

So, by the maximum principle

$$\exists C_1 > 0 \text{ independent of } \varepsilon : \quad \|w_t^\varepsilon\|_{L^\infty(0,T;L^2(\Omega))} \leq C_1. \quad (53)$$

The next step consists in obtaining uniform gradient estimates (independent of ε) for ∇w^ε . Let φ^ε be defined as in (24) such that $|\nabla \varphi^\varepsilon|^2 = g^2 + M\varepsilon$, where M is a constant to be chosen later. It is easy to verify that, for M sufficiently big ($M \geq \|\varphi_t^\varepsilon\|_{L^\infty(Q_T)} + \sup_{Q_T} \{\Delta \varphi^\varepsilon\} + \|f^\varepsilon\|_{L^\infty(Q_T)}$), independent of ε , then φ^ε is a supersolution of problem (52). On the other hand, h_ε is obviously a subsolution of the same problem. Then

$$h_\varepsilon \leq w^\varepsilon \leq \varphi^\varepsilon \quad \text{in } Q_T, \quad h_{\varepsilon|_\Sigma} = w_{|\Sigma}^\varepsilon = \varphi_{|\Sigma}^\varepsilon, \quad h_\varepsilon = w^\varepsilon(0) \leq \varphi^\varepsilon(0),$$

and so

$$|\nabla w^\varepsilon(x, t)|^2 \leq \max\{|\nabla h_\varepsilon(x)|^2, |\nabla \varphi^\varepsilon(x, t)|^2\} \leq g^2(x, t) + \varepsilon + M\varepsilon \quad \text{for } (x, t) \in \Sigma \cup \Omega_0.$$

Let $v = |\nabla w^\varepsilon|^2 - w^\varepsilon$. The maximum of v may be attained at the *parabolic boundary* $\Sigma \cup \Omega_0$ or at $Q_T \setminus (\Sigma \cup \Omega_0)$. If the first case happens, since $|\nabla w^\varepsilon|$ and w^ε are bounded independently of ε on $\Sigma \cup \Omega_0$, then the bound of v , and consequently, the bound of $|\nabla w^\varepsilon|$ is independent of ε . Let us consider now the second case, i.e. the maximum of v is attained at a point $(x_0, t_0) \notin \Sigma \cup \Omega_0$. Then, at this point (x_0, t_0) , we have

$$v_{x_i} = 0, \quad v_t = 0, \quad v_t - \Delta v \geq 0.$$

Since

$$v_t = 2w_{x_j}^\varepsilon w_{x_j t}^\varepsilon - w_t^\varepsilon, \quad \Delta v = 2\left(w_{x_i x_j}^\varepsilon\right)^2 + 2w_{x_j}^\varepsilon \Delta w_{x_j}^\varepsilon - \Delta w^\varepsilon,$$

and

$$\Delta w^\varepsilon = w_t^\varepsilon + \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) - f^\varepsilon, \quad \Delta w_{x_j}^\varepsilon = w_{x_j t}^\varepsilon + \gamma'_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)(|\nabla w^\varepsilon|^2 - g^2)_{x_j},$$

omitting the argument of γ_ε to simplify, we get at (x_0, t_0) ,

$$\begin{aligned} 0 \leq v_t - \Delta v &= 2w_{x_j}^\varepsilon w_{x_j t}^\varepsilon - w_t^\varepsilon - 2\left(w_{x_i}^\varepsilon w_{x_j}^\varepsilon\right)^2 - 2w_{x_j}^\varepsilon \left[w_{x_j t}^\varepsilon + \gamma'_\varepsilon(-)(|\nabla w^\varepsilon|^2 - g^2)_{x_j}\right] \\ &\quad + [w_t^\varepsilon + \gamma_\varepsilon(-) - f^\varepsilon] \\ &\leq -2\gamma'_\varepsilon(-)w_{x_j}^\varepsilon(|\nabla w^\varepsilon|^2 - g^2)_{x_j} + [\gamma_\varepsilon(-) + \|f^\varepsilon\|_{L^\infty(Q_T)}]. \end{aligned}$$

Since γ_ε is convex, we have, $\forall s \in \mathbb{R}$ $\gamma_\varepsilon(s) \leq \gamma'_\varepsilon(s)s$ and, since $v_{x_i} = 0$, we have $(|\nabla w^\varepsilon|^2)_{x_i} = w_{x_i}^\varepsilon$. On the other hand, we may suppose that $\gamma_\varepsilon(-) \geq 2\|f^\varepsilon\|_{L^\infty(Q_T)}$ at the point (x_0, t_0) , because, otherwise, we would have already obtained the bound for $|\nabla w^\varepsilon|$. So,

$$[\gamma_\varepsilon(-) + \|f^\varepsilon\|_{L^\infty(Q_T)}] \leq \gamma_\varepsilon(-) + \frac{\gamma_\varepsilon(-)}{2\|f^\varepsilon\|_{L^\infty(Q_T)}} \|f^\varepsilon\|_{L^\infty(Q_T)} = \frac{3}{2}\gamma_\varepsilon(-) \leq \frac{3}{2}\gamma'_\varepsilon(-)(|\nabla w^\varepsilon|^2 - g^2)$$

and so, at (x_0, t_0)

$$0 \leq v_t - \Delta v \leq \gamma'_\varepsilon(-) \left[-2|\nabla w^\varepsilon|^2 + 2\nabla g^2 \cdot \nabla w^\varepsilon + \frac{3}{2}(|\nabla w^\varepsilon|^2 - g^2) \right].$$

Then, $-\frac{1}{2}|\nabla w^\varepsilon|^2 + 2\nabla(g^2) \cdot \nabla w^\varepsilon - \frac{3}{2}g^2 \geq 0$ and we get

$$|\nabla w^\varepsilon(x_0, t_0)|^2 \leq 16|\nabla g^2(x_0, t_0)|^2 - 6g^2(x_0, t_0) \leq 16|\nabla g^2(x_0, t_0)|^2$$

and, since

$$v(x, t) = |\nabla w^\varepsilon(x, t)|^2 - w^\varepsilon(x, t) \leq v(x_0, t_0) = |\nabla w^\varepsilon(x_0, t_0)|^2 - w^\varepsilon(x_0, t_0),$$

because $|\nabla w^\varepsilon(x, t)|^2 = v(x, t) + w^\varepsilon(x, t)$ we get that

$$|\nabla w^\varepsilon(x, t)|^2 \leq v(x_0, t_0) + w^\varepsilon(x, t) \leq 16 \max_{(x,t) \in Q_T} |\nabla g^2(x, t)|^2 + 2\|w^\varepsilon\|_{L^\infty(Q_T)},$$

concluding then that ∇w^ε is bounded in Q_T , independently of ε .

Remark 3.13 Notice that, if $g \equiv 1$, then the maximum of v is attained at the parabolic boundary.

The next step consists in proving that $\gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)$ is locally bounded independently of ε : given $\Omega' \subset\subset \Omega$, let ζ belong to $\mathcal{D}(\Omega)$ be such that $\zeta|_{\Omega'} \equiv 1$. Define $v = \zeta^2 \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)$. As before, if $\max_{Q_T} v$ is attained at $(x_0, t_0) \in \Omega_0$ (notice that on Σ we have $v(x_0, t_0) = 0$) then $\max_{Q_T} v \leq \zeta^2(x_0, t_0) \gamma_\varepsilon(g^2(x_0, t_0) + M\varepsilon - g^2(x_0, t_0)) = \zeta^2(x_0, t_0)(M - 1)$ is independent of ε . If the maximum of v is attained at $(x_0, t_0) \in Q_T \setminus \Sigma \cup \Omega_0$ then, at this point, we have $v_t - \Delta v \geq 0$ and also $v_{x_i} = 0$.

Remarking that all the calculations below are done in the point (x_0, t_0) , we have

$$\begin{aligned} v_t &= \zeta^2 \gamma'_\varepsilon(-)(2w_{x_j}^\varepsilon w_{x_j t}^\varepsilon - (g^2)_t) \\ \Delta v &= \zeta^2 \gamma''_\varepsilon(-) \left[(|\nabla w^\varepsilon|^2 - g^2)_{x_i} \right]^2 + 2(\zeta^2)_{x_i} \gamma'_\varepsilon(-) (|\nabla w^\varepsilon|^2 - g^2)_{x_i} + \Delta \zeta^2 \gamma_\varepsilon(-) \\ &\quad + \zeta^2 \gamma'_\varepsilon(-) \left[2(w_{x_j x_i}^\varepsilon)^2 + 2w_{x_j}^\varepsilon \Delta w_{x_j}^\varepsilon - \Delta(g^2) \right], \end{aligned}$$

and, since $\zeta^2 \gamma''_\varepsilon(-) [(|\nabla w^\varepsilon|^2 - g^2)_{x_i}]^2 \geq 0$, using the calculations presented above for $\Delta w_{x_j}^\varepsilon$,

$$\begin{aligned} 0 &\leq v_t - \Delta v \leq \zeta^2 \gamma'_\varepsilon(-) \left[2w_{x_j}^\varepsilon w_{x_j t}^\varepsilon - (g^2)_t \right] - 2(\zeta^2)_{x_i} \gamma'_\varepsilon(-) (|\nabla w^\varepsilon|^2 - g^2)_{x_i} - \Delta \zeta^2 \gamma_\varepsilon(-) \\ &\quad - \zeta^2 \gamma'_\varepsilon(-) \left[2(w_{x_j x_i}^\varepsilon)^2 + 2w_{x_j}^\varepsilon (w_{x_j t}^\varepsilon + \gamma'_\varepsilon(-) (|\nabla w^\varepsilon|^2 - g^2)_{x_j}) - \Delta(g^2) \right]. \end{aligned}$$

Recall that, since γ_ε is a convex function, we have, $\forall s \in \mathbb{R}$, $\gamma'_\varepsilon(s)s \geq \gamma_\varepsilon(s)$. So, since $|\Delta \zeta^2| \leq C$, C independent of ε ,

$$\begin{aligned} 0 &\leq \gamma'_\varepsilon(-) \left[-\zeta^2 (g^2)_t - 2(\zeta^2)_{x_i} (|\nabla w^\varepsilon|^2 - g^2)_{x_i} + C(|\nabla w^\varepsilon|^2 - g^2) \right. \\ &\quad \left. - 2\zeta^2 (w_{x_i x_j}^\varepsilon)^2 - 2\zeta^2 \gamma'_\varepsilon(-) w_{x_i}^\varepsilon (|\nabla w^\varepsilon|^2 - g^2)_{x_i} + \zeta^2 \Delta g^2 \right] \end{aligned}$$

and, since $v_{x_i} = 0$ at the point considered, then, at that point, $\zeta^2 \gamma'_\varepsilon(-)(|\nabla w^\varepsilon|^2 - g^2)_{x_i} = -(\zeta^2)_{x_i} \gamma_\varepsilon(-)$ and we get

$$\begin{aligned} & -\zeta^2 (g^2)_t - 2 (\zeta^2)_{x_i} (|\nabla w^\varepsilon|^2 - g^2)_{x_i} + C(|\nabla w^\varepsilon|^2 - g^2) - 2\zeta^2 (w^\varepsilon_{x_i x_j})^2 \\ & + 2(\zeta^2)_{x_i} w^\varepsilon_{x_i} \gamma_\varepsilon(-) + \zeta^2 \Delta g^2 \geq 0. \end{aligned}$$

But

$$\begin{aligned} -2 (\zeta^2)_{x_i} (|\nabla w^\varepsilon|^2 - g^2)_{x_i} &= -4\zeta \zeta_{x_i} \left(2w^\varepsilon_{x_j} w^\varepsilon_{x_j x_i} - (g^2)_{x_i} \right) \\ &\leq \zeta^2 (w^\varepsilon_{x_i x_j})^2 + 16(w^\varepsilon_{x_j})^2 (\zeta_{x_i})^2 + 4\zeta \nabla \zeta \cdot \nabla g^2 \end{aligned} \quad (54)$$

and

$$2(\zeta^2)_{x_i} w^\varepsilon_{x_i} \gamma_\varepsilon(-) = 4\zeta \nabla \zeta \cdot \nabla w^\varepsilon \gamma_\varepsilon(-) \leq \frac{1}{4} \zeta^2 \gamma_\varepsilon^2(-) + 16|\nabla \zeta|^2 |\nabla w^\varepsilon|^2.$$

Then,

$$\begin{aligned} & -\zeta^2 (g^2)_t + \zeta^2 (w^\varepsilon_{x_i x_j})^2 + 16(w^\varepsilon_{x_j})^2 (\zeta_{x_i})^2 + 4\zeta \nabla \zeta \cdot \nabla g^2 + C(|\nabla w^\varepsilon|^2 - g^2) - 2\zeta^2 (w^\varepsilon_{x_i x_j})^2 \\ & + \frac{1}{4} \zeta^2 \gamma_\varepsilon^2(-) + 16|\nabla \zeta|^2 |\nabla w^\varepsilon|^2 + \zeta^2 \Delta g^2 \geq 0. \end{aligned}$$

So, there exists a constant C_0 , independent of ε , such that

$$\zeta^2 (w^\varepsilon_{x_i x_j})^2 \leq C_0 + \frac{1}{4} \zeta^2 \gamma_\varepsilon^2(|\nabla w^\varepsilon|^2 - g^2)$$

and then, at (x_0, t_0) , we have

$$\exists C_1 > 0 \text{ independent of } \varepsilon : \quad \zeta |w^\varepsilon_{x_i x_j}| \leq C_1 + \frac{1}{2} \zeta \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2).$$

Then, since $\zeta \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) = \zeta f - \zeta w_t^\varepsilon + \zeta \Delta w^\varepsilon$, we have, at the point (x_0, t_0) ,

$$\zeta \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) \leq \zeta \|f\|_{L^\infty(Q_T)} + \zeta \|w_t^\varepsilon\|_{L^\infty(Q_T)} + C_1 + \frac{1}{2} \zeta \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)$$

and so

$$\frac{1}{2} \zeta(x, t) \gamma_\varepsilon(|\nabla w^\varepsilon(x, t)|^2 - g^2(x, t)) \leq \frac{1}{2} \zeta(x_0, t_0) \gamma_\varepsilon(|\nabla w^\varepsilon(x_0, t_0)|^2 - g^2(x_0, t_0)) \leq C_2,$$

where C_2 is a constant independent of ε .

Now, if $\tilde{w} = \zeta w^\varepsilon$, we have,

$$\begin{cases} \tilde{w}_t - \Delta \tilde{w} = \zeta f - \zeta \gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) - (\Delta \zeta) w^\varepsilon - 2\nabla \zeta \cdot \nabla w^\varepsilon = \Phi, \\ \tilde{w}|_\Sigma = 0, \quad \tilde{w}(0) = \zeta h \end{cases}$$

and so, since Φ is bounded in $L^\infty(Q_T)$, independently of ε , then \tilde{w} is bounded in $W_p^{2,1}(Q_T)$, $1 < p < +\infty$, independently of ε (see [12], theorem 9.1, page 341).

Since $\{w^\varepsilon : \varepsilon \in [0, 1]\}$ is bounded in $W_{p,loc}^{2,1}(Q_T)$, let u be the weak limit of $(w^\varepsilon)_\varepsilon$ in this space (at least for a subsequence), when $\varepsilon \rightarrow 0$. Of course, we also have

$$w_t^\varepsilon \rightharpoonup u_t \quad \text{when } \varepsilon \rightarrow 0, \quad \text{in } L^\infty(Q_T) \text{ weak } - *,$$

$$w^\varepsilon \rightharpoonup u \quad \text{when } \varepsilon \rightarrow 0, \quad \text{in } L^p(0, T; W^{1,p}(\Omega')), \quad \text{for any } \Omega' \subset\subset \Omega, \quad 1 < p < +\infty,$$

and, since $\|w^\varepsilon\|_{W^{1,\infty}(\Omega \times [0, T])} \leq C$, C independent of ε , we also have, due to the compact inclusion $W^{1,\infty}(Q_T) \hookrightarrow C^{0,1}(\overline{Q_T})$,

$$w^\varepsilon \longrightarrow u \quad \text{uniformly in } \overline{Q_T}$$

and, in particular, $w^\varepsilon(x, 0) = h(x) \longrightarrow u(x, 0) = h(x)$, when $\varepsilon \rightarrow 0$.

Since $\gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2)$ is locally bounded, independently of ε , we must have $|\nabla u| \leq g$ a.e. in Q_T . On the other hand, $\gamma_\varepsilon(|\nabla w^\varepsilon|^2 - g^2) \rightharpoonup \chi$, in $L^\infty(\Omega' \times [0, T])$ weak-*, for every $\Omega' \subset\subset \Omega$, when $\varepsilon \rightarrow 0$. Letting $\varepsilon \rightarrow 0$ in problem (52), we see that

$$u_t - \Delta u + \chi = f, \quad |\nabla u| \leq g, \quad u|_\Sigma = 0, \quad u(0) = h.$$

It only remains to prove that, whenever $|\nabla u| < g$ we have $\chi = 0$. Given $x_0 \in \Omega$, let Ω' be such that $x_0 \in \Omega' \subset\subset \Omega$. Since $w^\varepsilon \rightharpoonup u$ in $W_p^{2,1}(\Omega' \times [0, T])$, for $1 < p < +\infty$, when $\varepsilon \rightarrow 0$, and $W_p^{2,1}(\Omega' \times [0, T]) \hookrightarrow C_{\alpha, \alpha/2}^{1,0}(\overline{\Omega'} \times [0, T])$, if $p > n$ (being this inclusion compact) then, if (x_0, t_0) is such that $|\nabla u(x_0, t_0)| < g(x_0, t_0)$, we have, for ε sufficiently small, $\gamma_\varepsilon(|\nabla w^\varepsilon(x_0, t_0)|^2 - g^2(x_0, t_0)) = 0$. So, letting $\varepsilon \rightarrow 0$, we conclude that $\chi(x_0, t_0) = 0$ and the result follows. \blacksquare

Proposition 3.14 *Suppose that the assumptions (3), (40) and (51) are verified. Then problems (2) and (29) are equivalent.*

Proof: Let u denote a solution of problem (29) and u^* the solution of problem (27).

Recall the family of penalized problems (36) for the double obstacle variational inequality problem:

$$\begin{cases} z_t^\varepsilon - \Delta z^\varepsilon + \frac{1}{\varepsilon} (z^\varepsilon - (z^\varepsilon \wedge \overline{\varphi}) \vee \underline{\varphi}) = f & \text{in } Q_T, \\ z^\varepsilon(0) = h, \quad z^\varepsilon|_\Sigma = 0. \end{cases}$$

Let us call $\Phi_\varepsilon(v) = \gamma_\varepsilon(|\nabla v|^2 - g^2)$ and $\Psi_\varepsilon(v) = \frac{1}{\varepsilon} (v - (v \wedge \overline{\varphi}) \vee \underline{\varphi})$.

Notice that:

- since $|\nabla u| \leq g$ then $\underline{\varphi} \leq u \leq \overline{\varphi}$ and so $\Psi_\varepsilon(u) = 0$;
- $u_t - \Delta u \leq f$, since u is solution of problem (29).

So, u is a subsolution of the problem (36) and, due to the monotonicity of Ψ_ε , we have $u \leq z^\varepsilon$ and, passing to the limit when $\varepsilon \rightarrow 0$,

$$u \leq u^*.$$

On the other hand

- since problems (2) and (27) are equivalent, we have $|\nabla u^*| \leq g$;
- $-\Delta h \leq f$ and $|\nabla h| \leq g$ (and consequently $\underline{\varphi} \leq h \leq \overline{\varphi}$) implies that h is a subsolution of the problem (36) and consequently, $\underline{\varphi} \leq \underline{h} \leq z^\varepsilon$;
- passing to the (weak) limit when $\varepsilon \rightarrow 0$ in the equation of problem (36), we conclude that $u_t^* - \Delta u^* + \chi^* = f$; notice that $\chi^* = \lim_\varepsilon \Psi_\varepsilon(z^\varepsilon) \geq 0$, since $z^\varepsilon \geq \underline{\varphi}$;
- since $u_t^* - \Delta u^* \leq f$ and $|\nabla u^*| \leq g$ we know that u^* is a subsolution of problem (52)

The monotonicity of Φ_ε implies that $u^* \leq w^\varepsilon$ and so, letting $\varepsilon \rightarrow 0$,

$$u^* \leq u.$$

So $u = u^*$ and both problems are equivalent.

In particular, since u^* is unique, we proved that problem (29) has a unique solution. ■

3.4 A counter-example

This subsection is dedicated to present a counter-example. We prove that problem (2) is not always equivalent to problem (27), as well as to problem (29), presenting data for which the solutions of problems (2), (27) and (29) are different. Detailing more, we are going to present an example to show that, if $(g^2)_t - \Delta(g^2) \not\geq 0$, then the problems considered in section 3 may not be equivalent.

It is important to note that the data chosen here do not satisfy completely the assumption (3), since the (very smooth) gradient constraint is zero in one point. Nevertheless, the nonzero gradient constraint condition is used in the previous sections only to prove existence of solution and not the equivalence among these problems. Since the solution of problem (2), for the chosen data, will be calculated here explicitly, there is no question about the existence of solution.

Let

$$\begin{array}{ll} f :]-1, 1[\times]0, T[& \rightarrow \mathbb{R} & g :]-1, 1[\times]0, T[& \rightarrow \mathbb{R} \\ x & \mapsto 2 & x & \mapsto 3x^2 \end{array}$$

Remark 3.15 *In fact $(g^2)_t(x, t) - \Delta(g^2)(x, t) = -108x^2 \not\geq 0$.*

Easy calculations show that the two obstacles (with respect to this function g) are

$$\overline{\varphi}(x, t) = \begin{cases} x^3 + 1 & \text{if } x \in [-1, 0[, \\ 1 - x^3 & \text{if } x \in [0, 1], \end{cases} \quad \text{and} \quad \underline{\varphi}(x, t) = \begin{cases} -x^3 - 1 & \text{if } x \in [-1, 0[, \\ x^3 - 1 & \text{if } x \in [0, 1]. \end{cases}$$

Let

$$h(x) = \begin{cases} 1 - x^2 & \text{if } |x| \geq \frac{2}{3} \text{ and } |x| \leq 1, \\ \overline{\varphi}(x, 0) - \frac{4}{27} & \text{otherwise.} \end{cases}$$

We would like to remark that the function h is a C^1 function. Defining $w(x, t) \equiv h(x)$, we see that $|\nabla w(x, t)| = 2|x| \leq g(x, t)$, if $|x| > \frac{2}{3}$ and $|\nabla w(x, t)| = 3x^2 = g(x, t)$ if $|x| < \frac{2}{3}$. So, $w(t) \in \mathbb{K}_{g(t)}$ for all $t \in [0, 1]$ and $w(0) = h$. We are going to prove that w is, in fact, the solution of problem (2), for the given data (f, g, h) . Recalling that w is of class C^1 , we have

$$\begin{aligned}
& \int_{-1}^1 w_t(t)(v(t) - w(t))dx + \int_{-1}^1 w_x(t)(v(t) - w(t))_x dx - \int_{-1}^1 2(v(t) - w(t))dx \\
&= \int_{-1}^1 h'(x)(v_x(x, t) - h'(x))dx - \int_{-1}^1 2(v(t) - h)dx \\
&= - \int_{-1}^1 h''(x)(v(x, t) - h(x))dx - \int_{-1}^1 2(v(t) - h)dx \\
&= \int_{\{|x| \leq \frac{2}{3}\}} (-h''(x) - 2)(v(x, t) - h(x))dx + \int_{\{|x| \geq \frac{2}{3}\}} (-h''(x) - 2)(v(x, t) - h(x))dx \\
&= \left[(-3x^2 - 2x)(v(x, t) - h(x)) \right]_{-\frac{2}{3}}^0 - \int_{-\frac{2}{3}}^0 (-3x^2 - 2x)(v_x(x, t) - h'(x)) \\
&\quad + \left[(3x^2 - 2x)(v(x, t) - h(x)) \right]_0^{\frac{2}{3}} - \int_0^{\frac{2}{3}} (3x^2 - 2x)(v_x(x, t) - h'(x))dx \\
&= \int_{-\frac{2}{3}}^0 (3x^2 + 2x)(v_x(x, t) - 3x^2) + \int_0^{\frac{2}{3}} (-3x^2 + 2x)(v_x(x, t) + 3x^2)dx \geq 0,
\end{aligned}$$

as long as v is such that $v(t) \in \mathbb{K}_{g(t)}$ for a.e. $t \in [-1, 1]$, since, in that case, we have $-3x^2 \leq v_x(x, t) \leq 3x^2$.

If u denotes the solution of problem (27), it is easy to verify that $u(t) \rightarrow u^\infty$, when $t \rightarrow +\infty$, in $L^2(\Omega)$, where u^∞ is the solution of the problem

$$\int_{-1}^1 u_x^\infty (v - u^\infty)_x \geq \int_{-1}^1 2(v - u^\infty), \quad \forall v \in \mathbb{K},$$

where $\mathbb{K} = \{v \in H_0^1(-1, 1) : \underline{\varphi} \leq v \leq \bar{\varphi} \text{ a.e.}\}$. Since $z(x) = 1 - x^2$ is such that $z \in \mathbb{K}$ and $z'' = -2$, obviously, $z = u^\infty$.

Now, if problems (2) and (27) were equivalent, we should have $\lim_{t \rightarrow +\infty} w(t) = u^\infty$ in $L^2(\Omega)$, which obviously does not happen.

On the other hand, w is not a solution of problem (29) since, although $|\nabla w| \leq g$ a.e., the function w does not verify $w_t - \Delta w \leq 2$ a.e. in Q_T .

It was then shown that, for the given data, problem (2) is not equivalent to problem (27) nor to problem (29).

References

- [1] Brandt, E. H., *Electric field in superconductor with regular cross-sections*, Phys. Reviews **B 52** (1995) 15442 – 15457.

- [2] Brézis, H., *Multiplicateur de Lagrange en torsion elastoplastique*, Arch. Rat. Mech. Analysis **49** (1972) 32 – 49.
- [3] Caffarelli, L. A. & Friedman, A., *Reinforcement problems in elasto-plasticity*, Roc. Mount. J. Math. **10** (1980) 155 – 184.
- [4] Choe, Hi Jun & Shim, Yong Sun, *Degenerate variational inequalities with gradient constraints*, Annali Scuola Normal Sup. Pisa (IV) **22** (1995) 25 – 53.
- [5] Evans, L., *A second order elliptic equation with gradient constraint*, Comm. Part. Diff. Eq. **4** (1979) 555 – 572.
- [6] Gerhardt, C., *On the Existence and Uniqueness of a Warpening Function in the Elastic-plastic Torsion of a Cylindrical Bar with a Multiply Connected Cross-section*, Proceedings of the “Joint Symposium IUTAM/IMU Marseille, Dold, A. and Eckmann, B. and Germain, P. and Nayroles, B. (editors), (1975).
- [7] Haraux, A., *Nonlinear evolution equations. Global behavior of solutions*, Lect. Notes in Math. **841**, Springer-Verlag, 1981.
- [8] Ishii, H. & Koike, S., *Boundary regularity and uniqueness for an elliptic equation with gradient constraint*, Comm. in Partial Diff. Eq. Vol.8 **4** (1983) 317 – 346.
- [9] Jensen, R., *Regularity for elasto-plastic type variational inequalities*, Indiana Univ. Math. Journal **32** 3 (1983) 407 – 423.
- [10] Kunze, M. M. & Monteiro Marques, M. D. P., *A note on Lipschitz continuous solutions of a parabolic quasi-variational inequality*, “Proceedings of the Luso-Chinese Symposium on Nonlinear Evolutionary Equations and their Applications” Macau, Ta-Tsien, L. & Long-Wei, L. & Rodrigues, J. F. (editors), World Scientific, 109 – 115.
- [11] Kunze, M. & Rodrigues, J. F., *An elliptic quasi-variational inequality with gradient constraints and some of its applications*, Math. Meth. Appl. Sci., to appear.
- [12] Ladyzenskaja, O. A., Solonnikov, V. A. & Ural’ceva, N. N., *Linear and Quasilinear Equations of Parabolic Type*, Translations of Mathematical Monographs **23**, AMS, 1968.
- [13] Lions, J. L., *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [14] Lions, J. L., *Sur le contrôle optimale de systèmes gouvernés par des équations aux dérivées partielles*, Dunod, Paris, 1968.
- [15] Lions, P. L., *Generalized solutions fo Hamilton-Jacobi equations*, Pitman Advanced Publishing Program, Boston-London-Melbourne, 1982.
- [16] Prighozin, L., *Variational model of sandpile growth*, Europ. J. Appl. Math. **7** (1996) 225 – 235.
- [17] Prighozin, L., *On the Bean critical state model in superconductivity*, Europ. J. Appl. Math. **7** (1996) 237 – 247.

- [18] Rodrigues, J. F., *Obstacle Problems in Mathematical Physics*, North Holland, Amsterdam, 1987.
- [19] Rodrigues, J. F. & Santos, L., *A parabolic quasi-variational inequality arising in a superconductivity model*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) Vol. XXIX (2000) 153 – 169.
- [20] Santos, L., *A diffusion problem with gradient constraint and evolutive Dirichlet condition*, Portugaliae Mathematica Vol.48 Fasc.4 (1991) 441 – 468.
- [21] Santos, L., *A parabolic variational inequality with non-constant gradient constraint*, “Proceedings of the Luso-Chinese Symposium on Nonlinear Evolutionary Equations and their Applications” Macau, Ta-Tsien, L. & Long-Wei, L. & Rodrigues, J. F. (editors), World Scientific, 203 – 212.
- [22] Simon, J., *Compact Sets in the Space $L^p(0, T; B)$* , Annali Mat. Pura et Appl. CXLVI, (1987), 65 – 96.
- [23] Zhu, H., *Characterization of variational inequalities in singular stochastic control*, Ph.D. thesis, Brown University, U.S.A., 1992.