

Geologia dell'Ambiente

Periodico trimestrale della SIGEA
Società Italiana di Geologia Ambientale



Supplemento al n. 3/2012

ISSN 1591-5352



Bari - Italy, 24-28 September 2012

Geoheritage: Protecting and Sharing

7th International Symposium ProGEO on the Conservation of the Geological Heritage
3rd Regional Meeting of the ProGEO SW Europe Working Group

www.geoheritagesymposium-bari2012.org



preprint

Poste Italiane S.p.a. - Spedizione in Abbonamento Postale - D.L. 353/2003 (conv. in L. 27/02/2004 n° 46) art. 1 comma 1 - DCB Roma

Geologia dell'Ambiente
Periodico trimestrale della SIGEA
Società Italiana di Geologia Ambientale

Supplemento al n. 3/2012
Anno XX - luglio-settembre 2012

a cura di **VILLAGGIO**
editore

Iscritto al Registro Nazionale della Stampa n. 06352
Autorizzazione del Tribunale di Roma n. 229
del 31 maggio 1994

Comitato scientifico

Mario Bentivenga, Aldino Bondesan,
Giancarlo Bortolami, Aldo Brondi,
Felice Di Gregorio, Giuseppe Gisotti,
Giancarlo Guado, Gioacchino Lena,
Giacomo Prosser, Giuseppe Spilotro

Consiglio Direttivo nazionale 2010-2013

Davide Baioni, Domenico Bartolucci,
Federico Boccalaro, Giancarlo Bortolami,
Antonio Fiore (*Tesoriere*), Fabio Garbin (*Segretario*),
Francesco Geremia, Giuseppe Gisotti (*Presidente*),
Gioacchino Lena (*Vice Presidente*),
Massimo Massellani, Vincent Ottaviani,
Andrea Vitturi, Francesco Zarlenga

Comitato di redazione

Federico Boccalaro, Giorgio Cardinali,
Giovanni Conte, Gioacchino Lena,
Paola Mauri, Maurizio Scardella

Direttore responsabile

Giuseppe Gisotti

Procedura per l'accettazione degli articoli

I lavori sottomessi alla rivista dell'Associazione,
dopo che sia stata verificata la loro pertinenza
con i temi di interesse della Rivista, saranno
sottoposti ad un giudizio di uno o più Referees.

Redazione

SIGEA: tel./fax 06 5943344
Casella Postale 2449 U.P. Roma 158
info@sigeaweb.it
www.sigeaweb.it

Progetto grafico e impaginazione

Angelo Perrini
angelo_perrini@fastwebnet.it

Pubblicità

SIGEA

Stampa

Tipolitografia Acropoli, Alatri - FR

Abbonamento annuale: Euro 30,00

Sommario

| | |
|--|-----|
| PLENARY SESSION - INVITED SPEAKERS | 5 |
| GEOSITES | 9 |
| GEOLOGICAL HERITAGE AND LAND-USE PLANNING | 90 |
| GEOPARK AND GEOTURISM | 130 |
| COOPERATION AND EDUCATION | 180 |
| FIELDTRIP GUIDES | 210 |

In copertina: Il promontorio di Capo Colonna Area calanchiva di Aliano

Mapping regional geodiversity in Brazil and Portugal

DIAMANTINO PEREIRA (1), LEONARDO SANTOS(2), JULIANA SILVA(3), PAULO PEREIRA(1), JOSÉ BRILHA(1), JÚLIO SILVA,(2) AND CLEIDE RODRIGUES(3)

(1) Earth Sciences Centre, University of Minho and Geology Centre of the University of Porto, Portugal, insuad@dct.uminho.pt

(2) Geography Department, Federal University of Paraná (Brazil) (3) Geography Department, University of São Paulo (Brazil)

KEYWORDS: GEODIVERSITY, ASSESSMENT, MAP, PARANÁ, XINGU, BRAZIL, PORTUGAL.

ABSTRACT

A methodology meant to be used in the quantitative assessment and mapping of geodiversity was defined for regional scale, following the initial proposal of Pereira et al. (2012). The method was tested in the Xingu Basin, Amazon, Brazil (about 510,000 km²), Paraná State, Brazil (about 200,000 km²), and Portugal mainland (about 89,000 km²). It is a GIS method intended to assess all features of geodiversity and to avoid overrating any particular one, such as lithology or relief, which is a common weakness in other methods. The procedure consists on the overlay of a grid over different types of maps at scales between 1:250 000 and 1:1 000 000. The number of geological units (stratigraphical and lithological) that occurs in each grid cell of the geological map is counted, producing a map of geological indexes. The geomorphological index map results from the sum of the relief and hydrographical indexes obtained from the geomorphological units map. Palaeontological and pedological index maps are obtained from counting palaeontological units and soil units, respectively. The singular occurrences index map is based on the number of occurrences such as precious stones and metals, energy and industrial minerals, mineral waters and springs. The final Geodiversity Map results from the combination of those five partial indexes. The Geodiversity Map is a GIS automatically generated map, which allows an easy interpretation by non specialists. The map can be used as a tool in land-use planning, particularly for the identification of priority areas for conservation, and for the use and management of natural resources.

INTRODUCTION

The concept of geodiversity is quite recent and considered by most experts as "the natural range (diversity) of geological (rocks, minerals, fossils), geomorphological (landforms, processes) and soil features. It includes their assemblages, relationships,

properties, interpretations and systems" (Gray 2004). Usually, geodiversity is considered only as a theoretical approach with no particular use or application and is frequently associated with geological heritage and geoconservation issues (e.g. Alexandrowicz & Kozłowski 1999; Carcavilla et al. 2008; Gray 2004, 2008a, 2008b). Nevertheless, these concepts should not be misinterpreted as being one and the same. Whereas geodiversity refers to all abiotic variety of nature, geological heritage is the set of the most relevant geodiversity elements with particular importance for science, education or tourism (Pereira et al., 2012). During recent years, some attempts were made in order to give geodiversity a more practical approach. For instance, the Brazilian Geological Survey (CPRM) published the Geodiversity Map of Brazil (at 1:2,500,000 scale; CPRM, 2006), which is a synthesis of the major geosystems that constitute the national territory, as well as their limitations and potential uses (Silva, 2008). However, the CPRM geodiversity map is based only on lithostratigraphical and mineral resources databases and does not take into account other geodiversity elements such as landforms, soils, and hydrography, which are also important to support decision-making and land-use management (Pereira et al., 2012). In what concerns geodiversity assessment, the most promising methods are based on the definition and calculation of geodiversity indexes. However, most of them (e.g. Serrano and Ruiz-Flaño 2007; Jackova and Romportl 2008; Benito-Calvo et al. 2009; Zwolinski 2009; Hjort and Luoto 2010) do not consider the whole range of geodiversity elements. Pereira et al. (2012) developed a first approach for the calculation of geodiversity indexes by assessing all geodiversity components and to avoid overrating any particular component, such as lithology or relief. A geodiversity map based on the calculation of a geodiversity index and the outline of isolines was also produced by the authors. This type of map is a good planning tool and allows an easy interpretation by those with little or even no geological background. The state of Paraná (Southern Brazil) with an area of

about 200,000 km² has a set of different cartographical data and for this reason was used in the work here presented as a first test for the methodology proposed. GIS software was used for counting the geodiversity occurrences and the indexes calculation on the Xingu River Basin, Amazon, Brazil, an area of about 510,000 km². The Geodiversity Map of the Xingu Basin consists of a GIS automatically generated polygon map.

METHODOLOGY

The above proposed method is based on the overlay of a grid over different types of maps, such as geological, geomorphological, and soil maps. The Geodiversity Map is an isolines map obtained from the calculation of a Geodiversity Index for each cell of the grid. Isolines join the central points of cells sharing the same geodiversity index (Pereira et al., 2012). Other thematic maps, such as the geological diversity map or the geomorphological diversity map can also be produced in a similar way. Maps at scales ranging from 1/1,000,000 to 1/250,000 were used. The grid gives raise to cells where units and occurrences can be counted and which allow the discrimination of results. Various grid sizes were tested in order to obtain the best balance between results discrimination and the number of cells. The best results were obtained with a grid-size of 25x25 km resulting in 371 cells for the Paraná State map. For the Xingu Basin, the GIS procedure has generated 2462 cells on a 13.8 x 13.8 km grid. For the Geodiversity Map of Portugal the grid size is still being tested. For each grid cell, the Geodiversity Index score is the sum of the following five partial indexes: i) The Geological Index is calculated by counting the number of geological units occurring in each cell of the grid, which is overlaid on the geological map. ii) The Geomorphological Index is the sum of two sub-indexes: Relief and Hydrography. The Relief Sub-index is calculated by counting units and contacts occurring in each cell of the grid overlaid on the Geomorphological Units Map, a three level classification of morphostructural units, morphosculptural units and morphosculptural sub-units (Santos et al.,

2009; Pereira et al., 2012). For this purpose, a brand new map of geomorphological units was produced for Portugal providing three 1st level units, nine 2nd level units, and 56 3rd level units. The Hydrography Sub-index is based on the assessment of stream categorisation using Strahler's method (Strahler, 1957). iii) The calculation of the Palaeontological Index follows a similar procedure to the one described for the assessment of the Geological Index: the number of different fossiliferous formations is counted in each grid cell overlaid on a geological map (Pereira et al., 2012). iv) The Pedological Index is obtained for each grid cell by counting the soil orders represented in the Map of Soils (Pereira et al., 2012). v) The Singular Occurrences Index is related with geodiversity features not covered in the previous indexes. It was considered: minerals such as precious stones, precious metals, metallic minerals, and industrial minerals; geological energy sources such as coal, oil shale, natural gas, and uranium; mineral waters and springs. Each map occurrence of any of the above items scores one point for the corresponding grid cell. Repeated occurrences of the same element in the same cell are not considered (Pereira et al., 2012).

RESULTS

Taking into account the minimum and maximum values obtained for the Geodiversity Index, five Geodiversity Index classes were considered: very low (<11), low (11-15), medium (16-20), high (21-25), and very high (>25). The Geodiversity Map of Paraná State shows some hot spots of very high geodiversity (> 25) in the east, a region with strong geomorphological contrasts and a large variety of geomorphological and stratigraphical units (Pereira et al., 2012). The Geodiversity Map of Xingu Basin also highlights a hot spot of geodiversity in a region with a larger diversity of rocks, soils and relief, as well as the presence of several mineral occurrences. The preliminary results for the Geodiversity Map of Portugal highlights the western region, which has a large diversity of stratigraphical and palaeontological Mesozoic and Cenozoic units. However, the rich geological diversity of Portugal also originates high values of the Geodiversity Index in other areas of the country. The three given examples show that areas where the occurrence of igneous plutonic rocks is higher have the lower Geodiversity Index.

CONCLUSION

Geodiversity Index maps can be produced for large territories if solid geological, geomorphological and soil units mapping is available. GIS procedures can be used to speed-up the calculation of the geodiversity index and

its cartographic representation. Geodiversity can be represented as isolines or polygon maps allowing an easy interpretation by those with no or little geological background. Geodiversity maps can be used as a tool in land-use planning, particularly in identifying priority areas for conservation and the use and management of natural resources.

REFERENCES

- Alexandrowicz Z, Kozłowski S (1999) From selected geosites to geodiversity conservation - Polish example of modern framework. In *Towards the balanced management and conservation of the geological heritage in the new millennium*. Baretino D., Vallejo M., Gallego E. (Eds). Sociedad Geológica de España, Madrid, 40-44.
- Benito-Calvo A, Pérez-González A, Magri O, Meza P (2009) Assessing regional geodiversity: the Iberian Peninsula. *Earth Surface Processes and Landforms* 34(10):1433-1445.
- Carcavilla L, Durán JJ y López-Martínez J (2008) Geodiversidad: concepto y relación con el patrimonio geológico. *Geo-Temas*, 10; 1299-1303 (in Spanish with English abstract).
- CPRM. Mapa Geodiversidade do Brasil (2006) escala 1:2.500.000. Legenda expandida. Brasília: CPRM/Serviço Geológico do Brasil, CD-ROM.
- Gray M (2004) *Geodiversity: valuing and conserving abiotic nature*. John Wiley and Sons, Chichester.
- Gray M (2008a) Geodiversity: the origin and evolution of a paradigm. In: Burek C, Prosser C (eds) *The history of geoconservation*, Geological Society, London, pp 31-36.
- Gray M (2008b) Geodiversity: a new paradigm for valuing and conserving geoheritage. *Geoscience Canada* 35(2/3); 51-59.
- Hjort J, Luoto M (2010) Geodiversity of high-latitude landscapes in northern Finland. *Geomorphology* 115(1-2):109-116.
- Jačková K, Romportl D (2008) The relationship between geodiversity and habitat richness in Sumava National Park and Kivoklátsko Pla (Czech Republic): a quantitative analysis approach. *Journal of Landscape Ecology* 1(1):23-38.
- Pereira D., Santos L., Pereira P. & Brilha J. (2012) Geodiversity assessment in Paraná State (Brazil). *Environmental Earth Sciences* (submitted).
- Santos LC, Oka-Fiori C, Canali N, Fiori A, Silveira C, Silva J (2009) Morphostructural Mapping of Paraná State, Brazil. *Journal of Maps*, 2009:170-178.
- Serrano E, Ruiz-Flaño P, Arroyo P (2009) Geodiversity assessment in a rural landscape: Tiermes-Caracena area (Soria, Spain). *Memorie Descrittive Della Carta Geologica d'Italia* 87:173-180.
- Silva CR (ed) (2008) *Geodiversidade do Brasil. Conhecer o passado, para entender o presente e prever o futuro*. CPRM, Geological Survey of Brazil, Rio de Janeiro (in Portuguese).
- Strahler AN (1957) Quantitative analysis of watershed geomorphology. *Transactions of the American Geophysical Union* 8(6):913-920.
- Zwoliński Z (2009) The routine of landform geodiversity map design for the Polish Carpathian Mts. *Landform Analysis* 11:77-85.